

**State of California  
Natural Resources Agency  
Department of Fish and Wildlife  
Wildlife Branch**

**Status and Distribution of the Light-footed Ridgway's  
(Clapper) Rail in California  
2015 Season**

**By**

**Richard Zembal, Susan M. Hoffman, and John Konecny**

# **Final Report**

To

State of California  
Department of Fish and Wildlife  
South Coast Region  
3883 Ruffin Road  
San Diego, CA 92123

## **Status and Distribution of the Light-footed Ridgway's (Clapper) Rail in California**

**2015 Season**

**Richard Zembal, Susan M. Hoffman, and John Konecny**

Clapper Rail Recovery Fund  
Huntington Beach Wetlands Conservancy  
24821 Buckboard Lane  
Laguna Hills, CA 92653

8 October 2015

**State of California  
The Resources Agency  
Department of Fish and Wildlife**

**Status and Distribution of the Light-footed Ridgway's  
(Clapper) Rail in California**

**2015 Season<sup>1</sup>**

by

Richard Zembal, Susan M. Hoffman, and John Konecny  
Clapper Rail Recovery Fund  
Huntington Beach Wetlands Conservancy  
24821 Buckboard Lane  
Laguna Hills, CA 92653

**ABSTRACT**

The thirty-sixth annual census of the Light-footed Clapper Rail in California was conducted from 23 February to 30 May 2015. Thirty coastal wetlands were surveyed by assessing call counts from Carpinteria Marsh in Santa Barbara County, south to Tijuana Marsh National Wildlife Refuge (NWR) on the Mexican border. This rail has been reclassified taxonomically and renamed by the American Ornithologist Union, ascribed to the Light-footed Ridgway's Rail, *Rallus obsoletus levipes*; formal wildlife agency recognition of this nomenclatural change is pending.

For the fourth year in a row the California population of the Light-footed Ridgway's Rail exceeded 500 breeding pairs and for the first time in 40 years exceeded 600 pairs. A total of 633 pairs exhibited breeding behavior in 22 marshes in 2015. This is the highest count on record, representing an increase of 105 pairs from the breeding population detected in 2014, and 43% larger than the former high count in 2007. The tally at Upper Newport Bay was the highest ever recorded at 234 pairs. The Newport subpopulation was once again the largest in California with 5.4% more rails exhibiting breeding behavior than in 2014, surpassing for the third year in a row and by 34.5% the long standing record of 174 pairs manifested in 2005. The calling in Tijuana Marsh NWR was again mediocre, yielding 98 breeding pairs, a 30.7% increase over 2014 but perhaps not completely reflective of the abundance of rails therein. Based on call counts, nest searches in Newport in 2014 and 2015 yielded much lower nesting activity and in Tijuana in 2014 yielded much higher nesting activity. The Newport subpopulation comprised 37% of the state population in 2015 and the subpopulation in the Tijuana Marsh NWR comprised a minimum of 15.5%, together accounting for 52.5% of the breeding population of this rail in

---

<sup>1</sup> Zembal, R., S.M. Hoffman, and John Konecny. 2015. Status and Distribution of the Light-footed Ridgway's (Clapper) Rail in California, 2015. California Department of Fish and Wildlife, Wildlife Branch, Nongame Wildlife Program Report, 2015-04. Sacramento, CA 26 pp.

California as compared to 56.2 % in 2014 and 56.4% in 2013.

Eleven of the small subpopulations increased in size from the 2014 totals, by a combined total of 82 breeding pairs in 2015. The Seal Beach National Wildlife Refuge (NWR) subpopulation was up by 17 to 66 pairs, ranking it the third largest in state. In response to the development of cordgrass cover that is unsurpassed in the range of this rail, the San Elijo Lagoon subpopulation doubled to 60 pairs and fourth largest in California. Batiquitos Lagoon was back up to its record high of 45 breeding pairs, ranking as the fifth largest subpopulation in 2015. The University of California's Kendall-Frost Reserve achieved a record high of 33 breeding pairs in 2015, a testament to strong management of this small Mission Bay remnant marsh. Nesting activity in Point Mugu was dismal in 2015 and it was back to 2010 levels of only 12 pairs after increasing steadily to a record high of 23 pairs in 2013. Declines totaling only 14 pairs were noted in 3 marshes including San Dieguito Lagoon (-8 pairs), Mugu Lagoon (-4 pairs), and San Luis Rey River (-2 pairs). Excluding the 2 largest subpopulations, there were 9 subpopulations in double figures, ranging from 10 to 66 pairs and totaling 264 breeding pairs or 41.7% of the state total. The remaining 11 subpopulations ranged from 1 to 8 pairs and totaled 37 breeding pairs, or 5.8% of the total.

The annual increases in the population total of the Light-footed Clapper Rail between 2002 and 2007 gave encouragement that restoration and management including breeding in zoological facilities were contributing to the recovery of this endangered bird. The 2008 crash was presumably weather-related and a harbinger of what could be in store if wide weather fluctuations are the future norm. Record high counts of over 500 pairs in 2012 through 2014, and of 633 pairs in 2015 is a manifestation of this subspecies' resiliency with consistent management.

## INTRODUCTION

The Light-footed Clapper Rail (*Rallus longirostris levipes*) is a state- and federally- listed endangered species that is resident in coastal wetlands in southern California and northern Baja, California, Mexico. This rail along with both of the other large rails of the western U.S. has been reclassified taxonomically and renamed by the American Ornithologist Union and ascribed to the Ridgway's Rail, *Rallus obsoletus* (Chesser et al. 2014). The common name for our southern California subspecies soon should be legally adopted by the wildlife agencies in recognition of this nomenclatural change. The Light-footed Clapper Rail will then be called the Light-footed Ridgway's Rail, *R. obsoletus levipes*.

Loss and degradation of habitat threaten the continued existence of this bird, although recent management efforts are reversing those trends. The California population of this endangered rail was at a former high of 325 pairs in 15 marshes in 1996, the largest number detected breeding since statewide annual surveys were begun in 1980, until 2004 when 350 pairs were detected in 15 marshes. Since then, there were annual increases until the record high in 2007, when 443 breeding pairs were detected in 19 marshes. There was a population crash in 2008 followed by recovery of 37% in 2009 to 320 breeding pairs, and annual increases since then through 2014 when a new high total of 528 pairs was reached.

One of the first major investigations of this rail identified the lack of suitable nesting habitat as a major, widespread limiting factor (Massey and Zembal 1980). Subsequent work demonstrated the need for emergency actions and recommended management strategies to stem the alarming population decline of this endangered bird in southern California. The actions taken have included: 1) habitat restoration, particularly through enhancement of tidal action to former wetlands; 2) study and control of introduced predators and unnaturally high predator populations; 3) provision of nesting sites in marshes with good habitat but limited options for protected nesting locations; 4) studies that have led to adaptive management strategies, benefiting the rail and the other co-inhabitants of these biologically-rich ecosystems; 5) development of a protocol for captive breeding and genetic and demographic augmentation of smaller subpopulations; and 6) surveys of the California population, in part to track the effects of management on annual recruitment.

Implementation of these measures has succeeded in protecting and maintaining the small subpopulations and in supporting the expansion of many of them, particularly because of the release of captive bred rails. However, the benefits of the associated habitat restoration and management go far beyond this single species. These endangered birds thrive in our most productive, remaining coastal wetlands. Measures that benefit this rail and its environs enhance conditions for a myriad of other species as well, including people. These places and the wildlife are cherished by hundreds of thousands of southern Californians for their inherent aesthetic, recreational, economic, scientific, educational, and ecological values. Furthermore, there are essential links between the coastal wetlands and vast acres of diverse upland habitats and wildlife located many miles from the coast (Soule et al. 1988, Zembal 1993). Restoring and maintaining the diversity and vital productivity of the coastal wetlands, while achieving the recovery of the Light-footed Ridgway's Rail, may only be possible in an environment that includes coastal southern California's complete wildlife heritage, fostered by a caring public who support the management necessary to maintain the interconnectedness and viability of the system.

Hundreds of wetland acres have undergone, or are being planned for restoration. However, full recovery and functionality of a coastal wetland may take decades to achieve. In the meantime, habitat suitability for the rail may be quite marginal. All but a few of the current subpopulations of Light-footed Ridgway's Rails depend upon a marginal habitat base and are too small to be expected to maintain themselves without management, particularly population augmentation.

Population monitoring is essential in understanding the effects of our management efforts and in stewardship of this critically endangered bird toward recovery. Reported herein are the results of the 2015 statewide survey of the Light-footed Ridgway's Rail.

## **METHODS**

The thirty-sixth consecutive annual census of Light-footed Ridgway's Rails in California was conducted from February 23 through May 30, 2015. Thirty coastal wetlands were surveyed by mapping territorial pairs based on their calls (Zembal and Massey 1981, 1985; Zembal 1992).

All of the coastal marshes with known or suspected rail subpopulations were surveyed until an evening or early morning with good calling activity was encountered. Small wetlands with no recent rail sightings that again yielded negative results were surveyed at least twice as were marsh parcels with lower than expected results on the first call count. Additionally, nesting data were considered in the assessment of the subpopulations inhabiting the four wetlands wherein such data were gathered in 2015 (Mugu, Seal Beach, Kendall-Frost, and Sweetwater) and a pre-nesting high tide count was accomplished on November 7, 2014 on the Seal Beach NWR; a post-nesting high tide count will be scheduled for fall/early winter 2015. This NWR is the only wetland inhabited by Ridgway's rails that is inundated enough during a 6.7 ft. tide or higher to get a relatively complete visual survey.

In Upper Newport Bay and Tijuana Marsh, mapping spontaneous calls was the prevalent technique. In marshes with fewer rails and along long, narrow strips of habitat, playbacks of taped "dueting" were used sparingly to elicit responses. In the Tijuana Marsh NWR, enough observers were stationed within potential hearing range of any calling rail to cover the entire marsh on a single evening. However, most of the marshes were surveyed by a single observer visiting discrete patches of habitat on consecutive evenings until all available habitat had been covered. Most of the observations were those of three observers, but primarily the principal investigator. Additional observers participated primarily in three of the year 2015 counts, those at Seal Beach NWR, Tijuana Slough NWR and Kendall-Frost Reserve.

The more movement required of an observer during a survey, the more likely that breeding, but infrequently-calling rails would be missed. Calling frequency and the detection of calls are influenced by the observer's hearing ability and experience with the calls, the stage of breeding of individual pairs, rail density, and weather conditions (Zembal and Massey 1987). Many surveys attempted on stormy, windy days needed to be repeated. When calling frequency is high with many rounds of calling as adjacent pairs respond to one another, it is possible to map the rails accurately and move on to survey more marsh. However, under usual circumstances approximately 20 ha (50 acres) of marsh can be adequately covered during a single survey.

Surveys are usually conducted in the 2 hrs before dark, but some are done from first light to about 2 hrs after sunrise. In the past, early morning and late evening surveys have been comparable, although evening calling by the rails is more intense and often ends with one or more flurries of intense calling (Zembal et al. 1989).

The playback of a taped "clapping" call appears to be responded to by the rails as if a living pair is calling nearby. However, work done with Yuma Ridgway's Rails (*Rallus obsoletus yumanensis*) strongly suggests that this closely-related species can become conditioned to the tape if it is used excessively (B. Eddleman, pers. comm., July 1993). During prime calling times in the evening or early morning, a playback sometimes elicits a single response or a round of calling. However, there are sometimes no vocal responses to the tape. If played at a time of day when the rails are not particularly prone to call, the only response likely to be elicited is that of the territorial pair intruded upon. Sometimes the response is non-vocal investigation by the pair or one member. Repeated playbacks are likely to elicit aggression. When used only once per

year at a given marsh and with minimal repetition, playbacks have yielded important results. Unmated Ridgway's rails, for example, often respond at considerable distances and may approach the tape. Isolated single rails often approach very closely and remain in the vicinity unless displaced.

In assessing the rail population, duets and some single "clapperings" were treated as territories. Since advertising singles are not indicative of an occupied territory with reproductive potential at the time of the survey, they are not included in the population total. However, a single "clapping" is as good an indicator of a territory as a duet, when advertising is not heard later from the same territory. Eventually, during a 2 – 4 hr census period, pairs often dueted from territories where only single pair members had called earlier. However, the fewer rails in a marsh, the more important it is to count only duets as pairs to avoid over-estimating the breeding subpopulation. The 2015 call counts were conducted on 41 dates and totaled approximately 412 field-hours, mostly from February 23 – May 30, 2015.

## Study Areas

Descriptions of all the marshes recently occupied by Light-footed Ridgway's Rails are available (U.S. Fish and Wildlife Service 1985 and Zembal and Massey 1981). Four of the current principal study areas are at the Naval Air Station Point Mugu (NASPM, also Point Mugu), the Seal Beach NWR, Upper Newport Bay Ecological Reserve, and Tijuana Slough NWR.

The marsh at Point Mugu is located in southeastern Ventura County on the 1,821 ha (4,500 acre) Naval Base Ventura County (NBVC), about 13 km (8 miles) west of the Los Angeles County line. There are 1,012 ha (2,500 acres) of jurisdictional wetlands in Point Mugu (USACOE/EPA 1994), including the largest functioning salt marsh in coastal southern California today. Considering the combined acreages of marshes that are regularly occupied, the vegetated marsh and most closely associated habitats at Mugu Lagoon represent more than 25% of the Ridgway's Rail's potential habitat base. The marsh is subject to nearly full tidal action in the central and eastern arms with a tidal amplitude of about 9 ft. The tides are dampened by constrictions at Laguna Road and farther west, resulting in an amplitude of only 4 - 5 ft. The wetland vegetation is dominated by pickleweed (*Salicornia virginica*) but scattered stands of spiny rush (*Juncus acutus* ssp. *leopoldii*) are critical for rail nest placement.

The Seal Beach NWR covers 369 ha (911 acres) of the 2,024 ha (5,000 acre) Seal Beach Naval Weapons Station in Orange County near the City of Seal Beach. About 299 ha (739 acres) of the refuge lands are subject to regular inundation by the tides. There are about 229 ha (565 acres) of salt marsh vegetation, 24 ha (60 acres) of mudflats that are exposed daily, and 46 ha (114 acres) of channel and open water. The wetlands are fully tidal, with a range of about - 0.5 m (1.7 ft) to + 2.2 m (7.2 ft) MLLW, and very productive with a high diversity and abundance of wildlife.

Upper Newport Bay is an Ecological Reserve of the California Department of Fish and Wildlife (CDFW), located approximately 22 km (13.7 mi) down coast of the Seal Beach NWR. Approximately 304 ha (750 acres) are fully tidal, including 105 ha (260 acres) of marsh. The bay

is bordered by bluffs, 9 - 18 m (30 - 59 ft) high, and surrounded by houses and roads. There are approximately 100 ha (247 acres) of shrublands remaining undeveloped on the edge of the wetlands and two local drainages, with some cover along them coursing into the bay.

Tijuana Slough NWR consists of 427 ha (1,056 acres) of open water, tidal salt marsh, beach dune, riparian, and maritime scrub habitats in the City of Imperial Beach in the extreme southwest corner of the U.S. The NWR is part of the 1,024 ha (2,530 acre) Tijuana River National Estuarine Research Reserve (NERR), one of only 26 such NERRs in the country. The fully tidal coastal salt marsh (influenced by a 7 ft tide MLLW) comprises 159 ha (392 ac) of the total area along with 41 ha (101 ac) of tidal creeks and mudflat. Tijuana Slough is the only coastal wetland in the southern California Bight that is not bisected or greatly impacted by a major paved road or the coast railroad.

## **RESULTS and DISCUSSION**

A total 633 pairs of Light-footed Ridgway's Rails exhibited breeding behavior in 22 marshes in 2015 (Table 1). This is the highest count on record, representing a 105 pair increase over the breeding population detected in 2014 (Zemba et al. 2014), and 43% larger than the former high count of 2007 that stood as the high for 31 years until 2012. Upper Newport Bay with 234 pairs was once again the largest subpopulation in California and at a record high for the third consecutive year with 5.4% more rails exhibiting breeding behavior than in 2014, 22.5% more than in 2013, and 34.5% higher than the long standing former high count of 174 in 2005.

Following three years of more than 100 pairs, the Tijuana Marsh NWR subpopulation count was 75 pairs in 2014 and back up to 98 pairs in 2015. The 2015 count was a 30.6% increase over 2014 but 31% lower than the record high of 142 pairs in 2007. The Newport subpopulation comprised 37% of the state population in 2015 and the subpopulation in the Tijuana Marsh comprised 15.5%, together accounting for 52.5% of the breeding population of this rail in California. In addition, nine subpopulations ranged in size from 10 to 66 pairs, totaling 264 breeding pairs or 41.7% of the state total. These nine included: Mugu Lagoon; Seal Beach NWR; Huntington Beach Wetlands; Buena Vista Lagoon; Batiquitos Lagoon; San Elijo Lagoon; San Dieguito Lagoon; Kendall-Frost Reserve; and the San Diego River. The remaining 11 subpopulations ranged from 1 to 8 pairs and totaled 37 breeding pairs (5.8% of the state total).

Breeding subpopulations in 2015 were male-skewed at 11 marshes and female-skewed at one marsh. One of the 11 male-skewed marshes also had three advertising females, a situation that was likely short lived. A minimum of 76 unmated males and four females were heard during the call counts including: two advertising males at Point Mugu; 24 single males at Seal Beach; two males at Bolsa Chica; two males at Huntington Beach Wetlands; 15 males at Upper Newport; one male in San Joaquin Reserve; three males in Buena Vista Lagoon; nine males in San Elijo Lagoon; eight males in San Dieguito River Valley; one female in Los Penasquitos Lagoon; two males at Kendall-Frost; and eight males and three females in Tijuana Marsh. The usual condition has been a slight male bias during most years in most marshes. An extreme male skew or a slight female skew could indicate major issues, possibly due to heavy depredation.



Table 1. Census of the Light-footed Ridgway's Rail in California, 1980-2015.

Part I: 1980 - 1989

Location	Number of Pairs Detected In:									
	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989
Santa Barbara County										
Goleta Slough	0	0	-	0	-	-	-	-	0	0
Carpinteria Marsh	16	14	20	18	26	7	4	5#	2#	0
Ventura County										
Ventura River Mouth	-	-	0	0	-	-	-	-	-	0
Santa Clara River Mouth	-	-	0	-	-	-	-	-	-	0
Mugu Lagoon	-	0	-	1	3	7	6	7#	7#	5
Los Angeles County										
Whittier Narrows Marsh	-	-	-	*	0	-	-	-	-	0
Orange County										
Seal Beach NWR	30	19	28	20	24	11	5	7	14	6#
Bolsa Chica	0	0	0	0	-	-	-	*	0	0*
Huntington Beach Wetlands	-	0	-	-	-	-	0	0	0	0
Upper Newport Bay	98	66	103	112	112	87	99	119	116	116
San Joaquin Reserve	-	-	5	4	1	2	1	0	0	0
Carlson Rd Marsh	-	-	5	4	2	0	0	1#	0	0
San Diego County										
San Mateo Creek Mouth	-	-	0	0	-	-	0	-	0	0
Las Pulgas Canyon Mouth	-	-	0	0	0	-	-	-	-	0
Las Flores Marsh	-	-	0	0	0	-	0	-	0	0
French Canyon Mouth	-	-	-	0	0	-	-	-	-	0
Cocklebur Canyon Mouth	-	-	1	0	0	-	-	0	0	0
Santa Margarita Lagoon	0	0	2	1	2	1	1	1	1	0
San Luis Rey River Mouth	-	-	0	0	-	-	0	0	0	0
Guajome Lake Marsh	-	-	0	1	2	0	0	0	0	0
Buena Vista Lagoon	0	0	0	*	0	-	-	-	0	0
Agua Hedionda Lagoon	1	2	1	7	6	1	0	0	0	0
Batiquitos Lagoon	0	0	0	0	0	-	-	-	-	0
San Elijo Lagoon	-	5a	4	4	10	1	0	2	5#	7#
San Dieguito Lagoon	-	-	-	-	-	-	-	*	0	0
Los Penasquitos Lagoon	-	0	-	0	0	-	0	-	1a#	0
Kendall-Frost Reserve	18	16	6	20	24	17	12	6a#	4a#	4#
San Diego River	-	3	1	2	2	1	0	0	1a#	0#
Paradise Creek Marsh	1	2	3	1	1	0	0	0	0	0
Sweetwater Marsh	4	5	7	6	14	3	9	5a#	5	5#
E Street Marsh	3	1	3	3	2	2	2	0a	1#	0
F Street Marsh	-	1	1	0	1	0	0	0	0	0
J Street Marsh	-	1	0	0	-	-	0	0	0	0
Otay River Mouth	3	4	5	3	5	1	1	0	0	0
South Bay Marine Reserve	3	3	1	1	2	1	1a	2#	5	5#
Dairymart Ponds	-	-	-	-	-	-	0	*	1a	0#
Tijuana Marsh NWR	26	31	25	41	38	0	2	23a#	14a#	15a#
Total: pairs	203	173	221	249	277	142	143	178	177	163
marshes	11	15	18	18	19	14	12	11	14	8

Table 1. Census of the Light-footed Ridgway's Rail in California, 1980-2015.  
(continued) Part II: 1990 - 1999

Location	Number of Pairs Detected In:									
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Santa Barbara County										
Goleta Slough	0	0	0	0	-	-	0	0	-	-
Carpinteria Marsh	0	0	0	0#	0	2#	3#	5#	3#	2#
Ventura County										
Ventura River Mouth	0	0	0	0	0	0	0	-	0	-
Santa Clara River Mouth	0	0	0	0	0	0	0	-	0	-
Mugu Lagoon	6#	4#	5#	5	6#	5#	3#	4#	4#	4#
Los Angeles County										
Whittier Narrows Marsh	-	-	-	0	0	-	0	0	-	-
Orange County										
Seal Beach NWR	16	28	36	65	66	51#	52#	37#	16#	15#
Bolsa Chica	0#	0*	0#	0#	0*	0*	0*	0*	0*	0
Huntington Beach Wetlands	0	0	0	0	0	0	0	0	0	-
Upper Newport Bay	131	128	136	142	129	117	158	149#	105#	104#
San Joaquin Reserve	0	0	0#	0	0	0	0	0	-	0
Carlson Rd Marsh	0	0	0	0	0	0	0	0	-	0#?
San Diego County										
San Mateo Creek Mouth	0	0	0	0	0	0	0	-	-	-
Las Flores Marsh	0	0	0	0	0	0	0	-	-	-
Cocklebur Canyon Mouth	0	0	0	0	0	0	0	0	0	0
Santa Margarita Lagoon	0	0	0	0#	0	0	0	0#	0	0
San Luis Rey River Mouth	0#	0	1	0	-	0	0	0	0	0
Guajome Lake Marsh	0	0	0	0	-	0	0	0	-	-
Buena Vista Lagoon	0a#	2#	5	2#	3#	1#	6#	7#	4	5#
Agua Hedionda Lagoon	0	0	0	0	0	0	0	1?	1	0
Batiquitos Lagoon	0#	0#	0	1#	1#	0#	2	2	1	3
San Elijo Lagoon	5#	5	4#	6#	1#	3#	3#	8	3#	5#
San Dieguito Lagoon	0	0	0	0	0	0	0	0	0	-
Los Penasquitos Lagoon	0	0#	0#	0#	1	1	1	2	2#	2
Kendall-Frost Reserve	5#	9	11	5#	5#	4#	1#	2	2	4#
San Diego River	2	5	1a	5	5#	6b	5	5#	4	3
Paradise Creek Marsh	0	0	1a	0a	0	1	2	0	0	0
Sweetwater Marsh	2#	4a	4a	3a	7#	7	8	3#	4	3
E Street Marsh	0	1a	1a	1	0#	2	1	1	1	2
F Street Marsh	0	0	0	0	0	0	0	0	1	0
J Street Marsh	0	0	0	0	0	0	0	0	0	0
Otay River Mouth	0	0	0	0	0	1	3	3	2	1
South Bay Marine Reserve	5	2	3a	1	0	0	0	1#	1	0
Dairymart Ponds	0a#	0#?	0#	1a	0	-	-	-	-	-
Tijuana Marsh NWR	17a#	47a	67a	63a	64	61	77	77#	68#	80#
Total: pairs	189	235	275	300	288	262	325	307	222	233
marshes	9	11	13	13	11	14	15	16	17	14

Table 1. Census of the Light-footed Ridgway's Rail in California, 1980 - 2015.  
(continued) Part III: 2000 - 2010.

Location	Number of Pairs Detected In:										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Santa Barbara County											
Goleta Slough	-	0	0	0	-	-	-	-	0	0	0
Carpinteria Marsh	1#	1#	2	0#	0#	0	0	0	0	0	0
Ventura County											
Ventura River Mouth	-	-	0	0	-	-	-	-	0	-	-
Santa Clara River Mouth	-	-	0	0	-	-	-	-	0	-	-
Mugu Lagoon	7#	7#	10#	14#	19#	14#	17#	15#	5#	9#	12#
Los Angeles County											
Whittier Narrows Marsh	-	-	0	-	-	-	-	0	-	0	0
Orange County											
Seal Beach NWR	10#	11#	24#	23#	16#	15#	21#	24#	17#	19#	25
Bolsa Chica	0	0	0*	0	0	0	*	*	*	*	1
Huntington Beach Wetlands	-	0	0	0	0	0	4#	4	1#	5#	6#
Upper Newport Bay	150#	124#	129#	144#	165#	174#	158#	165#	88#	148#	131#
San Joaquin Reserve	0	0	0	0	-	0	0	0	*	0	#
Carlson Rd Marsh	0#	0	0	0	-	0	0	0	0	0	0
San Diego County											
San Mateo Creek Mouth	0	0	0	0	0	-	-	-	0	-	-
Las Flores Marsh	0	0	0	0	0	-	-	-	0	-	-
Cocklebur Canyon Mouth	0	0	0	0	0	-	-	-	0	-	-
Santa Margarita Lagoon	0	0	1	2	1	2	1	1	1#	-	-
San Luis Rey River Mouth	0	0	0	0	0	0	0	0	0	0	2#
Guajome Lake Marsh	0	-	-	0	-	-	0	0	0	-	-
Buena Vista Lagoon	5#	3#	6#	5#	5#	6#	8#	8#	9#	9#	6
Agua Hedionda Lagoon	2	2	1	4	5	4#	7#	4	7	6	2#
Batiquitos Lagoon	2#	3#	3#	5	11	16#	19#	22	22	26#	36#
San Elijo Lagoon	1#	1#	2	7#	7#	6#	15#	12#	5#	8	15#
San Dieguito Lagoon	0#	0#	0	0#	6	12#	31#	15#	21#	12#	28#
Los Penasquitos Lagoon	1	1	2	1#	2#	2	7#	12#	2#	4#	9#
Kendall-Frost Reserve	4	4	5#	6#	14	14	5#	4#	2#	7	10#
San Diego River	3#	4	6	6#	8#	5	4	6	4#	3	7#
Paradise Creek Marsh	0	0	0	0	0	0	0	0	0	-	0
Sweetwater Marsh	2	3#	3#	1#	3#	1	4#	4#	3	5	6#
E Street Marsh	2	0	1	1	0	0	2	1	0	0	2
F Street Marsh	0	0	0	0	0	0	0	0	0	0	0
J Street Marsh	1	0	0	1	0	0	0	0	0	0	0
Otay River Mouth	1	1	1	0	0	1	2	1	0	1	1
South Bay Marine Reserve	0	0	0	0	0	0	1	2	0	1	1
Dairymart Ponds	-	-	-	2	1	1	0	1	-	0	0
Tijuana Marsh NWR	61#	52#	78#	64#	87	87#	102#	142#	47#	57#	76#
Total: pairs	253	217	274	286	350	360	408	443	234	320	376
marshes	16	14	16	16	15	16	18	19	15	16	19

Table 1. Census of the Light-footed Ridgway's Rail in California, 1980 - 2015.  
(continued) Part IV: 2011 - 2015.

Location	Number of Pairs Detected In:				
	2011	2012	2013	2014	2015
Santa Barbara County					
Goleta Slough	-	0	0	0	0
Carpinteria Marsh	0	0	0	0	0
Ventura County					
Ventura River Mouth	-	0	0	0	0
Santa Clara River Mouth	-	0	0	0	0
Mugu Lagoon	16#	22#	23	16	12 2K
Los Angeles County					
Whittier Narrows Marsh	-	-	-	-	-
Orange County					
Seal Beach NWR	34#	42#	40	49	66 24K
Bolsa Chica	*	*	1	2	7 2K
Huntington Beach Wetlands	6#	6	7	9	12 2K
Upper Newport Bay	137#	165#	191	222	234 15K
San Joaquin Reserve	2#	1#	2	1	1 1K
Carlson Rd Marsh	0	0	0	0	0
San Diego County					
San Onofre Creek Mouth	0	-	1	-	0
Las Flores Marsh	0	-	0	-	0
Cocklebur Canyon Mouth	0	-	0	-	0
Santa Margarita Lagoon	2	0	0	-	3
San Luis Rey River	3	3	4	5	3
Guajome Lake Marsh	-	-	-	-	0
Buena Vista Lagoon	3#	9#	2	4	10 3K
Agua Hedionda Lagoon	7	9	8	6	8
Batiquitos Lagoon	43#	43#	45	40	45
San Elijo Lagoon	15#	31#	20	30	60 9K
San Dieguito Lagoon	12#	45#	37	23	15 8K
Los Penasquitos Lagoon	12#	11#	12	5	5 1KB
Kendall-Frost Reserve	19	16#	8	23	33 2K
San Diego River	6#	6#	10	9	11
Paradise Creek Marsh	0	0	0	0	0
Sweetwater Marsh	7#	4#	4	4	5
E Street Marsh	1	1	1	1	1
F Street Marsh	0	0	0	0	0
J Street Marsh	1	1	1	1	1
Otay River Mouth	1	1	1	1	1
South Bay Marine Reserve	1	3	2	2	2
Dairymart Ponds	-	0	-	0	-
Tijuana Marsh NWR	113#	101#	105	75	98 8K 3KB
Total: pairs	441	520	525	528	633
marshes	21	20	22	21	22

- indicates that no census was taken.

\* indicates a fall or winter occurrence.

# indicates the detection of unpaired rails (used beginning in 1987).

K = Kecking by advertising male; KB = keck-burr by advertising female.

a Paul Jorgensen Unpublished data; b 2 pairs are in Famosa Slough.

## **Four Principal Study Areas; Largest Subpopulations and/or Habitat Acreage**

### Upper Newport Bay

The 2015 Newport count was taken on 10 dates in February and March during this fourth year of drought and the rails were responding to neighboring calls in enthusiastic fashion, yielding the highest count ever tallied in a single wetland since these surveys began in 1979. This was followed later in the spring by nest searching in order to re-verify the one to one relationship documented in the 1980s between calling and nesting; a pair's duet means there is, or shortly thereafter, will be a nesting attempt in close proximity. Nest searching in Upper Newport Bay had been very unfruitful in recent years compared to earlier years. Four to six nests were found annually 2009 – 2013 with 60 field-hours or more of effort but few egg nests were discovered before hatching (none in 2013) and several had been depredated by raccoons, *Procyon lotor*. Apparently, the super abundance of raccoons (there are tracks and sign everywhere well out into the marsh) has been spawned in part by the illegal release of rehabilitated raccoons into Upper Newport Bay. Nest searches were done mostly by RZ in support of and with assistance from Michelle Barton, a graduate student at California State University Long Beach who is examining nest site characteristics. In 2014 we found a total of 8 nesting sites on Upper Island in about 30 hours of searching over eight dates; we expected to find 18 nesting sites, based upon calling. The cordgrass on Upper Island was stunted; tall enough patches to support a rail nest were very scarce in 2014 and 2015. Even the bulrush cover was mediocre and late sprouting probably due to the lack of rain and runoff. With the threat of raccoon depredation, adequate nest cover in flotsam, wrack, or on high marsh berms, places that have been used in the past, were not used in 2014 or 2015. It currently appears that tidal wash threatens cordgrass nests with highs as low as 6.3 ft MLLW.

Big winter rail survival in recent years may be one positive aspect of having no winter storm flows ripping up the marsh. With so little nesting activity on Upper Island we searched additional areas and found nest numbers closer to the expectations yielded by earlier call counts in those same areas. Over three dates in May and early June we found 12 nests, four with eggs and four that had been predated. Nesting rails in Upper Newport Bay are being increasingly affected by tidal wash with limited alternatives because of predator abundance, particularly raccoons. It is time to consider deployment of nesting rafts in Upper Newport Bay and the institution of predator management.

### Tijuana Marsh

Tijuana Marsh's subpopulation was 87 pairs for two consecutive years prior to the 2006 count of 102 breeding pairs, then the record count of 142 pairs in 2007. That 40-pair increase in 2007 was unprecedented at any marsh except Upper Newport Bay; the 95 pair crash in 2008 was totally unprecedented. This subpopulation had not been that small since 1991. Rail numbers increased annually to more than 100 pairs by 2011 – 2013 and probably stayed near there since. The 2014 call count was shown to be low by comparing nest search results; calling was not great in 2015 either. The Tijuana Marsh subpopulation came back from a crash in 1985, suffering the effects of ocean inlet closure. Today, this is the second most secure and resilient subpopulation in California. Four of the pairs detected in 2011 - 2015 were in the restored "Model Marsh" south of the river where nesting was confirmed in 2010 with the discovery of several hatched egg nests.

Observed differences between habitat conditions and nest placement in the Tijuana Marsh and Upper Newport Bay in recent years have been compelling. Most of the nests in Tijuana Marsh were in cordgrass but many also were in rank pickleweed, isolated out in the marsh with little indication of tidal over-wash. There was also little evidence of predation or raccoon presence and encounters with adult rails, adults displaying in defense of chicks, and with young of the year were very common in Tijuana Marsh but rare in Upper Newport Bay.

### Seal Beach

The Seal Beach NWR subpopulation was 20 pairs or slightly more for most of the 2000s until 2011 and 2012 when 34 and 42 pairs bred there, respectively. The 2012 total was the highest since 1996 and the fifth highest count ever. The 2013 total was just two pairs short of the 2012 counts with 40 pairs tallied combining the nesting rafts and call count data. The 2014 total of 49 pairs was the new high count since 1996 and the 66 pairs documented in 2015 ties for the highest count on record, last observed back in 1994. Evening call count results have generally been poor and we have had to rely upon nesting data obtained through monthly visits to the nesting rafts, upon which most of this subpopulation nests.

With ample habitat available to the rails on the Seal Beach NWR and strong monitoring and management programs in place, a large, resilient breeding population has been expected to develop. Raptor predation is suspected to be limiting rail survival in part because the marsh cover has been short; the cover, particularly of cordgrass has improved over time. Ongoing raptor monitoring has documented very high raptor numbers particularly in winter; seasonal high tide counts of rails and raptors have also continued. Another rail kill by a Red-tailed Hawk, *Buteo jamaicensis*, was observed in 2015; the Red-tail was nesting on the edge of the NWR in a Eucalyptus apparently left for such purpose. The most recently observed rail kill by a Northern Harrier (*Circus cyaneus*) in the NWR was in 2013, witnessed by Kirk Gilligan. Many more raptor kills than observed undoubtedly occur.

Seal Beach is one of few marshes currently occupied by Light-footed Ridgway's Rails that gets fully inundated during a high tide of about 6.7 ft (MLLW), or higher. Tides of this height occur regularly in the late summer, usually in darkness, and in the fall and winter in the early morning. The rails are forced onto debris or to the edge of the marsh where there is little cover and busy roads just beyond. This greatly exposes the rails to potential predation and vehicle collision. For example, a juvenile rail was found dead on the edge of PCH just off the NWR in September 2015 by Bob Schallman. However, the completeness of inundation also allows fairly dependable surveying of the subpopulation outside of the breeding season. Accordingly, the rails were counted again from canoes after the 2014 breeding season, but before the 2015 breeding season; the post-breeding high tide count will be done in early winter 2015. The pre-nesting count was on 7 November 2014 and 102 rails were counted (Table 2).

Table 2. High Tide and Call Counts of Ridgway's Rails on the Seal Beach National Wildlife Refuge, 1975 - 2014.

Date	Tidal Height	Ridgway's Breeding Pair			Notes
		Rails Counted	Members Before	Members After	
2 Dec 1975	7.0	22	-	-	
31 Dec 1975	6.7	12	-	-	
21 Nov 1976	7.1	24	-	-	
20 Dec 1976	7.1	35	-	-	
21 Dec 1976	7.0	34	-	-	
10 Dec 1977	7.1	16	-	-	
11 Dec 1977	7.1	40	-	-	
18 Jun 1978	6.8	16	-	42	+6 youngsters
30 Nov 1978	6.7	38	-	42	
1 Dec 1978	6.7	32	-	42	
3 Sep 1979	6.4	20	42	60	Tide too low
3 Nov 1979	6.6	56	42	60	
2 Dec 1979	6.7	32	42	60	
3 Dec 1979	6.7	44	42	60	
21 Nov 1980	6.9	55	60	38	First red fox den found
29 Jun 1981	7.0	34	60	38	Tide too late, dark
12 Nov 1981	6.9	43	38	56	
29 Dec 1982	7.0	23	56	40	
18 Jan 1984	6.9	23	40	48	
21 Nov 1984	6.7	5	48	22	+ 7 red foxes
13 Nov 1985	7.1	2	22	10	+ 2 red foxes
12 Dec 1985	7.2	2	22	10	+ 2 red foxes
30 Dec 1986	7.2	7	10	14	Begin red fox trapping, 59 foxes removed in 1986
28 Jan 1987	7.0	7	10	14	63 red foxes removed in 1987
8 Aug 1987	7.3	8	14	14	Tide too late, dark
22 Nov 1987	6.7	12	14	28	
21 Dec 1987	7.0	8	14	28	+ 2 red foxes
16 Feb 1988	6.8	10	14	28	
22 Nov 1988	6.9	6	28	12	128 red foxes removed in '88
16 Oct 1989	6.9	59	12	32	Record High Tide Count; 25 red foxes removed in 1989
5 Oct 1990	6.4	57	32	56	Tide too low
2 Nov 1990	6.8	69	32	56	Record High Tide Count
22 Nov 1991	6.9	98	56	72	Highest Population Total
26 Oct 1992	6.8	159	72	130	Highest Population Total
15 Oct 1993	6.8	143	130	132	Highest Population Total Ever
4 Nov 1994	7.0	150	132	102	220 Red-tailed Hawks counted On the NWS on 11 December 1994
25 Oct 1995	6.5	53	102	104	Tide too low
22 Nov 1995	6.9	55	102	104	
10 Dec 1996	6.7	55	104	74	
17 Oct 1997	6.6	40	74	32	
04 Nov 1998	6.8	30	32	30	

Table 2 (continued). High Tide and Call Counts of Ridgway's Rails on the Seal Beach National Wildlife Refuge, 1975 - 2014.

Date	Tidal Height	Ridgway's Rails Counted	Breeding Pair Members		Notes
			Before	After	
23 Nov 1999	7.0	17	30	20	
11 Dec 2000	6.9	30	20	22	
15 Nov 2001	6.7	35	22	48	
04 Dec 2002	7.1	62	48	46	
26 Oct 2003	6.7	96	46	32	
12 Nov 2004	6.7	52	32	30	
15 Nov 2005	6.7	57	30	42	
09 Oct 2006	6.6	103	42	48	
06 Nov 2006	7.0	95	42	48	
26 Oct 2007	7.1	32	48	34	
12 Nov 2008	6.9	20	34	38	
01 Dec 2009	6.8	50	38	50	
05 Nov 2010	7.0	51	50	68	
26 Oct 2011	6.9	96	68	84	
14 Nov 2012	7.1	145	84	80	
04 Nov 2013	6.7	121	80	98	
07 Nov 2014	6.8	102	98	132	

The pre-nesting high tide count of 102 rails in 2014 was only the seventh highest on record. Since 132 paired rails and 24 males made it into the breeding season, at least 54 rails hid without being counted during the high tide survey. Potential rail predators were out in abundance during the winter count, hunting the marsh and edges, including Red-tailed Hawks (*Buteo jamaicensis*), Northern Harriers, Peregrine Falcon (*Falco peregrinus*), Cooper's hawk (*Accipiter cooperi*), American kestrels (*Falco sparverius*), and Short-eared Owls (*Asio flammeus*). Continued upgrading and maintenance of the artificial rafts on the Seal Beach NWR is essential to the protection of the wintering rails and success of the breeding rails. More than half of the rails observed during winter high-tide counts are sequestered on the rafts.

### Point Mugu

Since doubling in size between 2001 and 2003, the Point Mugu subpopulation fluctuated between 14 and 19 pairs, from 2003 - 2007. It had been much smaller, 3 - 7 pairs for nearly 20 years until augmentations with captive-bred rails fostered its growth. There was a crash in 2008 back to 5 pairs, but the subpopulation was back up to 9 pairs in 2009, 12 pairs in 2010, a minimum of 16 pairs in 2011, 22 pairs in 2012, an all-time high of 23 pairs in 2013, but down to 16 pairs in 2014 and 12 pairs in 2015. There is an efficient predator management program in place, consistent rail and marsh management, but issues, perhaps mostly raptor predation, prevent this subpopulation from exploding into full occupation of the largest contiguous patch of potential habitat in southern California. There was no activity detected in the eastern arm/central lagoon and two pairs attempted to breed in freshwater marsh vegetation on the west side along Perimeter Road. Raptor depredation appears to have been a long standing issue in Mugu. Consequently, the rails depend upon the heavy cover provided by spiny rush (*Juncus acutus leopoldii*) but many of the



spiny rush stands are greatly degraded by competing vegetation that should be weeded out of these stands. In addition, the freshwater marsh dewatered in dry years and could be kept viable through the entire breeding season with flood irrigation if possible there.

There were regular re-sightings of banded rails at Point Mugu up until 2008 when captive-bred rails were no longer released there. Although some of the captive-bred rails appeared to have stayed in Mugu, some definitely left after release. In 2008, for example, Martin Ruane re-sighted a banded rail four days after its release on August 22 near the release site. However, at least one banded rail, a female banded 1035-8878, did not stay at Point Mugu. A photograph was taken of this rail at Upper Newport Bay on December 12, 2004 by Steve Metz. This female was bred at the Chula Vista Nature Center and released into the eastern arm of Point Mugu on August 28, 2004, 106 days before her picture was taken at Newport. This shattered the old long-distance movement of 13.5 miles recorded for the subspecies *levipes* (Zembal et al. 1983). The distance from Point Mugu to Upper Newport Bay is approximately 90 miles along the coast. The long distance record, 160 miles was traveled by a female banded 1065-39863 (“Amelia”), released at Point Mugu August 25, 2009 and recaptured November 4, 2010 at the Chula Vista Nature Center (now Living Coast Discovery Center). Amelia had returned to the facility where she was hatched and reared.

### **Eighteen Additional Study Areas, North to South**

#### **Carpinteria Marsh**

The last known Ridgway’s Rail call from Carpinteria Marsh was from an unmated female vocalizing constantly with no answering call in 2003. In 2004, there was total silence until April 13, when two males were released in the hope that the female was still alive. Occasional reports of Ridgway’s Rail vocalizations have been investigated in 2005 through 2015 but could not be corroborated. This northern wetland is plagued with domestic cats in the marsh and other predators of concern, most notably red fox. At least one red fox den location is known on the very edge of the marsh. Without consistent predator management, the chances for the reoccurrence of a viable subpopulation in Carpinteria Marsh are poor.

#### **Bolsa Chica**

Attempts to elicit responses to a tape-playback of a duet were unsuccessful at Bolsa Chica in 2011 and 2012, when only males were detected. However, Ridgway’s Rail breeding behavior was observed definitively in Bolsa Chica in 2010, 2013, 2014, and particularly 2015 when a record seven pairs were documented. The rails on the up-coast side of the boardwalk were extremely visible foraging, nest building, feeding young, and throughout the season. Kelly O’Reilly sent along multiple photos of the first brood there of seven chicks. They may be the most observed and photographed family of rails ever. There were three pairs of rails south and four pairs north of the boardwalk including one on Rabbit Island. The habitat is close enough to PCH, to be of major concern for collision hazard with vehicles (see 2015 recovery of a dead rail from PCH at Seal Beach NWR). The near constant noise masks predator cues and the fast moving vehicles would dispatch any rail that flushed that way. Recent reports of rails vocalizing from south Bolsa Chica below the bluffs in the freshwater reed stands again could not be corroborated.

### Huntington Beach Wetlands

One of the highlights of the 2006 survey of Light-footed Ridgway's Rails was the discovery of yet another breeding location in the Santa Ana River Marsh, also previously known as Newport Slough and listed in Table 1 under the Huntington Beach Wetlands (HBW). Four pairs were detected there in 2006 and 2007, up to six pairs in 2010 – 2012, a record nine pairs in 2014 (again including one pair in the Brookhurst Marsh), and the new high of 12 pairs in 2015. The Santa Ana Marsh is at the southern terminus of the Huntington Beach Wetland Complex, comprised of several wetland patches strung along the coast totaling more than 200 acres. The 92-acre Santa Ana Marsh was restored as part of the Federal Flood Control Project on the Santa Ana River. Dampened tidal influence was re-established and cordgrass was planted primarily along a narrow eastern portion of the marsh that lies between an oil field and the south dike of the river. This cordgrass marsh is extremely well-developed and patches have grown into the main marsh as well. Although the main marsh area is heavily impacted by human residents and their dogs from just across the main channel, one of the pairs detected in most years called from the largest patch of cordgrass in the center of the main marsh.

Restoration of the Huntington Beach Wetlands is continuing and one of the pairs counted in the tally for this marsh complex was actually in the Brookhurst Marsh in 2010. Lena Hyashi reported a pair on April 19, 2010 vocalizing and observed along the larger stand of Spiny Rush near the dunes and Pacific Coast Highway (PCH). This was the first record for Ridgway's Rails potentially breeding in the HBW Complex outside the Santa Ana River Marsh since the 1970s. Unfortunately, late in the 2010 season and in 2011 we were only able to elicit "kecking" from a male, so breeding was not confirmed. However, a pair was back again in the Brookhurst Marsh in 2012 – 2015 with a second pair in the Talbert Marsh in 2015.

### San Joaquin Reserve

Ridgway's Rail vocalizations were reported for the San Joaquin Reserve in 2010 - 2015. Nancy Kenyon reported and Jeff Bray photographed an adult there on 30 March 2015. The calling reported in the Reserve was likely an unmated male in 2010 but in 2011 breeding was documented by Barry Nerhus. A 9-egg nest was found in the southwest corner of cell 6 in bulrush in April; it subsequently hatched and chicks were observed. At least two pairs bred in the Reserve in 2011 and 2013 along with advertising males; one pair was detected in 2012; a pair and advertising female were detected in 2014; and a pair and at least one male were detected in 2015.

With increased management for edge foraging habitat, this extensive freshwater marsh system has good future potential for rails, marauding raccoons notwithstanding.

### Santa Margarita River

The salt marsh at the mouth of the Santa Margarita River typically held a single pair of nesting rails for many years and occasionally there have been two. These pairs are invariably in the same spots from year to year; at the river mouth in freshwater marsh in the Sweetwater Marsh section of the estuary and/or between Stuart Mesa Road and the railroad tracks on the north side of the river in the freshwater marsh that rims a pond. Unusually, in 2008 a single pair was located on the channel surrounding the least tern island at the junction of the inlet channel. We did not gain access to do surveys in 2009 or 2010 but did a base-wide survey of the potential habitat on base

in 2011. Once again, John Konecny found two nesting pairs in the Sweetwater Marsh section of the river mouth and nothing in the many little pocket wetlands scattered along the Pendleton coast. The Sweetwater Marsh Complex was checked once by Barry Nerhus in 2012 with negative results. Tom Ryan checked the Pendleton coast in 2013 and reported three points of calling to the state. Two points were south of the river along the little channels in the vicinity of the tern island and were described as a “purr” which must mean two advertising females; the third rail apparently uttered a single clapping at the mouth of San Onofre Creek. Access was again not gained in 2014. In 2015, Tom Ryan reported two points of calling from the usual places and John Konecny’s assistant found a pair inland of the 5 freeway.

### San Luis Rey River

Historic detections of Ridgway’s Rails on the San Luis Rey River have been rare and mostly confined to the freshwater marsh at the river mouth in Oceanside. Past reports of inland sightings could not be corroborated until recently when John Konecny found 2 pairs defending inland freshwater marsh habitat in 2010, three pairs in 2011 and 2012 (RZ), four pairs in 2013, a record five pairs and a male in 2014 and three pairs in 2015. The freshwater marsh is being invaded by willows and will probably not survive many more years unless the hydrology changes with higher flows.

### Buena Vista Lagoon

As in San Dieguito, rail numbers in the freshwater marsh habitat of Buena Vista Lagoon have fluctuated widely over the years. The high count was nine pairs in 2008, 2009, and 2012 but only two pairs in 2013, four pairs in 2014, and now a new high of 10 pairs in 2015. Two pairs were detected in the western lagoon between the railroad tracks and the coast route; three pairs were detected in the big central lagoon out from the Nature Center; four pairs were detected in the eastern lagoon; and a pair was detected on the creek. The habitat on the creek comes and goes with flood control maintenance and high winter flows. There had been a 4-acre fire in the marsh adjacent to the interpretive center in 2013 but the vegetation recovered; the entire wetland abounds with extremely abundant raccoon sign. The many management issues at this little marsh are shared with most of the other coastal wetlands including abundant non-native trees and shrubs that harbor perching predators and homeless people. In order to potentially bolster the subpopulation in this freshwater system, there was a release of 15 rails bred in zoological facilities on July 19, 2011 into the central lagoon. No releases have been allowed since then until the planned restoration is completed or the California Department of Fish and Wildlife supports a release of rails carrying telemetry or satellite tags.

### Agua Hedionda Lagoon

The marsh at Agua Hedionda Lagoon previously held a maximum of seven pairs in 1983 followed by a development project that greatly affected the drainage that used to support the brackish marsh and rails were barely detected in the 1990s. The numbers built slowly but fluctuated widely to a former high again of seven pairs in 2006, 2008, and 2011. The count hit the all-time high of nine pairs in 2012 and was just under that in 2013 and 2015 with eight pairs. With the recently increased street runoff from adjacent housing, the main freshwater marsh has rejuvenated to some extent, perhaps to the benefit of the rails as evidenced by the record number

in 2012. More recently, the drought has taken a toll on the habitat. The salt marsh on the eastern rim of the inner lagoon has developed some nice cordgrass stands from which four of the rail pairs were calling in 2015. Unfortunately four of those pairs are being regularly impacted by paddle boarders, beach goers, and their dogs accessing the marsh off the sand spit at Bayshore Drive. This subpopulation was augmented with the release of five rails from the breeding program in 2004, six in 2011, 16 in 2012, and nine in 2013 on the inland edge of the inner lagoon.

### Batiquitos Lagoon

The rails increased gradually in Batiquitos Lagoon as the wetland habitat continued to improve over time following the major restoration project implemented there by December 1996. The lagoon has remained mostly tidal and rail habitat has been generally increasing and improving. Breeding rails were detected on the north side of the lagoon for the first time in 2004 and a total of 11 pairs was detected. Ridgway's Rail numbers grew to 22 pairs in 2007 and 2008 and Batiquitos Lagoon was the third largest subpopulation in the state 2008 - 2010. New annual high counts continued into 2011 and 2012 with 43 pairs detected each year and a new recorded high of 45 pairs was documented in 2013 and now again in 2015.

We covered this marsh over multiple visits, usually with two observers using duet playback. The use of multiple observers on a single count was attempted in 2013, mostly without playback and few pairs were documented. The multiple-observer approach to surveying is very dependent on conditions on the day of the survey and the results might be greatly enhanced with playback. In 2015, there were 6 breeding pairs vocalizing from habitat adjacent to and south of the western tern island; 12 pairs along the north edge of the inner lagoon; 23 pairs along the southern edge but with no advertising females this year; 2 pairs in the northeast corner of the middle basin just west of the freeway in the extensive reed stand there; a pair on the edge of the eastern tern island; and a pair on the creek off Levante and El Camino Real in the freshwater reeds. The cordgrass in the west basin is extensive and looks vigorous, although most of it is too submerged during higher tides to provide adequate nest cover. Batiquitos Lagoon received rails bred in the Zoological facilities in 2004 and 2005 (8 rails each year), again in 2013 (6 rails) and 2014 (12 rails).

### San Elijo Lagoon

The San Elijo Lagoon subpopulation was back up to its former record high level of 15 nesting pairs in 2010 and 2011; the former high was more than doubled in 2012 with the detection of 31 breeding pairs; was down to 20 pairs in 2013; back up to one pair shy of the record high in 2014 with 30 pairs; and doubled to 60 pairs in 2015, the record high count for this wetland and fourth largest subpopulation in California. San Elijo Lagoon has had major efforts to maintain tidal function and suitable rail habitat in the central lagoon has expanded greatly; the cordgrass there is as lush as can be found anywhere in the California range of this rail. However, the lagoon does still close off to the ocean regularly resulting in poor hydrologic conditions for variable periods of time. The area inland of the weir becomes a lake during high rainfall years. Of the total, 13 pairs were in the east basin with 6 males and including two pairs along the creek; 40 pairs were in the central basin, mostly in great cordgrass; and 7 pairs were in the west basin, also mostly in lush

cordgrass. San Elijo received an augmentation of 8 rails bred in Zoological Facilities in 2004, 5 in 2006, 4 in 2007, 16 in 2009, and 7 in 2012 mostly at the weir in the inner lagoon. One of the 2004 rails was re-sighted near the railroad tracks in the central lagoon on December 13, 2004, 6 months following release, and one of the 2006 rails was observed repeatedly over 6 months off of the Rios Avenue trail. However, there have been no reported re-sightings of live banded rails since then. A dead rail was retrieved in May of 2010 that was banded and released into San Elijo on June 16, 2009.

### San Dieguito River Valley

The subpopulation of Light-footed Ridgway's Rails discovered in the San Dieguito River Valley in 2004, inland of the lagoon and El Camino Real, was first reported at six breeding pairs and then conservatively, at 12 pairs in 2005. In 2006, there was abundant calling indicative of at least 31 breeding pairs ranking this as the third largest subpopulation that year and the largest ever reported in a freshwater marsh system. This freshwater marsh fared better than the tidal marshes in the crash year of 2008 and reached its height in 2012 at 45 pairs. The 2015 count of 15 pairs was a partial count because the golf course on the south side would not allow access. However, doing playback along the south side generally adds only about 5 pairs. The surveys here are generally high one year and low the next and counts invariably include a lot of advertising males, particularly in low pair count years. In 2011, for example, there were 12 pairs and 33 advertising males. Such an abundance of unmated males is indicative of female-skewed predation, probably suffered during egg depredation. These widely fluctuating annual totals and abundance of males indicate a general lack of stability, probably due to extreme vulnerability to predators in this type of wetland; raccoon sign is very abundant along the marsh. Usually, at least one pair is detected in habitat on pond edges in the golf course; perhaps a disincentive for allowing the survey. Additional Ridgway's Rail detections are still being reported from the San Dieguito Creek Watershed but have yet to be corroborated since they would not respond to callback. Reported locations have included Lusardi Creek, the pond at 4S Ranch Community Park on Dove Creek Road, and at 4 Gee Road just north of Camino Del Sur.

The freshwater marsh system in San Dieguito Creek above El Camino Real is enigmatic in the broad swings in rail abundance. However, it is paramount to maintain this important freshwater marsh system for the rails. When the largest rail subpopulations crashed in 2008, the one in San Dieguito went up 40%. The current hydrologic regime provides the conditions sustaining this one-of-a-kind wetland; the current hydrology needs to be understood and maintained. The invasion of non-native plants should be managed; the marsh is succeeding slowly toward a woodland. The most pervasive invader is *Tamarix* sp., occurring along with pampas grass (*Cortaderia* sp.), eucalyptus (*Eucalyptus* sp.), palms (mostly *Washingtonia* sp.), and more limited giant reed (*Arundo donax*), and castor bean (*Ricinus communis*). The tamarisk in particular provides cover, shelter, and perch sites for raccoons; it needs to be removed. The restoration area between El Camino Real and the freeway has developed good cordgrass cover that appears of suitable quality for rails except that it may be inundated too frequently.

### Los Penasquitos Marsh

Los Penasquitos Marsh is dominated by vegetation indicative of prolonged closure to the ocean, particularly pickleweed. However, freshwater influence and freshwater marsh edge are increasing and the rails currently appear to be using mostly the freshwater marsh habitat. The detection of 12 pairs was a record high for this wetland in 2007. The number plummeted to only two pairs in 2008, but built back gradually to 12 pairs by 2011 and 2013 (four of which were on the creek above the lagoon in both years), but much lower in 2014 and 2015 with five pairs including one on the creek. In most years but particularly wet ones like 2011, the lagoon fills with runoff and much of the marsh remains inundated until late spring. Under these conditions, the rails do not call much and are difficult to detect until the marsh drains, later in the season; the conditions are too lake-like for breeding and foraging for a good part of the spring and early summer. With the prolonged drought over recent years, an additional problem has been extreme dryness in most areas untouched by the tides. Four rails bred at the zoological facilities were released into Los Penasquitos in 2004, four more in 2007, and nine in 2009. There was a re-sighting of a banded female hatched at the Wild Animal Park and released in 2007 at Los Penasquitos. She was photographed with her mate and three downy chicks on the edge of the pond below the San Diego Water Utilities Pump Station on Sorrento Valley Road on July 10, 2009 by Eric Kallen.

### Kendall-Frost

The annual subpopulation total in the University of California Reserve at Kendall-Frost has fluctuated widely. The rails struggled there in the 1990s with a high of 11 pairs in 1992 and mostly 2 – 5 pairs annually. In the early 2000s this subpopulation hit a low of 2 pairs in 2008, following two years of 14 pairs in 2004 and 2005. The former high count was 24 pairs back in 1984, a count nearly reclaimed in 2014 with 23 pairs but also with 16 males advertising. Now in 2015 a new all-time high of 33 pairs was recorded. This is about as densely packed as rails get in a southern California wetland. This marsh is small, very isolated, and bordered by urban housing, but it is also well managed under the University of California Reserve System. The stewardship includes appropriate predator management, habitat restoration, and research management to assure minimal human disturbance to the rails and their habitat. Additionally, nesting rafts have been provided (22 rafts in 2015) and used heavily by the rails there since 1987. There have also been translocations of eggs and adults bred in zoological facilities (5 rails in 2003, 7 in 2009, and 14 rails were released there in 2013). Additional monitoring of this remnant Mission Bay wetland is planned using winter high tide counts with the aid of the San Diego Audubon Society. A count conducted from kayaks on December 4, 2013 revealed 28 Ridgway's Rails at least two of which were banded, undoubtedly in 2013. Banded rails were observed with chicks in 2014.

### San Diego River

Cordgrass continues to dominate a significant portion of the western end of the San Diego River at the bay and a high count of 8 pairs of breeding Light-footed Ridgway's Rails were there in 2004. The numbers varied since then with seven breeding pairs detected in 2010, six pairs in 2011 and 2012, a record 10 pairs in 2013, nine pairs in 2014, and a new record of 11 pairs in 2015. One of the breeding pairs continues to be detected in little Famosa Slough, south of the 8 Freeway, where breeding was confirmed again by observers and reported by Jim Peugh; one of

the adults seen with chicks was banded. One of the pairs detected in 2010 was well west of the others, close to the ocean at the dog park. A previously unknown population of Salt Marsh Bird's Beak, *Cordylanthus maritimus maritimus*, was also discovered there in 2010 just off one of the foot trails. There were several hundred plants but unfortunately they are being smothered out by the clumped invasive Algerian Sea Lavender, *Limonium ramosissimum*. Rails bred in the zoological facilities have been released in the cordgrass marsh to potentially spawn a larger, more viable subpopulation. Five rails were released in each of three years, 2005, 2007, and 2010; 11 rails were released in 2011 including five females; and nine more were released in 2012. One of these more recently released rails was likely the banded rail observed with chicks in Famosa Slough.

The habitat in the river west of the 5 Freeway appears quite suited for rails but management may be required to reach full potential. There are large rat and ground squirrel populations inhabiting the riprap along the channel, a known drop and feeding station for bolstering the tortured lives of feral domestic cats, and a large raccoon population. We are examining the prospects of filling and vegetating the riprap with pickleweed and maritime scrub, limiting the habitat suitability for egg-eating rats and expanding native habitat. However, the river is operated in part for flood control and regular high flows in wet years could greatly affect the rails therein. Any potential project would need to be well-coordinated among many agencies.

#### Sweetwater Marsh

None of the breeding pairs of Ridgway's Rails reported for the Sweetwater Marsh NWR were inland along the Sweetwater River in 2013 or 2015. They had been detected annually for many years along the river above 2<sup>nd</sup> Street and a single pair was there again in 2014. In 2015 there were two pairs in the main marsh near the bay and another north of the Vener Pond marsh; three pairs in Vener Pond marsh; and a pair in the E Street Marsh parcel. Breeding was documented on five rafts in 2015 with signs of at least partial hatches in 10 of 11 nests; three egg clutches were found, comprised of six, six, and seven eggs, respectively. The Sweetwater Marsh Complex is endowed with a thriving raptor population, fully in evidence on every visit with ample good hunting perches spaced regularly along the marsh edge. The marsh growth is low and the rails are quite vulnerable. Four captive-bred Ridgway's Rails were released into Sweetwater in 2002, 11 in 2005, six in 2008, 14 in 2010, three in 2011, and nine in 2012 (eight of nine in Paradise Marsh) but none has been re-sighted.

#### J Street Marsh

The J Street Marsh parcel is the marsh just north of the former power plant site and salt works, dominated by cordgrass, and probably has regular presence by Ridgway's Rails but is difficult to access and survey. Single pairs were detected annually in 2011 - 2015 next to the small park at the north terminus of the marsh. This little wetland currently sports some of the most vigorous cordgrass growth in the south bay and should be a focus site for future management.

### Otay River

The Otay River is channelized, typically 100 ft wide or less where it runs under the 5 Freeway, coursing northwest for about 3,200 ft to the salt works. Most of the vegetation along this stretch is dominated by cattails with willow over-story near the freeway. The channel continues another 10,200 ft until it opens to south San Diego Bay. This latter, longer stretch is dominated by upper salt marsh plants. Single pairs of rails were detected in 2011, 2012, 2014, 2015, and in many previous years calling from the vicinity of the bike trail overcrossing of the channel just south of the salt works. No presence was detectable there in 2013 but a single clapping and a male were heard on Otay Lake on a north finger near Route 9 and Otay Lake Road. The lake is lined with a narrow fringe of reeds that may harbor more rails than detected but the habitat is narrow and marginal. Reports of rails have been annual in recent years on a developing marsh portion of the river where it flows into the salt works; the area is most easily accessible by kayak. A juvenile was reported from the newly restored western ponds in south San Diego Bay by Robert Patton on June 17, 2015. This older chick was still darkly-colored and so hatched very nearby.

### South Bay Marine Reserve

An adult Ridgway's Rail and a chick were observed in the South Bay Marine Reserve in 2005 after the survey report had already been finalized. There have been one to three pairs detected annually 2005 – 2015, except in the 2008 crash; two pairs have been reported for the past three years. This small isolated marsh is not so isolated anymore with the restoration of the two ponds to the south where a young rail was observed in 2015.

### **Additional Rail Sightings**

The continued annual release of captive-bred Ridgway's Rails is co-occurring with increased detections of rails in new locations, particularly inland sites on creeks, rivers, and lake edges. Some of the recent detections of interest are as follows. Rachel Woodfield photographed a single Ridgway's Rail at the Ballona Wetlands in August 2008. There have been repeated sightings on the edge of Point Mugu at Ormond Beach since 2009. A Ridgway's Rail was heard and observed in Bolsa Chica at the foot bridge in October 2009, bred near there in 2010, and there are annual reports of sightings since then culminating in numerous sightings in 2015 from the Bolsa foot bridge. There was also a rail reported in brackish marsh on Aera Energy property below Sea Point Avenue. Sue Hoffman flushed a single Ridgway's Rail adjacent to the mouth of the Santa Ana River in the plover yard at the Huntington State Beach California Least Tern nesting colony in 2008; a dead rail was reported between PCH and the Tern colony in July 2009. A rail was reported from the lake at Laguna Niguel in 2011. Ridgway's Rails are still reportedly vocalizing in the reeds at Kumeyaay Lake on the San Diego River including at least one advertising female in 2011. Ridgway's Rails are reported regularly in the San Dieguito River Watershed well inland of the Polo Club. Steve Brad reported a Ridgway's Rail in Encinitas Creek under the Calle Barcelona Bridge in 2011. Paul Lehman and others reported seeing a Ridgway's Rail at the northern end of Upper Otay Lake on April 20, 2009 and there have been reports there almost annually. Phil Unitt reported a dead rail retrieved on July 15, 2015 from the south side of a large building at 9791 Towne Center Drive near Los Penasquitos Lagoon. Finally, Bob Schallman recovered a dead rail from the edge of PCH near the Seal Beach NWR in September 2015.



## **Conclusion**

The Light-footed Ridgway's Rail population in California increased annually beginning in 2001, coincidentally the year of the first release of captive-bred rails into the wild, to a high count of 443 pairs in 2007 followed by the crash of 2008. The state population recovered from the crash with a 37% increase in 2009, growing annually thereafter to within two pairs of the 2007 record in 2011. In 2012 it reached a new high, for the first time exceeding 500 pairs statewide and has maintained 500+ breeding pairs annually for four consecutive years, exceeding 600 pairs in 2015 for the first time in recent record. However, many of the extant Ridgway's Rail subpopulations today remain too small for long-term viability; 15 of 22 subpopulations were 12 pairs or fewer in 2015. On the other hand, the subpopulation in Upper Newport Bay is at a record high; Tijuana Slough was near or exceeded 100 breeding pairs for at least four consecutive years recently; Batiquitos Lagoon has supported 40+ breeding pairs for five consecutive years; San Elijo Lagoon exceeded all expectations with 60 pairs; and several subpopulations are either expanding, holding, or fluctuating but at relatively high totals compared to the past, particularly in the Seal Beach NWR, and Kendall-Frost Reserve. The future outlook for the Light-footed Ridgway's Rail is stronger now than at any other former time.

## **ACKNOWLEDGEMENTS**

We thank Jim Robins, Diane Zembal, John Zembal, Martin Ruane, Charles Gailband, Brian Collins, Laurie Conrad, and Michael Mace for consistent support and participation; Michelle Barton, Mark Beatty, Josie Bennett, Mark Berger, Robin Booth, Slader Buck, Brian Collins, Lisa Cox, Jonathan Dwyer, John Fitch, Charles Gailband, Kirk Gilligan, Debbie Good, Kate Goodenough, Susan Kaveggia, Isabel Kay, Peter Knapp, Carolyn Lieberman, Kaye London, Jessica Looney, Robert Mraz, Barry Nerhus, Joy Parkes, Chris Redfern, Jim Robins, Tom Ryan, Bob Schallman, Rebecca Schwartz, Patti Smith, Dave Telford, Sharon Telford, Susan Tinel, and Christine Whitcraft for their support and participation in essential activities. Special acknowledgment goes to the staff of the Living Coast Discovery Center; Sea World, particularly Laurie Conrad; San Diego Safari Park, particularly Michael Mace; Fish and Wildlife Service; California Department of Fish and Wildlife, particularly Nancy Frost; and the Huntington Beach Wetlands Conservancy, particularly Ann McCarthy, for their contributions to the efforts for Ridgway's Rails in 2015. These activities are conducted under Master Bird Banding Permit No. 22420, Federal Fish and Wildlife Permit No. TE839480, and a Scientific Collecting Permit and Memorandum of Understanding issued by the California Department of Fish and Wildlife to Richard Zembal. Funding for this project was provided by the U. S. Fish and Wildlife Service Grant-in-Aid for threatened and endangered species program (Section 6). This report is dedicated to the memories of Loren Hays and Kristen Bender whose encouragement and insight helped keep the senior author involved in rail and wetland conservation.

## **LITERATURE CITED**

Chesser T.R., R.C. Banks, C. Cicero, J.L. Dunn, A.W. Kratter, I.J. Lovette, A.G. Navarro-Sigüenza, P.C. Rasmussen, J.V. Remsen, Jr., J.D. Rising, D.F. Stotz, and K. Winker. 2014. Fifty-Fifth Supplement to the American Ornithologists' Union *Check-list of North American*

*Birds*. The Auk: October 2014, Vol. 131, No. 4, pp. CSi-CSxv.

Massey, B.W., and R. Zembal. 1980. A comparative study of the Light-footed Clapper Rail in Anaheim Bay and Upper Newport Bay, Orange County, CA. Contract Rep., End. Spp. Office, U. S. Fish and Wildl. Serv., Sacramento, CA. 69 pp.

Massey, B.W., R. Zembal, and P.D. Jorgensen. 1984. Nesting habitat of the Light-footed Clapper Rail in southern California. *J. Field Ornithol.* 55: 67-80.

Soule, M.E., D.T. Bolger, A.C. Alberts, J. Wright, M. Sorice, and S. Hill. 1988. Reconstructed dynamics of rapid extinctions of chaparral-requiring birds in urban habitat islands. *Conservation Biology* 2(1): 75 - 92.

U. S. Fish and Wildlife Service. 1985. Recovery Plan for the Light-footed Clapper Rail. Portland, OR. 121 pp.

Zembal, R., and B. W. Massey. 1981. A census of the Light-footed Clapper Rail in California. *West. Birds* 12: 87-99.

Zembal, R., J.M. Fancher, C.S. Nordby, and R.J. Bransfield. 1983. Intermarsh movements of Light-footed Clapper Rails indicated in part through regular censusing. *California Fish and Game* 71: 164 - 171.

Zembal, R., and B.W. Massey. 1985. Distribution of the Light-footed Clapper Rail in California, 1980 - 1984. *Amer. Birds* 39: 135-137.

\_\_\_\_\_. 1987. Seasonality of vocalizations by Light-footed Clapper Rails. *J. Field Ornith.* 58: 41 – 48.

Zembal, R., B.W. Massey, and J.M. Fancher. 1989. Movements and activity patterns of the Light-footed Clapper Rail. *J. Wildl. Manage.* 53: 39 – 42.

Zembal, R. 1992. Light-footed Clapper Rail census and study, 1991. Contract Report to Calif. Dep. Fish and Game, Wildl. Manage. Div., Nongame Bird and Mammal Section Rep. 92-08. 32pp.

\_\_\_\_\_. 1993. The need for corridors between southern California's coastal wetlands and uplands, in J. E. Keeley, ed., *Interface between Ecology and Land Development in California*, Symposium proceedings, Southern California Academy of Sciences meetings at Occidental College, 1992.

Zembal, R., S.M. Hoffman, and J. Konecny. 2014. Status and distribution of the Light-footed Clapper Rail in California, 2014. CA Department of Fish and Game, Nongame Wildlife Unit Report, 2014-05. Sacramento, CA. 26 pp.