California Wildlife Habitat Relationships System California Department of Fish and Wildlife California Interagency Wildlife Task Group

SOUTHERN CALIFORNIA SLENDER SALAMANDER Family: PLETHODONTIDAE Order: CAUDATA A013 Batrachoseps major Class: AMPHIBIA

Written by: S. Morey, H. Basey Reviewed by: T. Papenfuss Edited by: R. Duke, J. Harris Updated by: CWHR Program Staff, September 2005

DISTRIBUTION, ABUNDANCE, AND SEASONALITY

What was Batrachoseps pacificus major is now recognized as its own species, Batrachoseps major. Current phylogenetic research indicates that what was Batrachoseps aridus now belongs as a subspecies of B. major, called B. major aridus. B. major aridus (endangered desert slender salamander) is found only in Hidden Palm Canyon, a tributary of Deep Canyon, and nearby Guadalupe Canyon, Riverside Co., at an elevation of 750 m (2500 ft) (Stebbins 2003). The entire population of this species may be less than 1,000 individuals. Habitats occupied include barren, palm oasis, desert wash, and desert scrub. The 57 ha (141 ac) Hidden Palm Ecological Reserve is the only known habitat of this salamander (California Department of Fish and Game 1980). B. major major and other subspecies are distributed along the southern California coast from Los Angeles Co. south to San Diego Co. and east to elevations of 1500 m (5000 ft) in the Santa Monica, San Gabriel and San Bernardino mountains (Stebbins 2003). Habitats occupied by B. major include oak woodland, chaparral, and grasslands of washes, canyons and mountain slopes.

SPECIFIC HABITAT REQUIREMENTS

Feeding: No information, but probably takes a variety of small arthropods, annelids, and molluscs.

Cover: B. major aridus occurs under limestone sheets, rocks, and talus, usually at the base of shaded, water-soaked, north- and west-facing cliff-like walls (Brame 1970). B. major found under rocks, bark and logs where there is sandy or gravelly soil, and formerly common in gardens under stones, boards and potted plants (Stebbins 2003).

Reproduction: Egg-laying occurs in fall and winter.

Water: Always associated with moist seepages.

Pattern: Prefers moist areas with rocky cover.

SPECIES LIFE HISTORY

Activity Patterns: Nocturnal surface activity during the wetter periods of winter and spring. During wetter, cooler months (late winter, early spring) B. major aridus individuals may be found primarily under rocks, but appear to retreat to moist areas underneath limestone slabs by late summer (Brame 1970).

Seasonal Movements/Migration: None.

Home Range: Unknown, but probably small (less than one acre).

Territory: Unknown.

Reproduction: Little information. Related forms lay eggs in clusters of from 4 to 21 (Stebbins 1954).

Niche: No data.

Comments: The limestone sheets utilized by B. major aridus may be critical as cover during summer. Destruction of these sheets by collectors, or other human activities, could cause the extinction of the entire species.

REFERENCES

Brame, A. H., Jr. 1970. A new species of Batrachoseps (slender salamander) from the desert of southern California. Los Angeles Co. Mus. Contrib. Sci. 200:1-11.

California Department of Fish and Game. 1980. At the crossroads. A report on the status of California's endangered and rare fish and wildlife. Sacramento. 149pp.

Stebbins, R. C. 1954. Amphibians and reptiles of western North America. McGraw-Hill, New York. 536pp.

Stebbins, R.C. 2003. Western reptiles and amphibians, 3rd edition. Houghton Mifflin, Boston. 533 pp.

A013

Life history accounts for species in the California Wildlife Habitat Relationships (CWHR) System were originally published in: Zeiner, D.C., W.F.Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. California's Wildlife. Vol. I-III. California Depart. of Fish and Game, Sacramento, California. Updates are noted in accounts that have been added or edited since original publication.