

State of California  
The Resources Agency  
DEPARTMENT OF FISH AND GAME

MONITORING OF JUVENILE ANADROMOUS SALMONID  
EMIGRATION IN THE SACRAMENTO RIVER  
NEAR HAMILTON CITY, CALIFORNIA  
JULY 2002 THROUGH JUNE 2003

by

Diane Coulon  
North Central Region

MONITORING OF JUVENILE ANADROMOUS SALMONID  
EMIGRATION IN THE SACRAMENTO RIVER  
NEAR HAMILTON CITY, CALIFORNIA  
JULY 2002 THROUGH JUNE 2003

by

Diane Coulon<sup>1</sup>

ABSTRACT

This report covers monitoring of juvenile Chinook salmon, (*Oncorhynchus tshawytscha*) trapped in an oxbow bypass channel from July 2002 through June 2003. The trap was downstream from the Glenn-Colusa Irrigation District water control structure at River Mile 205.5. The trap was operated 24 hours a day, seven days per week except during high water flow events and for trap repairs.

For the 2002/2003 sampling period, 9,195 juvenile winter-run Chinook salmon were captured at the oxbow site; 567 of those fish were adipose-fin clipped (ad-clipped). In addition, 27,290 other Chinook were captured (282 ad-clipped).

---

<sup>1</sup>

CDFG North Central Region, 2545 Zanella Way, Chico, CA 95928

## INTRODUCTION

This report is an annual compilation of monitoring done from July 2002 through June 2003 at the Glenn-Colusa Irrigation District (GCID) fish screen bypass channel site off the Sacramento River. This site has been monitored since 1991. Although all species of fish captured in the trap were recorded (Table 5), the primary focus of this study was winter-run Chinook salmon.


The winter-run Chinook salmon (WRCS), (*Oncorhynchus tshawytscha*), was listed as endangered under the Federal Endangered Species Act in January 1994, and State-listed as endangered in September 1999 under the California Endangered Species Act. Genetic research involving analysis of nuclear DNA has provided evidence that the Sacramento River winter-run Chinook salmon is distinct from other Chinook salmon in the Central Valley.

The U.S. Fish and Wildlife Service (USFWS) has a propagation program for WRCS at Livingston Stone National Fish Hatchery at the base of Shasta Dam. The program was suspended in 1996 due to concerns about potential hybridization with spring-run salmon and imprinting problems with juvenile winter-run Chinook salmon. In 1998, after considerable advances in genetic analyses to assure genetic integrity of artificially propagated WRCS, the program was re-authorized by the National Marine Fisheries Service (NMFS) and the California Department of Fish and Game (DFG).

In addition to the GCID monitoring site, there are two other monitoring sites located on the Sacramento River:

1. Red Bluff Diversion Dam, River Mile 243. Monitoring conducted year-round by the USFWS, four 8' rotary screw traps.
2. Knights Landing, River Mile 88. Monitoring conducted nine months per year by the DFG, two 8' rotary screw traps.

Figure 1.



The Sacramento River system (Figure 1) which flows through California's Central Valley, is the principal producer of Chinook salmon caught in the state's ocean fisheries. There are four distinct runs of Chinook salmon in the Sacramento River system and are distinguished as follows:

1. Fall run. These salmon are the most numerous in the Sacramento River system. They return from the ocean during June through November and spawn from September through January.
2. Late-fall run. These salmon spawn in the upper Sacramento River an upstream of Red Bluff, arriving in November through February and spawning from January through April.
3. Winter run. These salmon spawn almost exclusively in the Sacramento River and tributaries upstream of Red Bluff, spawning from July through September.
4. Spring run. These salmon return to the river from the ocean from mid-February through June and spawn from August through October.

The GCID has been diverting Sacramento River water since 1883. The GCID conveys Sacramento River water through irrigation canals to approximately 141,000 acres of agricultural land and delivers water to 20,000 acres of wildlife habitat comprising the Sacramento, Delevan and Colusa national wildlife refuges.

The GCID's Hamilton City pump station is located approximately 100 miles north of the City of Sacramento. The pump station is situated on an oxbow off the main stem of the Sacramento River at River Mile 205.5. Water flow passes through the approximately 1,100 foot length of flat plate fish screen where a portion of the water is pumped into GCID's main irrigation canal. The remaining flow in the oxbow passes by the screens and then back into the mainstem of the Sacramento River.

The GCID diverts a maximum of 3,000 cfs from the Sacramento River, with the peak demand occurring in the spring. Because GCID diverts up to 25 percent of the Sacramento River flow at Hamilton City, its pumping operations were identified as a significant impediment to the downstream migration of juvenile Chinook salmon. All four runs of salmon have declined over the past 25 years. One reason for the decline was the poor performance of an existing 20-year old drum screen. In August 1993 an interim flat-plate screen was installed across the trashrack in front of the rotary drum screen. The bypass return channel was altered to reduce the time it takes for fish to return to the river. In May of 1998, construction was begun on an approximately 520-foot extension to the existing 600-foot fish screen and three bypass structures were installed from the screens to the oxbow bypass channel. Construction was completed in May 2002.

## MATERIALS AND METHODS

### Trapping Site

Fish were trapped in an oxbow bypass channel of the Sacramento River. The site was approximately 300 feet downstream from the flat-plate fish screen structure and 100 feet downstream from the water control structure. Sampling was conducted using one 8-foot diameter rotary screw trap (RST) with a live box measuring 4'x4'x3' manufactured by EG solutions (Eugene, Oregon). Steel cable (1/4 inch diameter) connected the RST to a 1/2 inch steel high-line cable suspended across the channel and anchored on the opposite banks by secured tower structures. The trap was fished 24 hours per day, seven days a week, except during extraordinarily high water flows or during periods of excessive debris (Table 1).

### Processing Captured Fish

All fish were netted from the live box and placed into a shallow tub of river water. A subsample of approximately 50 salmon juveniles was placed in a bucket containing a solution of tricaine methanesulfonate (MS-222) and anesthetized (Nine grams of MS-222 powder are dissolved in one liter river water). Upon mobilization, the juveniles were individually placed onto a plexiglass measuring board and measured to the nearest mm fork length. Salmon race was determined using size-at-time criteria developed by Frank Fisher (DFG), Northern California, North Coast Region, unpublished data). After the subsample of 50 juvenile salmon were measured, all the juvenile salmon not anesthetized or measured were grouped and counted according to race (Table 2) and it was noted if the fish was ad-clipped (Table 3). The first ten of each non-salmonid fish species were anesthetized and measured to the nearest mm fork length, and the remainder were not anesthetized or measured but were counted. All other species captured were recorded (Table 5). After full recovery, all fish were transported downstream by boat and released into the mainstem Sacramento River at approximate River Mile 206.

### Results

This report is quantitative and does not predict future spawner populations. The data gathered in this monitoring project is used to determine trends in the numbers, timing, and presence of the Chinook salmon runs and threatened and endangered fishes.

The daily data obtained from the Glenn-Colusa fish screen bypass canal trap from January 1, 1996 to the present has been entered into the California Department of Water Resources Interagency Ecological Program's database and is available on the internet at <http://BayDelta.water.ca.gov>.

Table 1. Trap operation July 2002 through June 2003.

July 2002 – Trap in operation 25 days. No sampling conducted six days.

07/03 – GCID screen performance study.

07/10 – GCID screen performance study.

07/12 – GCID screen performance study.

07/17 – GCID screen performance study.

07/19 – GCID screen performance study.

07/25 – GCID screen performance study.

August 2002 – Trap in operation 31 days.

September 2002 – Trap in operation 30 days.

October 2002 – Trap in operation 31 days.

November 2002 – Trap in operation 30 days.

December 2002 – Trap in operation 14 days. No sampling conducted 17 days.

12/15 through 12/31 – Heavy debris, high flows.

January 2003 – Trap in operation 16 days. No sampling conducted 15 days.

01/01 through 01/07 – High flows, heavy debris, trap repairs.

01/12 through 01/16 – High flows, heavy debris.

01/23 through 01/25 – High flows, heavy debris.

February 2003 – Trap in operation 26 days. No sampling conducted two days.

02/17 through 02/18 – High flows, heavy debris.

March 2003 – Trap in operation 24 days. No sampling conducted seven days.

03/14 through 03/18 – High flows, heavy debris.

03/26 – Screen performance study.

03/28 – Screen performance study.

Table 1. Trap operation July 2002 through June 2003 (continued).

April 2003 – Trap in operation 15 days. No sampling conducted 15 days.

04/02 – Screen performance study.

04/04 – Screen performance study.

04/09 – Screen performance study.

04/11 – Screen performance study.

04/16 – Screen performance study.

04/17 – Lack of personnel.

04/18 – Screen performance study.

04/19 through 04/20 – Avoiding mortalities due to large numbers of Coleman fish and debris.

04/25 – Screen performance study.

04/26 through 04/30 – Avoiding mortalities due to large numbers of Coleman fish and debris.

May 2003 – Trap in operation 14 days. No sampling conducted 17 days.

05/01 through 05/13 – High flows, heavy debris.

05/21 – GCID screen performance study.

05/23 – GCID screen performance study.

05/28 – GCID screen performance study.

05/30 – GCID screen performance study.

June 2003 – Trap in operation 23 days. No sampling conducted 7 days.

06/04 – GCID screen performance study.

06/06 – GCID screen performance study.

06/11 – GCID screen performance study.

06/13 – GCID screen performance study.

06/18 – GCID screen performance study.

06/25 – GCID screen performance study.

06/27 – GCID screen performance study.



Table 2. Unmarked juvenile Chinook salmon recoveries at the Glenn-Colusa Irrigation District bypass channel July 2002 through June 2003. Fish were assigned to race based on size-at-time criteria developed by Frank Fisher (California Department of Fish and Game, Northern California – North Coast Region, unpublished data).

Time Period	Winter-run	Spring-run	Fall-run	Late fall-run
July 1-15	9	0	832	260
July 16-31	51	0	131	120
Aug. 1-15	260	0	73	165
Aug. 16-31	253	0	27	69
Sept. 1-15	658	0	10	29
Sept 16-30	2581	0	6	91
Oct. 1-15	3087	0	2	73
Oct. 16-31	716	13	0	32
Nov. 1-15	273	5	2	62
Nov. 16-30	609	6	0	38
Dec. 1-15	49	12	15	10
Dec. 16-31	0	0	0	0
Jan. 1-15	31	1	2542	2
Jan. 16-31	11	4	8027	0
Feb. 1-15	16	4	4484	0
Feb. 16-29	7	1	1724	0
Mar. 1-15	1	26	186	0
Mar. 16-31	16	38	360	0
Apr. 1-15	0	1162	876	3
Apr. 16-30	0	565	3786	0
May 1-15	0	0	134	1
May 16-31	0	0	599	6
June 1-15	0	0	139	1
June 16-30	0	0	254	0
TOTAL	8628	1837	24209	962

Table 3. Adipose fin-clipped juvenile Chinook salmon recoveries at the Glenn-Colusa Irrigation District bypass channel July 2002 through June 2003. Fish were assigned to race based on size-at-time criteria developed by Frank Fisher (California Department of Fish and Game, Northern California – North Coast Region, unpublished data).

Time Period	Winter-run	Spring-run	Fall-run	Late fall-run
July 1-15	0	0	0	0
July 16-31	0	0	0	0
Aug. 1-15	0	0	0	0
Aug. 16-31	0	0	0	0
Sept. 1-15	0	0	0	0
Sept 16-30	0	0	0	0
Oct. 1-15	0	0	0	0
Oct. 16-31	0	0	0	0
Nov. 1-15	0	0	0	29
Nov. 16-30	0	0	0	27
Dec. 1-15	0	0	1	13
Dec. 16-31	0	0	0	0
Jan. 1-15	47	0	0	26
Jan. 16-31	26	0	0	7
Feb. 1-15	470	46	11	0
Feb. 16-29	15	20	56	0
Mar. 1-15	3	6	12	0
Mar. 16-31	6	13	10	0
Apr. 1-15	0	5	0	0
Apr. 16-30	0	0	0	0
May 1-15	0	0	0	0
May 16-31	0	0	0	0
June 1-15	0	0	0	0
June 16-30	0	0	0	0
<b>TOTAL</b>	567	90	90	102

Figure 2. Summary of unmarked Chinook salmon recoveries in the Department of Fish and Game's rotary screw trap located at the GCID fish screen facility, July 2002 through June 2003.




Figure 3. Summary of marked Chinook salmon recoveries in the Department of Fish and Game's rotary screw trap located at the GCID fish screen facility, July 2002 through June 2003.




Table 4. GCID bypass channel other species inventory.

Acipenseridae

sturgeon (*Acipenser spp*)

Catostomidae

sucker, Sacramento (*Catostomus occidentalis*)

Centrarchidae

bass, largemouth (*Micropterus salmoides*)

bass, smallmouth (*Micropterus dolomieu*)

bluegill (*Lepomis macrochirus*)

crappie, black (*Pomoxis nigromaculatus*)

crappie, white (*Pomoxis annularis*)

sunfish, green (*Lepomis cyanellus*)

sunfish, redear (*Lepomis microlophus*)

Clupeidae

shad, American (*Alosa sapidissima*)

shad, threadfin (*Dorosoma petenense*)

Cottidae

sculpin (*Cottus spp*)

Cyprinidae

carp, common (*Cyprinus carpio*)

hardhead (*Mylopharodon conocephalus*)

hitch (*Lavinia exilicauda*)

roach, California (*Hesperoleucus symmetricus*)

shiner, golden (*Notemigonus chrysoleucas*)

splittail (*Pogonichthys macrolepidotus*)

pikeminnow, Sacramento (*Ptychocheilus grandis*)

dace, speckled (*Rhinichthys osculus*)

Embiotocidae

perch, tule (*Hysterocarpus traski*)

Gasterosteidae

stickleback, threespine (*Gasterosteus aculeatus*)

Ictaluridae

bullhead, brown (*Ictalurus nebulosus*)

catfish, channel (*Ictalurus punctatus*)

catfish, white (*Ictalurus catus*)

Petromyzontidae

Lamprey (*Lampetra spp*)

Poeciliidae

Mosquitofish, western (*Gambusia affinis*)

Salmonidae

salmon, Chinook (*Oncorhynchus tshawytscha*)

trout, rainbow (*Oncorhynchus mykiss*)

Table 5. Monthly summary of other species captured in the Department of Fish and Game's rotary screw trap located at the GCID fish screen facility, July 2002 through June 2003.

SPECIES	July	August	September	October	November	December	January	February	March	April	May	June	TOTAL
bass, largemouth	1	5	4	1	3	0	0	0	0	0	0	26	40
bass, smallmouth	0	0	0	0	7	0	2	4	0	0	0	1	14
bluegill	1	2	0	3	10	0	1	6	38	13	7	11	92
bullhead, brown	0	0	0	0	1	0	0	0	0	0	0	0	1
carp, common	0	0	1	0	1	0	0	1	0	0	0	0	3
catfish, channel	0	0	0	0	1	0	2	1	0	0	1	2	7
catfish, white	0	0	0	0	0	0	0	0	0	0	5	0	5
crappie, black	0	0	0	1	0	0	0	1	1	0	0	0	3
crappie, white	0	0	0	1	0	0	0	0	0	0	0	1	2
dace, speckled	0	0	0	0	0	0	0	0	0	0	0	0	0
fathead minnow	0	0	0	0	0	0	0	0	0	0	0	0	0
hardhead	4	17	6	2	5	0	9	18	10	9	14	20	114
hitch	0	0	0	0	0	0	0	2	0	0	0	0	2
Lamprey	39	31	24	64	42	32	3	0	1	10	29	66	341
Mosquitofish, western	3	9	5	4	2	0	0	0	10	0	0	0	33
perch, tule	2	7	3	6	0	3	1	0	2	3	2	3	32
pikeminnow, Sacramento	175	493	36	23	11	2	41	105	113	28	27	553	1607
roach, California	0	0	0	0	0	0	0	0	0	0	0	0	0
sculpin, prickly	1	3	0	1	1	1	2	23	37	16	3	6	94
sculpin, riffle	0	0	0	0	0	0	0	0	0	0	0	0	0
shad, American	0	4	0	0	0	0	0	0	0	0	0	1	5
shad, threadfin	2	0	0	2	0	1	17	6	0	0	12	1	40
shiner, golden	3	3	1	0	1	0	4	1	1	0	1	4	19
splittail	0	0	0	0	0	0	1	0	0	0	0	0	1
stickleback, threespine	0	10	7	7	3	0	2	11	20	7	1	0	68
sturgeon (unidentified)	0	0	0	0	0	0	0	0	0	0	0	4	4
sucker, Sacramento	114	101	232	79	15	7	2	5	0	0	62	53	670
sunfish, green	0	0	0	1	0	0	1	0	0	1	0	0	3
sunfish, redear	0	0	0	0	0	0	0	0	0	0	0	0	0
trout, rainbow (unmarked)	11	16	2	1	2	0	5	1	4	2	4	0	48

