

California Wildlife Habitat Relationships System
California Department of Fish and Wildlife
California Interagency Wildlife Task Group

BROWN PELICAN

Pelecanus occidentalis

Family: PELECANIDAE
B043

Order: PELECANIFORMES

Class: AVES

Written by: S. Granholm
Reviewed by: D. Raveling
Edited by: R. Duke

DISTRIBUTION, ABUNDANCE, AND SEASONALITY

The brown pelican is found in estuarine, marine subtidal, and marine pelagic waters along the California coast. Rare to uncommon on Salton Sea from July to September. Breeds on Channel Islands: Anacapa, Santa Barbara, and Santa Cruz (Garrett and Dunn 1981) from March to early August; most numerous then within 20 km (12 mi) of those islands (Briggs et al. 1981). In southern California, common along coast from June to October, especially within 30 km (19 mi) of shore, but regularly out to 175 km (109 mi) (Briggs et al. 1981). Fairly common the rest of year, with lowest numbers April to May (Garrett and Dunn 1981). In northern California, fairly common to common June to November, rare to uncommon December to February and May, and very rare and irregular March and April (Anderson and Anderson 1976, Cogswell 1977, McCaskie et al. 1979).

SPECIFIC HABITAT REQUIREMENTS

Feeding: Forages mainly in early morning or late afternoon, or when tide is rising. Feeds almost entirely on fish, caught by diving from 6-12 m (20-40 ft) in the air, and occasionally from up to 20 m (66 ft). May completely, or only partially, submerge and water may be shallow or deep. Occasionally feeds on crustaceans, carrion, and young of its own species (Palmer 1962).

Cover: Usually rests on water or inaccessible rocks (either offshore or on mainland), but also uses mudflats, sandy beaches, wharfs, and jetties. Evidently does not roost overnight on water, but concentrates at a few traditional roosts on mainland or islands (Briggs et al. 1981).

Reproduction: The nest is a small mound of sticks or debris on rocky, or low, brushy slopes of undisturbed islands (Cogswell 1977); usually on the ground, but less often in bushes (Palmer 1962).

Water: No additional data found.

Pattern: Needs undisturbed islands adjacent to good marine fishing areas.

SPECIES LIFE HISTORY

Activity Patterns: Yearlong, diurnal activity.

Seasonal Movements/Migration: After breeding, beginning as early as mid-May, individuals leave colonies in the Channel Islands and in Mexico, and disperse along the entire California coast. Small numbers visit the Salton Sea and Colorado River reservoirs. Most return to breeding colonies by March or April.

Home Range: During nesting season, most numerous within 20 km (12 mi) of nesting

islands (Briggs et al. 1981).

Territory: Nests are spaced a minimum distance "equivalent to 2 birds' necks plus bills" (Palmer 1962).

Reproduction: Present at nesting islands March to early August, and lays eggs March to April, but perhaps as late as June (Palmer 1962). Clutch size usually 3 eggs, sometimes 2; single-brooded. Incubation lasts about 4 wk. Young are altricial and tended by both parents; first fly at 9 wk, and are independent sometime later. Young breed first at about 2-3 yr.

Niche: Gulls, especially the Heermann's, frequently steal fish from pouches of brown pelicans immediately after a dive. Contamination of food supply by DDT and other chlorinated hydrocarbons reduced nesting productivity in California nearly to zero in 1969-71, from eggshell thinning and altered parental behavior. Since then, contamination has been reduced and productivity has increased, but it is too low for population stability (Anderson 1977). The California population is augmented by individuals moving north from Mexico each year. Gulls and vultures are typical nest predators, and eggs and nestlings sometimes are lost in storms (Palmer 1962).

REFERENCES

- Anderson, D. W. 1977. The status of brown pelicans at Anacapa Island in 1975. *Calif. Fish and Game* 63:4-10.
- Anderson, D. W., and I. T. Anderson. 1976. Distribution and status of brown pelicans in the California Current. *Am. Birds* 30:3-12.
- Anderson, D. W., and F. Gress. 1983. Status of a northern population of California brown pelicans. *Condor* 85:79-88.
- Anderson, D. W., J. R. Jehl, Jr., R. W. Risebrough, L. A. Woods, Jr., L. R. Deweese, and W. G. Edgecomb. 1975. Brown pelicans: improved reproduction off the southern California Coast. *Science* 190:806-808.
- Briggs, K. T., D. B. Lewis, W. B. Tyler, and G. L. Hunt, Jr. 1981. Brown pelicans in southern California: habitat use and environmental fluctuations. *Condor* 83:1-15.
- Cogswell, H. L. 1977. *Water birds of California*. Univ. California Press, Berkeley. 399pp.
- Garrett, K., and J. Dunn. 1981. *Birds of southern California*. Los Angeles Audubon Soc. 408pp.
- Jehl, J. R., Jr. 1973. Studies of a declining population of brown pelicans in northwestern Baja California. *Condor* 75:69-79.
- McCaskie, G., P. De Benedictis, R. Erickson, and J. Morlan. 1979. *Birds of northern California, an annotated field list*. 2nd ed. Golden Gate Audubon Soc., Berkeley. 84pp.
- Palmer, R. S., ed. 1962. *Handbook of North American birds*. Vol. 1. Yale University Press, New Haven, CT. 567pp. W. Collins Sons and Co., Cleveland OH. 416pp.
- Knopf, F. L. 1979. Spatial and temporal aspects of colonial nesting of white pelicans. *Condor* 81:353-363.
- Lingle, G. R., and N. F. Sloan. 1980. Food habits of white pelicans during 1976 and 1977 at Chase Lake National Wildlife Refuge, North Dakota. *Wilson Bull.* 92:123-125.
- Low, J. B., L. Kay, and D. I. Rasmussen. 1950. Recent observations on the white pelican on Gunnison Island, Great Salt Lake, Utah. *Auk* 67:345-356.
- Marshall, D. B., and L. W. Giles. 1953. Recent observations of birds of Anaho Island, Pyramid Lake, Nevada. *Condor* 55:105-116.
- McCaskie, G., P. De Benedictis, R. Erickson, and J. Morlan. 1979. *Birds of northern California, an annotated field list*. 2nd ed. Golden Gate Audubon Soc., Berkeley. 84pp.
- Palmer, R. S., ed. 1962. *Handbook of North American birds*. Vol. 1. Yale University Press, New Haven CT. 567pp.
- Remsen, J. V., Jr. 1978. Bird species of special concern in California. *Calif. Dept. of Fish and Game, Sacramento. Wildl. Manage. Admin. Rep. No. 78-1*. 54pp.
- Schaller, G. B. 1964. Breeding behavior of the white pelican at Yellowstone Lake, Wyoming. *Condor* 66:3-23.
- Sloan, N. F. 1982. Status of breeding colonies of white pelicans in the United States through

1979. *Am. Birds* 36:250-257.
Tait, I. C., F. L. Knopf, and J. L. Kennedy. 1978. White pelicans nesting at Honey Lake,
California. *West. Birds* 9:38-40.

B043

Life history accounts for species in the California Wildlife Habitat Relationships (CWHHR) System were originally published in:
Zeiner, D.C., W.F.Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. *California's Wildlife*. Vol. I-III. California Depart.
of Fish and Game, Sacramento, California. Updates are noted in accounts that have been added or edited since original publication.