

**California Wildlife Habitat Relationships System**  
**California Department of Fish and Wildlife**  
**California Interagency Wildlife Task Group**

---

AMERICAN COOT

*Fulica americana*

Family: RALLIDAE  
B149

Order: GRUIFORMES

Class: AVES

Written by: M. Rigney  
Reviewed by: L. Mewaldt  
Edited by: R. Duke

#### DISTRIBUTION, ABUNDANCE, AND SEASONALITY

The American coot is a common to abundant resident throughout most of the state below 2150 m (7100 ft). It is found in fresh, and saline emergent wetlands, wet grasslands, pastures, lacustrine, estuarine, cropland, and urban habitats. Uncommon on bay tideflats and salt marshes, except in winter months. May form huge concentrations at larger lakes, and at the Salton Sea before heavy winter freezes (Cogswell 1977, Garrett and Dunn 1981).

#### SPECIFIC HABITAT REQUIREMENTS

**Feeding:** Forages underwater on the foliage and roots of submerged aquatic plants such as filamentous algae, pondweeds, and watermilfoil (Fitzner et al. 1979). Also feeds by gleaned seeds, insects, and small fish from water, soil and vegetation. Large groups may feed along wetland edges, and in open fields.

**Cover:** Retreats to open water, or into dense wetland vegetation, when disturbed (Gullion 1953).

**Reproduction:** Nests over water in dense, emergent wetlands, preferably among bulrushes or cattails. Often nests in rice croplands. Builds in succession, a display platform, an egg nest, and a brood nest. Egg nest usually 0.6 to 0.9 m (2-3 ft) from open water (Bent 1926, Gullion 1954). Egg and brood nests are large, woven platforms usually placed on a floating foundation of dry plants.

**Water:** No additional data found.

**Pattern:** Breeds primarily in dense, fresh emergent wetland. Requires permanent, or semi-permanent, emergent vegetation, particularly cattails or bulrushes, and stable water levels during the breeding season. May be found in autumn and winter in *Salicornia* and *Spartina* associations with open water.

#### SPECIES LIFE HISTORY

**Activity Patterns:** Yearlong, diurnal activity.

**Seasonal Movements/Migration:** Population breeding in colder locations in the state migrates in September and October to areas where water remains unfrozen. Inland population may move to the coast for winter. Migratory patterns not well known.

**Home Range:** No data found.

**Territory:** Breeding territory in a California study averaged 0.43 ha (1.06 ac), and ranged from 0.22 to 0.56 ha (0.54 to 1.39 ac) (Gullion 1953). Aggressively defends territory intra-

and interspecifically (Ryder 1959, Ehrlich et al. 1988).

Reproduction: Monogamous; breeds April-September, with a peak May-June. First clutch averages 9 eggs, and second clutch 6.4 eggs. Incubation period 23-24 days, by both sexes. Precocial young tended by both parents (Gullion 1954, Fredrickson 1970). Young are independent at 5 wk.

Niche: Raccoons, hawks, owls, and eagles are common predators. Hunted regularly, but not highly prized.

## REFERENCES

- Bent, A. C. 1926. Life histories of North American marsh birds. U.S. Natl. Mus. Bull. 135. 490pp.
- Cogswell, H. L. 1977. Water birds of California. Univ. California Press, Berkeley. 399pp.
- Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1988. The birder's handbook. Simon and Schuster, New York. 785pp.
- Fitzner, R. E., E. T. Sipco, and R. G. Schrecknise. 1979. American coot nesting and feeding habits in southeastern Washington. Northwest Sci. 54:244-252.
- Fredrickson, L. H. 1970. Breeding biology of American coots in Iowa. Wilson Bull. 82:445-457.
- Garrett, K., and J. Dunn. 1981. Birds of southern California. Los Angeles Audubon Soc. 408pp.
- Gullion, G. W. 1953. Territory behavior of the American coot. Condor 55:169-186.
- Gullion, G. W. 1954. The reproductive cycle of American coots in California. Auk 71:366-412.
- Ryder, R. A. 1959. Interspecific intolerance of the American coot in Utah. Auk 76:424-442. Valley has reduced available habitat; densest populations remain in hunting clubs, wildlife areas, and along other managed water channels (Cogswell 1977).

Comments: Formerly called common gallinule (Cogswell 1977).

## REFERENCES

- Bent, A. C. 1926. Life histories of North American marsh birds. U.S. Natl. Mus. Bull. 135. 490pp.
- Cogswell, H. L. 1977. Water birds of California. Univ. California Press, Berkeley. 399pp.
- Dawson, W. L. 1923. The birds of California. 4 Vols. South Moulton Co., San Diego. 2121pp.
- Eckert, A. W., and K. E. Karalus. 1981. The wading birds of North America. Doubleday and Co., Garden City, NY. 252pp.
- Fredrickson, L. H. 1971. Common gallinule: breeding biology and development. Auk 88:914-919.
- Garrett, K., and J. Dunn. 1981. Birds of southern California. Los Angeles Audubon Soc. 408pp.
- Grinnell, J., and A. H. Miller. 1944. The distribution of the birds of California. Pac. Coast Avifauna No. 27. 608pp.
- Gullion, G. W. 1953. Territory behavior of the American coot. Condor 55:169-186.
- Harrison, C. 1978. A field guide to the nests, eggs and nestlings of North American birds. W. Collins Sons and Co., Cleveland, OH. 416pp.
- Harrison, C. J. O., ed. 1978. Bird families of the world. Harry N. Abrams, Inc., New York. 264pp.
- Johnsgard, P. A. 1975a. North American game birds of upland and shoreline. Univ. Nebraska Press, Lincoln. 183pp.
- McCaskie, G., P. De Benedictis, R. Erickson, and J. Morlan. 1979. Birds of northern California, an annotated field list. 2nd ed. Golden Gate Audubon Soc., Berkeley. 84pp.
- Ripley, S. D. 1977. Rails of the world: Monograph of the family Rallidae. D. R. Godine, Boston, MA. 406pp.
- Terres, J. K. 1980. The Audubon Society encyclopedia of North American birds. A.

Knopf, New York. 1100pp.

B149

---

Life history accounts for species in the California Wildlife Habitat Relationships (CWHR) System were originally published in: Zeiner, D.C., W.F.Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. California's Wildlife. Vol. I-III. California Department of Fish and Game, Sacramento, California. Updates are noted in accounts that have been added or edited since original publication.