

State Water Project Incidental Take Permit Risk Assessment for Delta Smelt and Longfin Smelt

Section 1: Overview

Date: 02/08/2022

Life Stages Present:

Delta Smelt (DS): Adults, sub-adults

Longfin Smelt (LFS): Adults, sub-adults, and larvae

Advice to Water Operations Management Team (WOMT):

No Advice.

Risk Assessment:

Delta Smelt: Based on distribution patterns over the past decade and rare detections, DS are unlikely to be prevalent in the south Delta. Limited detection data support DS being present in the Sacramento Deep Water Ship Channel (SDWSC), lower San Joaquin River, south Delta, Suisun Marsh and the lower Sacramento River. Two marked individuals have been collected since 2/1/2022. Turbidity is expected to remain stable and a turbidity bridge avoidance action will not be possible since the OMR Index will be equal to or less negative than -2,000 cfs. The expected less negative Old and Middle River (OMR) Index should maintain a low overall likelihood of entraining adults. Water temperatures are within the range for DS spawning.

Longfin Smelt: OMR Index is projected to be between -500 cfs and -2,000 cfs due to compliance with D-1641. X2 has moved downstream to 76 km due to low exports and increased upstream releases. The Smelt Monitoring Team (SMT) has determined that the overall risk of entrainment is low for larvae in the lower San Joaquin River and low for sub-adults, and adults based on low exports. LFS adults are present in the Delta based on detections by Chipps Island Trawl and Enhanced Delta Smelt Monitoring (EDSM). There is evidence that spawning has occurred in the south Delta due to detections by Smelt Larva Survey (SLS) 2 and Larval Entrainment Pilot Study (LEPS). SLS 2 triggered 8.4.2 on 1/31/2022 however, the SMT determined that advice was not warranted during the 2/1/2022 meeting because of low exports.

Barker Slough: SLS 2 did detect larvae at station 716. The relevant Condition of Approval (COA 8.12) is not active based on the current Sacramento River Index Forecast as of January 1, 2022, being Below Normal. If subsequent water year forecasts reflect a drier hydrology of Dry or Critical, then 8.12 will become active.

Section 1-A: Sacramento River and Confluence

Risk of entrainment into the central Delta and export facilities for DS and LFS in Sacramento River (8.1.5.2 C ii, iii, iv)

- Exposure Risk (Hydrology):
 - DS: Low. DS are expected to have made their migration in response to first flush and be holding in anticipation for temperatures conducive to spawning. Current water temperatures are within the range associated with spawning as reported in Damon et al. (2016).
 - LFS: Risk for larvae is low due to low exports and a resulting positive Qwest. See 'Routing Risk' for more information on adults and sub-adults.
- Routing Risk (Behavior and life history):
 - DS: Low. The daily average turbidity at Old River at Bacon Island (OBI) has not exceeded 12 FNU.
 - LFS: Low risk of entrainment. Larvae do not exhibit swimming behaviors that would result in volitional movement into areas with a higher risk of entrainment. Spawning migration is underway. There is potential for adult and sub-adult movement into the central Delta however, risk remains low due to low exports and X2 shifting downstream.
- Overall Entrainment Risk:
 - DS: Low.
 - LFS: Low for adults, sub-adults and larvae, due to:
 - Exports targeting Net Delta Outflow Index (NDOI) of 11,400 cfs which will result in an OMR Index no more negative than -2,000 cfs.
 - Adult salvage has been rare following the Pelagic Organism Decline.

Section 1-B: Central Delta

Risk of entrainment into the export facilities for DS and LFS in the central Delta (8.1.5.2 D iii, iv, v)

- Exposure Risk (Low, Medium, High):
 - DS: Low. DS are present in the south Delta based on the detection of a marked fish in salvage on 01/16/2022 and the detection of a marked DS in the lower San Joaquin River on 2/4/2022. However, the likelihood of adult and sub-adult DS entrainment remains low due to projected operations resulting in an OMR Index no more negative than -2000 cfs. There is a high degree of uncertainty regarding the response of cultured fish to environmental cues typically applied to wild DS.
 - LFS: Low risk for adult and sub-adult LFS entrainment. EDSM collected one sub-adult LFS in the lower San Joaquin River on 2/4/2022.
 - Low risk for larvae observed in the lower San Joaquin River by SLS 2. Qwest remains positive and exports have decreased since 02/01/2022.
 - SLS 2 triggered COA 8.4.2 on 1/31/2022 (See Discussion of Conditions of Approval) likely due to continued emergence in the South and Central Delta.

- Change in exposure from previous week: (Note: The change in risk compared to previous weeks is not required by the Incidental Take Permit [ITP]).
 - DS: No change from last week. X2 has shifted downstream compared to last week.
 - LFS: No change from last week due to low exports. The presence of LFS larvae in south and central Delta has been confirmed by recent sampling, though reduced exports minimizes the risk of entrainment.
- Reporting OMRI (Number and range of OMRI bins will vary based on anticipated hydrology and operations)
 - The SMT determined that risk of entrainment is low across the range of expected OMRI values. Projected exports limit the range of possible OMR Index values.

Section 2: Basis for Advice

The 2020 ITP ([Incidental Take Permit for Long-Term Operation of the State Water Project in the Sacramento-San Joaquin Delta 2081-2019-066-00](#)) states that advice to WOMT shall be based on the following Conditions of Approval:

List relevant Condition of Approval number and title based on species/life stage, time of year, etc.

8.3.1 Integrated Early Winter Pulse Protection. Between December 1 and January 31 each year Permittee shall reduce south Delta exports for 14 consecutive days to maintain a 14-day average OMR index no more negative than -2,000 cfs, and convene the Smelt Monitoring Team within one day of triggering the following criteria:

- Three-day running average daily flows at Freeport greater than, or equal to, 25,000 cfs, AND
- Three-day running average of daily turbidity at Freeport greater than, or equal to, 50 Nephelometric Turbidity Units (NTU), OR
- The Smelt Monitoring Team determines that real-time monitoring of abiotic and biotic factors indicates a high risk of DS migration and dispersal into areas at high risk of future entrainment.

After maintaining a 14-day average OMR index no more negative than -2,000 cfs for 14 days, Permittee shall maintain a 14-day average OMR index no more negative than -5,000 cfs, initiating the OMR Management season, until the OMR Management Season ends (Condition of Approval 8.8).

The Integrated Early Winter Pulse Protection Action may only be initiated once during the December 1 through January 31 time period each year.

8.3.3 Adult Longfin Smelt Entrainment Protection. After December 1, if an Integrated Early Winter Pulse Protection (Condition of Approval 8.3.1) has not yet initiated, Permittee shall

reduce south Delta exports to maintain a 14-day average OMR index no more negative than -5,000 cfs and initiate OMR Management (Condition of Approval 8.3) if:

- Cumulative combined LFS expanded salvage (total estimated LFS counts at the CVP and SWP salvage facilities beginning December 1 through February 28 exceeds the most recent Fall Midwater Trawl (FMWT) LFS index¹ divided by 10, OR
- Real-time monitoring of abiotic and biotic factors indicates a high risk of LFS movement into areas at high risk of future entrainment, as determined by DWR and CDFW Smelt Monitoring Team staff.

When evaluating the possibility of LFS movement into areas that may be subject to an elevated risk of entrainment, the Smelt Monitoring Team shall evaluate catch of LFS with fork length ≥ 60 mm by the Chipps Island Trawl (conducted by USFWS) as an early warning indicator for LFS migration movement into the Delta, in addition to other available survey and abiotic data. The Smelt Monitoring Team shall communicate the results of these risk assessments and advice to the WOMT (Condition of Approval 8.1.3), and operational decisions shall be made as described in Condition of Approval 8.1.4 (Collaborative Approach to Real-time Risk Assessment).

8.4.1 OMR Management for Adult Longfin Smelt. From the onset of OMR Management (Condition of Approval 8.3) through February 28, the Smelt Monitoring Team shall conduct weekly, or more often as needed, risk assessments (see Condition of Approval 8.1.5.2) and decide whether to recommend an OMR flow requirement between -5,000 cfs and -1,250 cfs to minimize entrainment and take of adult LFS. The Smelt Monitoring Team may provide advice to restrict south Delta exports for seven consecutive days to achieve a seven-day average OMR index within three risk categories:

- Low risk: OMR between -4,000 cfs to -5,000 cfs
- Medium risk: OMR between -2,500 cfs to -4,000 cfs
- High risk: OMR between -1,250 cfs to -2,500 cfs

If a risk assessment conducted by the Smelt Monitoring Team determines that a more restrictive OMR flow requirement is needed to minimize take of adult LFS, the Smelt Monitoring Team shall provide its advice to WOMT (Condition of Approval 8.1.3) and operational decisions shall be made following the process described in Condition of Approval 8.1.4 (Collaborative Approach to Real-time Risk Assessment).

¹ The Fall Midwater Trawl (FMWT) Survey annual abundance index for LFS is calculated as the sum of September through December monthly abundance indices and is typically reported at about the same date as adult salvage begins in December. The FMWT Index available beginning on December 1 each year shall be used to establish this threshold.

This Condition will terminate when a high-flow off-ramp occurs (Condition of Approval 8.4.3), or when LFS spawning has been detected in the system, as determined by the Smelt Monitoring Team, or, if there is disagreement and resolution is not reached within WOMT, as determined by CDFW. The Smelt Monitoring Team shall consider results from Additional LFS Larval Sampling (Condition of Approval 7.6.1) to inform its assessment of the start of LFS spawning. After LFS spawning has been observed, Permittee shall implement Condition of Approval 8.4.2 to minimize take of larval and juvenile LFS.

8.4.2 Larval and Juvenile Longfin Smelt Entrainment Protection. From January 1 through June 30, when a single Smelt Larva Survey (SLS) or 20 mm Survey (20 mm) sampling period exceeds one of the following thresholds:

- LFS larvae or juveniles found in four or more of the 12 SLS or 20 mm stations in the central Delta and south Delta (Stations 809, 812, 815, 901, 902, 906, 910, 912, 914, 915, 918, 919), or
- LFS catch per tow exceeds five LFS larvae or juveniles in two or more of the 12 stations in the central Delta and south Delta (Stations 809, 812, 815, 901, 902, 906, 910, 912, 914, 915, 918, 919).

Permittee shall restrict south Delta exports for seven consecutive days to maintain a seven-day average OMR index no more negative than -5,000 cfs. Permittee shall also immediately convene the Smelt Monitoring Team to conduct a risk assessment (see Condition of Approval 8.5.1.2) to assess the risk of larval and juvenile LFS entrainment into the South Delta Export Facilities, determine if an OMR flow restriction is warranted, and recommend an OMR flow limit between -1,250 and -5,000 cfs. The Smelt Monitoring Team risk assessment and operational advice shall be reviewed by the WOMT (Condition of Approval 8.1.3) via the Collaborative Real-time Decision-making process (Condition of Approval 8.1.4). Permittee shall operate to the export restriction and OMR flow target approved through Conditions of Approval 8.1.3 and 8.1.4. Each week the Smelt Monitoring Team shall convene to conduct a new risk assessment and determine whether to maintain, or off ramp from, export restrictions based on the risk to LFS, or until the DS and LFS off-ramp has been met as described in Condition of Approval 8.8 (End of OMR Management).

From January 1 through June 30, DWR and CDFW Smelt Monitoring Team staff shall conduct weekly, or more often as needed, risk assessments (see Condition of Approval 8.5.1.2) to assess the risk of larval and juvenile LFS entrainment into the South Delta Export Facilities. As a part of the risk assessment the Smelt Monitoring Team shall provide advice on the appropriate OMR flow targets to minimize LFS entrainment or entrainment risk, or both. The Smelt Monitoring Team shall provide its advice to WOMT (Condition of Approval 8.1.3) and use the Collaborative Approach to Real-time Risk Assessment process described in Condition of Approval 8.1.4 to determine if an OMR flow restriction is warranted and determine OMR flow limit between -1,250 and -5,000 cfs. The OMR flow limit shall be in place until the next risk assessment conducted by the Smelt Monitoring Team determines that it is no longer necessary to minimize

take or related impacts to LFS, or until the DS and LFS off-ramp has been met as described in Condition of Approval 8.8 (End of OMR Management).

8.4.3 High Flow Off-Ramp from Longfin Smelt OMR Restrictions. OMR management for adult, juvenile, or larval LFS as described in Conditions of Approval 8.4.1 and 8.4.2 are not required, or would cease if previously required, when river flows are (a) greater than 55,000 cfs in the Sacramento River at Rio Vista or (b) greater than 8,000 cfs in the San Joaquin River at Vernalis. If flows subsequently drop below 40,000 cfs in the Sacramento River at Rio Vista or below 5,000 cfs in the San Joaquin River at Vernalis, the OMR limit previously required as a part of Conditions of Approval 8.4.1 and 8.4.2 shall resume.

8.5.1 Turbidity Bridge Avoidance. The purpose of this Condition is to minimize the risk of entrainment of adult DS in the corridors of the Old and Middle rivers into the south Delta export facilities. This Condition is intended to avoid the formation of a turbidity bridge from the San Joaquin River shipping channel to the south Delta export facilities, which historically has been associated with elevated salvage of pre-spawning adult DS.

After the Integrated Early Winter Pulse Protection (Condition of Approval 8.1.3) or February 1 (whichever comes first), until April 1, Permittee shall manage exports to maintain daily average turbidity in Old River at Bacon Island (OBI) at a level of less than 12 NTU. If the daily average turbidity at OBI is greater than 12 NTU, Permittee shall restrict south Delta exports to achieve an OMR flow that is no more negative than -2,000 cfs until the daily average turbidity at OBI is less than 12 NTU.

If, after five consecutive days of OMR flow that is less negative than -2,000 cfs and the daily average turbidity at OBI is not less than 12 NTU, the Smelt Monitoring Team may convene to assess the risk of entrainment of DS (Condition of Approval 8.1.5.2). The Smelt Monitoring Team may provide advice to WOMT regarding changes in operations that could be conducted to minimize the risk of entrainment of DS (Condition of Approval 8.1.3). The Smelt Monitoring Team may also determine that OMR restrictions to manage turbidity are infeasible and may instead provide advice for a different OMR flow target that is between -2,000 and -5,000 cfs and is protective based on turbidity and adult DS distribution and salvage to the WOMT for consideration (Condition of Approval 8.1.3). Operational decisions shall be made following the process described in Condition of Approval 8.1.4 (Collaborative Real Time Risk Assessment).

Turbidity readings at individual sensors can generate spurious results in real time. Spurious results could be incorrectly interpreted as a turbidity bridge, when in fact the cause is a result of local conditions or sensor error. To assess whether turbidity readings at OBI are attributable to a sensor error or a localized turbidity spike, Permittee, in coordination with Reclamation, may consider and review data from other nearby locations and sources. Additional information that will be reviewed include regional visualizations of turbidity, alternative sensors, and boat-based turbidity mapping, particularly if there was evidence of a local sensor error. Permittee may bring data from these additional sources to the Smelt Monitoring Team for consideration during the development of a risk assessment to be provided to the WOMT for evaluation (Condition of Approval 8.1.3).

Permittee shall use the decision-making process described in Condition of Approval 8.1.4 (Collaborative Real-time Risk Assessment) to determine if south Delta exports may increase after five days of OMR no more negative than -2,000 cfs, or to determine that this action is not warranted due to a sensor error or localized turbidity event. Permittee shall implement this action until CDFW is in agreement that the action may be ended or modified.

8.5.2 Larval and Juvenile Delta Smelt Protection. If the five-day cumulative salvage of juvenile DS at the CVP and SWP facilities is greater than or equal to one plus the average prior three years' FMWT index (rounded down), Permittee shall restrict south Delta exports for seven consecutive days to maintain a seven-day average OMR index no more negative than -5,000 cfs. Additionally, if the five-day cumulative salvage threshold is met or exceeded, Permittee shall immediately convene the Smelt Monitoring Team to conduct a risk assessment (Condition of Approval 8.1.5.2) and determine the future risk of entrainment and take of larval and juvenile DS. The Smelt Monitoring Team may provide advice to further restrict south Delta exports to maintain a more positive OMR than -5,000 cfs. The Smelt Monitoring Team may provide advice for further restrictions within three risk categories:

- Low risk: Limit OMR between -4,000 cfs to -5,000 cfs
- Medium risk: Limit OMR between -2,500 cfs to -4,000 cfs
- High risk: Limit OMR between -1,250 cfs to -2,500 cfs

The duration and magnitude of operational advice shall be provided to the WOMT (Condition of Approval 8.1.3) and decisions shall be made following the process described in Condition of Approval 8.1.4 (Collaborative Real Time Risk Assessment). When conducting risk assessments to evaluate the risk of entrainment and take of juvenile DS, the Smelt Monitoring Team shall evaluate the following information sources, in addition to any other models or surveys they deem appropriate and those listed in Condition of Approval 8.1.5.2:

- Results from a CDFW approved DS life cycle model.
- DS recruitment levels identified by the Smelt Monitoring Team using the CDFW-approved life cycle model that links environmental conditions to recruitment, including factors related to loss as a result of entrainment such as OMR flows. In this context, recruitment is defined as the estimated number of post-larval DS in June per number of spawning adults in the prior February-March period.
- Hydrodynamic models and forecasts of entrainment informed by the EDSM or other relevant survey data to estimate the percentage of larval and juvenile DS that could be entrained.

If expanded salvage at the CVP and SWP facilities of juvenile DS exceeds 11 within a three-day period under this condition, Permittee shall restrict south Delta exports for seven consecutive days to maintain a seven-day average OMR index no more negative than -3,500 cfs. If juvenile

DS continue to be salvaged at the CVP and SWP facilities during the seven days of OMR restrictions, then Permittee shall continue restrictions and request a risk assessment by the Smelt Monitoring Team to determine if additional advice and subsequent restrictions are warranted and provide advice to WOMT (see Condition of Approval 8.1.3) and follow the decision-making process described in Condition of Approval 8.1.4.

8.12 Barker Slough Pumping Plant Longfin and Delta Smelt Protection. Permittee shall operate the BSPP to protect larval LFS from January 15 through March 31 of dry and critical water years. Permittee shall operate to protect larval DS from March 1 through June 30 of dry and critical years. If the water year type changes after January 1 to below normal, above normal or wet, this action will be suspended. If the water year type changes after January to dry or critical, Permittee shall operate according to this Condition of Approval.

From January 15 through March 31 of dry and critical water years, Permittee shall reduce the maximum seven-day average diversion rate at BSPP to less than 60 cfs when larval LFS are detected at Station 716. In addition, in its weekly meetings from January 15 through March 31, the Smelt Monitoring Team shall review LFS abundance and distribution survey data and other pertinent abiotic and biotic factors that influence the entrainment risk of larval LFS at the BSPP. When recommended by the Smelt Monitoring Team, and as approved through the decision-making processes described in Conditions of Approval 8.1.3 and 8.1.4, Permittee shall reduce the maximum seven-day average diversion rate at BSPP according to the advice provided by the Smelt Monitoring Team.

From March 1 through June 30 of dry and critical water years, Permittee shall reduce the maximum seven-day average diversion rate at BSPP to less than 60 cfs when larval DS are detected at Station 716. In addition, in its weekly meetings from March 1 through June 30, the Smelt Monitoring Team shall review DS abundance and distribution survey data and other pertinent abiotic and biotic factors that influence the entrainment risk of larval DS at the BSPP (including temperature and turbidity). When recommended by the Smelt Monitoring Team, and as approved through the decision-making processes described in Conditions of Approval 8.1.3 and 8.1.4, Permittee shall reduce the maximum seven-day average diversion rate at BSPP to less than 60 cfs.

The DS requirements described in this condition may be adjusted to align with USFWS requirements to minimize take of DS through an amendment to this ITP.

8.13 Water Year Type Definition. All references to water year type in this ITP shall be defined based on the Sacramento Valley Index unless otherwise noted.

Discussion of Conditions of Approval

Provide discussion addressing criteria for each Condition of Approval listed in "Basis for Advice" section. Refer to data below where appropriate.

COAs relevant to initiating OMR management went into effect December 1st. The Smelt Monitoring Team conducted a Risk Assessment based on COA 8.1.5.2.

8.3.1: This COA was triggered by conditions measured on 12/17/2021 when the running three-day average of daily flow and turbidity reached 27,152 cfs and 66.79 FNU respectively. Operations were reduced on 12/20/2021 targeting a 14-day average OMR index no more negative than -2,000 cfs for 14 consecutive days. After maintaining a 14-day average OMR index no more negative than -2,000 cfs for 14 days, Permittee shall maintain a 14-day average OMR index no more negative than - 5,000 cfs, initiating the OMR Management season, until the OMR Management Season ends (Condition of Approval 8.8).

8.3.3: This COA is no longer active due to the initiation of an Integrated Early Winter Pulse Protection (COA 8.3.1).

8.4.1: This COA is no longer active due to the detection of larval LFS by Smelt larva Survey (SLS).

8.4.2: This COA went into effect on 01/03/2022 following the 14-day Integrated Early Winter Pulse Protection (COA 8.3.1). SLS 1 was initially canceled due to COVID mitigation, however, the 12 south and central Delta stations listed in this COA were sampled on 1/18/2022. The resulting data triggered this COA by detection of larval LFS at more than four stations (809, 812, 815, 901, 906, and 910) and larval catch exceeded 5 fish per tow at two stations (809 and 812). Data collected by SLS 2 (See Table 3 in Attachments below) triggered this action for the second time this season on 1/31/2022. Larval LFS were detected at four of the relevant stations (809, 812, 815 and 906) and catch per tow exceeded five LFS at two stations (809 and 812). The SMT did not advise a less negative OMR Index under this COA due to reduced exports.

8.5.1: This COA went into effect on 01/03/2022 following the 14-day Integrated Early Winter Pulse Protection (COA 8.3.1). Current OBI turbidity levels are below the threshold.

8.5.2: The 2021 FMWT Annual Index for DS is zero for the fourth consecutive year. The salvage threshold is one Juvenile DS. No juvenile DS have been salvaged this water year. One cultured subadult DS (fork length = 54 mm, adipose fin clipped) was salvaged on 1/16/2022.

8.12: This COA is not active due to water year type. The current water year type forecast is Below Normal. This COA may become active as the Water Year Type forecast is updated. SLS 2 did detect larvae at station 716.

8.13: The Sacramento Valley Water Year Type Index (SVI) January forecast corresponding to the 50% probability of exceedance is 7.5 which is in the range for a Below Normal water year classification. The forecast was reported on the California Data Exchange Center (CDEC) [Water Supply Index Webpage](#), accessed on 01/11/2022.

Section 3: Hydrology and Operations Assessment of hydrologic, operational, and meteorological information. 8.1.5.2 A.

Section 3-A: Water operations conditions. 8.1.5.2.A. i

- Antecedent Actions: (e.g. Delta Cross Channel [DCC] gate closure and actions such as integrated early winter pulse protection, etc.)
 - COA 8.4.2 was triggered for a second time this season by data reported to the SMT 1/31/2022. No advice was provided.
 - OMR Management was initiated on 01/03/2021 following the 14-day Integrated Early Warning Pulse Protection action (COA 8.3.1).
 - COA 8.3.1 was triggered by conditions measured on 12/17/2021. Exports were reduced to comply with this COA on 12/20/2021 through 01/02/2021.
 - DCC gates closed on 11/30/2021.
 - The Drought barrier at False River has been notched. The notch is quite large, such that hydrodynamically no barrier is present.
- Controlling Factors: Delta outflow/X2 (D-1641)
- Water Temperature:
 - Clifton Court Forebay (CCF) Daily Average Water Temperature = NA
 - 3 Station Average = 10.05°C
- Tidal Cycle: In a weak spring tide transitioning to a neap tide next week.
- Turbidity:
 - 8.3.1 Freeport 3-day average = NA
 - 8.5.1 OBI Turbidity Daily Average = 5.56 FNU.
- Salinity: X2 = 76 km.
- Hydrologic Footprint: No new particle tracking models were requested or reviewed.

Section 3-B: Water operations outlook. 8.1.5.2.A. ii

- Outages:
 - State Water Project (SWP): None
 - Central Valley Project (CVP): None.
- Exports: Combined exports are targeting an NDOI of 11,400 cfs
 - SWP: 300 to 800 cfs
 - CVP: 1,800 cfs

Meteorological Forecast: No significant precipitation is in the forecast. Section 3-C: Projected conditions. 8.1.5.2.A. iii

- DCC Gates position: Closed 11/30/2021.
- Sacramento River flow at Freeport is 12,000 to 14,500 cfs
- San Joaquin River flow at Vernalis is 1,000 to 2,000 cfs.
- Qwest: +1,600 cfs
- Expected changes in South Delta Exports: Exports are not expected to change substantially in the next 7 days.

- NDOI: 11,800 cfs
- Upstream releases:
 - Keswick = 3,250 cfs
 - Nimbus = 2,000 cfs and may decrease to 1,750 cfs
 - Goodwin = 1,500 cfs.
 - Oroville = Increased to 3,250 cfs.

Table 1: Comparison of OMR and OMR Index (5-day and 14-day averages in this table for OMR Index and USGS gauge were reported on [SacPAS website](#), accessed 02/08/2022.

Date	Averaging Period	USGS gauges (cfs)	Index (cfs)
02/08/2022	Daily	Not Reported	-1,600 cfs
2/5/2022	5-day	-3,670 cfs	-1,330 cfs
2/4/2022	14-day	-4,040 cfs	-2,020 cfs

Distribution and Biology.

8.1.5.2.B. Assessment of biological information for Delta Smelt and Longfin Smelt

Section 4-A: Delta Smelt population status 8.1.5.2.B. i

- The last DS (adipose fin clipped) detection occurred on 02/07/2022 in the lower Sacramento River.
- EDSM: From 01/30/2022 through 02/04/2022, EDSM completed sampling at 18 sites and collected 2 marked DS in the SDWSC and one marked DS in the lower San Joaquin River. NOTE: This report contains data that were collected today (02/07/22) by EDSM, one DS (marked) was captured in the Lower Sacramento River Stratum. Sampling this week will occur Monday through Thursday (February 7- 10). See Table 1 in Attachments for details.
- Chipps: From 01/30/2022 through 02/04/2022, Chipps Island Trawl completed 50 tows and did not collect any DS.
- Salvage: No DS have been salvaged at either facility in the past seven days.
- Fall Mid-water Trawl (FMWT) Index for DS = 0
- DS life cycle model (LCM) discussion: NA
- Biological Conditions: Water temperatures are within the range conducive to spawning as reported in Damon et al. (2016).
- % of population in Delta zones: NA

Section 4-B: Longfin Smelt population status 8.1.5.2.B. ii.

- FMWT Index for LFS = 323
- Other Surveys:
- EDSM: From 01/31/2022 through 02/04/2022, EDSM sampled 18 sites and collected 6 LFS in Lower Sacramento, 5 LFS in Suisun Bay, 1 LFS in Lower San Joaquin, and 48 LFS in Suisun Marsh. See Table 1 in the Attachments for details.
- Chipps: From 01/30/2022 through 02/04/2022 Chipps Island Trawl conducted 50 tows and collected 18 LFS. See Table 2 in the Attachments for details. SLS 2 sample collection is complete. Sample processing is completed for most stations except 602, 520 and stations in the Napa River and San Pablo Bay. See Table 3 below for details.
- LEPS has completed a 24-hour sampling effort. Data will be shared upon completion of sample processing.
- Salvage: No LFS have been salvaged at either facility.

Section 4-C: Additional data sources to assess sensitivity to entrainment Delta.8.1.5.2.C & D. i

Notes:

- SMT members discussed the difference between the OMR Index range reported during last week's call and actual operations. The range reported in the WOMT notes was closer to actual operations. A wider range is reported in the weekly Operations Outlook than is reported during SMT calls.
- The SMT will review literature related to swimming ability of larval Longfin Smelt to inform movement.
- Experimental Release: 6,400 marked DS were released in the SDWSC on 02/03/2022. An additional 12,800 marked DS are scheduled to be released in Montezuma Slough this week. The last release for this season will consist of approximately 9,000 fish that will be released in the SDWSC.
- Qualitative larval sampling has begun at both facilities. Data will be reported during the Salvage update.

Literature cited:

Damon, L. J., S. B. Slater, R. D. Baxter, and R. W. Fujimura. 2016. Fecundity and reproductive potential of wild female Delta smelt in the upper San Francisco Estuary, California. *California Fish and Game* 102(4):188–210.

Attachments: Table 1: EDSM Catch Table, Figure 1: EDSM Sample Locations, Table 2: Chipps Island Catch Table, Table 3: SLS 2 Catch Table

Table 1. Delta Smelt (DSM) and Longfin Smelt (LFS) catch per station for EDSM 2022 Phase 1 Kodiak trawls, from 1/31/2022–2/7/2022. These data are preliminary and subject to change.

Water Year	Phase	Station Code	Date	# Tows	Species	Mark Type	Fork Length	Total Catch	Stratum
2022	1	22-27-SBM01	2/1/2022	4	NA	NA	NA	NA	Suisun Bay
2022	1	22-27-SBM04	2/1/2022	4	LFS	None	68	1	Suisun Bay
2022	1	22-27-SBM04	2/1/2022	4	LFS	None	74	2	Suisun Bay
2022	1	22-27-SBM04	2/1/2022	4	LFS	None	75	1	Suisun Bay
2022	1	22-27-SBM04	2/1/2022	4	LFS	None	81	1	Suisun Bay
2022	1	22-27-GB01	1/31/2022	4	LFS	None	51	1	Suisun Marsh
2022	1	22-27-GB01	1/31/2022	4	LFS	None	66	1	Suisun Marsh
2022	1	22-27-GB01	1/31/2022	4	LFS	None	69	1	Suisun Marsh
2022	1	22-27-GB01	1/31/2022	4	LFS	None	70	1	Suisun Marsh
2022	1	22-27-GB01	1/31/2022	4	LFS	None	72	1	Suisun Marsh
2022	1	22-27-GB01	1/31/2022	4	LFS	None	78	1	Suisun Marsh
2022	1	22-27-GB01	1/31/2022	4	LFS	None	82	1	Suisun Marsh
2022	1	22-27-GB01	1/31/2022	4	LFS	None	96	1	Suisun Marsh
2022	1	22-27-GB01	1/31/2022	4	LFS	None	106	1	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	58	1	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	61	1	Suisun Marsh

Water Year	Phase	Station Code	Date	# Tows	Species	Mark Type	Fork Length	Total Catch	Stratum
2022	1	22-27-GB02	1/31/2022	4	LFS	None	63	1	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	64	2	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	66	1	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	67	1	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	69	1	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	70	1	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	71	1	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	72	1	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	73	1	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	74	2	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	75	3	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	76	2	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	77	1	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	78	2	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	80	1	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	82	1	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	87	1	Suisun Marsh
2022	1	22-27-GB02	1/31/2022	4	LFS	None	88	1	Suisun Marsh
2022	1	22-27-SM01	1/31/2022	4	LFS	None	43	1	Suisun Marsh

Water Year	Phase	Station Code	Date	# Tows	Species	Mark Type	Fork Length	Total Catch	Stratum
2022	1	22-27-SM01	1/31/2022	4	LFS	None	68	1	Suisun Marsh
2022	1	22-27-SM01	1/31/2022	4	LFS	None	72	1	Suisun Marsh
2022	1	22-27-SM01	1/31/2022	4	LFS	None	76	2	Suisun Marsh
2022	1	22-27-SM01	1/31/2022	4	LFS	None	77	2	Suisun Marsh
2022	1	22-27-SM01	1/31/2022	4	LFS	None	78	1	Suisun Marsh
2022	1	22-27-SM01	1/31/2022	4	LFS	None	80	1	Suisun Marsh
2022	1	22-27-SM01	1/31/2022	4	LFS	None	84	2	Suisun Marsh
2022	1	22-27-SM01	1/31/2022	4	LFS	None	86	1	Suisun Marsh
2022	1	22-27-SM01	1/31/2022	4	LFS	None	87	1	Suisun Marsh
2022	1	22-27-RV02	2/1/2022	3	NA	NA	NA	NA	Lower Sac River
2022	1	22-27-LSR01	2/3/2022	4	NA	NA	NA	NA	Lower Sac River
2022	1	22-27-LSR02	2/3/2022	4	LFS	None	71	1	Lower Sac River
2022	1	22-27-LSR02	2/3/2022	4	LFS	None	72	1	Lower Sac River
2022	1	22-27-LSR02	2/3/2022	4	LFS	None	79	1	Lower Sac River
2022	1	22-27-LSR02	2/3/2022	4	LFS	None	80	1	Lower Sac River
2022	1	22-27-LSR02	2/3/2022	4	LFS	None	84	1	Lower Sac River

Water Year	Phase	Station Code	Date	# Tows	Species	Mark Type	Fork Length	Total Catch	Stratum
2022	1	22-27-LSR02	2/3/2022	4	LFS	None	87	1	Lower Sac River
2022	1	22-27-RV01	2/3/2022	4	NA	NA	NA	NA	Lower Sac River
2022	1	22-28-LSR01	2/7/2022	4	DSM	AdClipped	55	1	Lower Sac River
2022	1	22-27-PP01	1/31/2022	4	NA	NA	NA	NA	Lower San Joaquin River
2022	1	22-27-PP02	1/31/2022	4	NA	NA	NA	NA	Lower San Joaquin River
2022	1	22-27-PP03	1/31/2022	4	NA	NA	NA	NA	Lower San Joaquin River
2022	1	22-27-LSJ01	2/4/2022	4	DSM	AdClipped	57	1	Lower San Joaquin River
2022	1	22-27-LSJ02	2/4/2022	4	LFS	None	76	1	Lower San Joaquin River
2022	1	22-27-SJT01	2/4/2022	4	NA	NA	NA	NA	Lower San Joaquin River
2022	1	22-27-LSSC01	1/31/2022	4	NA	NA	NA	NA	Sac DW Ship Channel

Water Year	Phase	Station Code	Date	# Tows	Species	Mark Type	Fork Length	Total Catch	Stratum
2022	1	22-27-LSSC02	1/31/2022	4	NA	NA	NA	NA	Sac DW Ship Channel
2022	1	22-27-LSSC03	1/31/2022	2	DSM	AdClipped	66	1	Sac DW Ship Channel
2022	1	22-27-LSSC03	1/31/2022	2	DSM	AdClipped	70	1	Sac DW Ship Channel

Unmarked DSM collected during Phase 1 are transferred alive to FCCL to contribute to DSM broodstock if tow temperatures are below 17°C. If tow temperatures are above 17°C, unmarked DSM are flash frozen in liquid nitrogen and transferred to the UC Davis Aquatic Health Program for processing. All marked DSM are flash frozen in liquid nitrogen and transferred to UC Davis for processing.

DRAFT




Figure 1. Delta Smelt (DSM) and Longfin Smelt (LFS) catch per station for EDSM 2022 Phase 1 Kodiak trawls, from 1/31/2022–2/4/2022. Sites with no DSM or LFS catch are indicated with squares.

Table 2. Delta Smelt (DSM) and Longfin Smelt (LFS) catch in Chipps Island mid-water trawls from 1/30/2022–2/04/2022. These data are preliminary and subject to change.

Year	Station Code	Date	Species	Fork Length	Total Catch	Mark Type	Special Study	Location
2022	SB018M	2/3/2022	LFS	82	1	None	n/p	Chipps Island
2022	SB018M	2/3/2022	LFS	88	1	None	n/p	Chipps Island
2022	SB018N	2/3/2022	LFS	83	1	None	n/p	Chipps Island
2022	SB018N	2/3/2022	LFS	82	1	None	n/p	Chipps Island
2022	SB018N	2/3/2022	LFS	79	1	None	n/p	Chipps Island

Year	Station Code	Date	Species	Fork Length	Total Catch	Mark Type	Special Study	Location
2022	SB018N	2/3/2022	LFS	81	1	None	n/p	Chippis Island
2022	SB018S	2/3/2022	LFS	73	1	None	n/p	Chippis Island
2022	SB018M	2/4/2022	LFS	84	1	None	n/p	Chippis Island
2022	SB018M	2/4/2022	LFS	90	1	None	n/p	Chippis Island
2022	SB018M	2/4/2022	LFS	68	1	None	n/p	Chippis Island
2022	SB018M	2/4/2022	LFS	72	1	None	n/p	Chippis Island
2022	SB018M	2/4/2022	LFS	82	1	None	n/p	Chippis Island
2022	SB018N	2/4/2022	LFS	88	1	None	n/p	Chippis Island
2022	SB018N	2/4/2022	LFS	88	1	None	n/p	Chippis Island
2022	SB018N	2/4/2022	LFS	70	1	None	n/p	Chippis Island
2022	SB018N	2/4/2022	LFS	79	1	None	n/p	Chippis Island
2022	SB018S	2/4/2022	LFS	75	1	None	n/p	Chippis Island
2022	SB018S	2/4/2022	LFS	87	1	None	n/p	Chippis Island

As requested, LFS > 80 mm fork length collected in Chippis Island trawls during Dec-Apr are transferred alive to FCCL to contribute to LFS broodstock if tow temperatures are below 14.5°C. All DSM are flash frozen in liquid nitrogen and transferred to the UC Davis Aquatic Health Program for processing. L*: Fish dead, taken back to LFWO lab for preservation.

Table 3. Longfin Smelt catch per station from 2022 Smelt Larva Survey, Survey 2. Longfin Smelt incidental take permit criteria stations are highlighted in blue (Barker Slough Pumping Plant) and yellow (South Delta exports). Survey 2 was conducted between 1/24/2022 - 1/28/2022

Year	Survey #	SLS Station	Turbidity (NTU)	Sample Status	Species	Smelt Catch	Min Length	Max Length	Mean Length
2022	2	340	22.0	Not yet processed	NA	NA	NA	NA	NA
2022	2	342	18.2	Not yet processed	NA	NA	NA	NA	NA
2022	2	343	16.7	Not yet processed	NA	NA	NA	NA	NA
2022	2	344	20.6	Not yet processed	NA	NA	NA	NA	NA
2022	2	345	18.1	Not yet processed	NA	NA	NA	NA	NA
2022	2	346	18.3	Not yet processed	NA	NA	NA	NA	NA
2022	2	347	14.1	Not yet processed	NA	NA	NA	NA	NA
2022	2	348	9.7	Not yet processed	NA	NA	NA	NA	NA
2022	2	349	9.3	Not yet processed	NA	NA	NA	NA	NA
2022	2	405	70.4	Not yet processed	NA	NA	NA	NA	NA
2022	2	411	23.3	Processed	Longfin Smelt	1	7	7	7.0
2022	2	418	10.6	Processed	NA	No Smelt Catch	NA	NA	NA
2022	2	501	19.2	Processed	Longfin Smelt	17	7	10	8.0
2022	2	504	22.4	Processed	Longfin Smelt	6	7	9	8.3
2022	2	508	29.9	Processed	Longfin Smelt	22	6	10	7.5
2022	2	513	39.2	Processed	Longfin Smelt	48	6	12	7.4
2022	2	519	28.9	Processed	Longfin Smelt	14	7	9	8.1
2022	2	520	34.8	Not yet processed	NA	NA	NA	NA	NA
2022	2	602	35.2	Not yet processed	NA	NA	NA	NA	NA
2022	2	606	47.6	Processed	Longfin Smelt	56	7	12	8.4
2022	2	609	73.6	Processed	Longfin Smelt	8	8	14	11.0
2022	2	610	46.6	Processed	NA	No Smelt Catch	NA	NA	NA

Year	Survey #	SLS Station	Turbidity (NTU)	Sample Status	Species	Smelt Catch	Min Length	Max Length	Mean Length
2022	2	703	37.8	Processed	Longfin Smelt	4	6	7	6.8
2022	2	704	41.9	Processed	Longfin Smelt	14	6	7	6.7
2022	2	705	16.1	Processed	Longfin Smelt	1	7	7	7.0
2022	2	706	37.3	Processed	Longfin Smelt	15	6	7	6.4
2022	2	707	18.5	Processed	Longfin Smelt	41	6	7	6.5
2022	2	711	6.8	Processed	NA	No Smelt Catch	NA	NA	NA
2022	2	716	12.7	Processed	Longfin Smelt	5	6	7	6.6
2022	2	723	11.6	Processed	Longfin Smelt	5	6	7	6.2
2022	2	801	28.9	Processed	Longfin Smelt	63	6	10	7.3
2022	2	804	21.9	Processed	Longfin Smelt	26	6	8	6.7
2022	2	809	29.1	Processed	Longfin Smelt	6	6	8	7.2
2022	2	812	16.7	Processed	Longfin Smelt	24	6	8	7.5
2022	2	815	13.8	Processed	Longfin Smelt	2	6	7	6.5
2022	2	901	17.4	Processed	NA	No Smelt Catch	NA	NA	NA
2022	2	902	19.6	Processed	NA	No Smelt Catch	NA	NA	NA
2022	2	906	9.0	Processed	Longfin Smelt	1	8	8	8.0
2022	2	910	7.0	Processed	NA	No Smelt Catch	NA	NA	NA
2022	2	912	11.0	Processed	NA	No Smelt Catch	NA	NA	NA
2022	2	914	9.1	Processed	NA	No Smelt Catch	NA	NA	NA
2022	2	915	13.6	Processed	NA	No Smelt Catch	NA	NA	NA
2022	2	918	7.8	Processed	NA	No Smelt Catch	NA	NA	NA
2022	2	919	5.0	Processed	NA	No Smelt Catch	NA	NA	NA

Processing is complete through 2/07/2022