

California Wildlife Habitat Relationships System
California Department of Fish and Wildlife
California Interagency Wildlife Task Group

CHESTNUT-BACKED CHICKADEE

Poecile rufescens

Family: PARIDAE
B357

Order: PASSERIFORMES

Class: AVES

Written by: M. Green

Reviewed by: L. Mewaldt, D. Airola

Edited by: R. Duke, D. Winkler

DISTRIBUTION, ABUNDANCE, AND SEASONALITY

Common to abundant throughout most of its range; but rare and local at eastern and southern limits. Ranges inland from north coast area into Siskiyou and Trinity cos. and extends down western slope of the Sierra Nevada to Madera Co., where it is rare. Range extends along north coast area to Lake and Solano cos., then tapering toward the coast from Contra Costa Co. south to Santa Barbara Co. Prefers closed-cone pine-cypress, Douglas-fir, and redwood habitats, but often found in riparian, hardwood, and other habitats (Grinnell and Miller 1944, Crase 1976, McCaskie et al. 1979).

SPECIFIC HABITAT REQUIREMENTS

Feeding: Gleans insects and spiders from foliage, twigs, and branches of trees, especially conifers; eats some conifer seeds and fruits especially in fall and winter. Feeds mostly in middle and upper canopy levels (Airola and Barrett 1985).

Cover: Cover provided by trees and shrubs in humid forests of the coast and mixed forests of the western Sierra Nevada. Frequents closed-cone pine-cypress, redwood, riparian, and Douglas-fir habitats. Also, regularly found in moist hardwood habitats.

Reproduction: Builds nest of moss with cup of fur, hair, feathers, plant fibers (Harrison 1978); usually in old woodpecker hole, or excavates own cavity in decaying wood. Nest height usually less than 3 m (10 ft), but has been found up to 24 m (80 ft) (Bent 1946). Readily uses nest boxes.

Water: Bathes and drinks occasionally.

Pattern: Frequents conifer habitats, especially those with oaks, maples, and other hardwoods. Also feeds regularly in alders, willows, and other riparian vegetation.

SPECIES LIFE HISTORY

Activity Patterns: Yearlong, diurnal activity.

Seasonal Movements/Migration: A yearlong resident in California; appears to wander very little. Existence on the western slope of Sierra Nevada results from a recent, major range expansion (Crase 1976).

Home Range: Apparently smaller in summer.

Territory: Hertz et al. (1976) reported a mean breeding territory of 1.3 ha (3.3 ac) (N = 2) in an oak woodland in San Mateo Co.

Reproduction: Breeds from mid-March into July; peak in May and June. Lays 5-9 eggs; average 6-7 (Bent 1946). Altricial young cared for by both parents (Harrison 1978).

Niche: Subject to same predators as other small, forest-dwelling passerines: adults to predation from accipiters, and eggs and nestlings to predation from small mammals, snakes, and corvids. May join flocks of mixed species in winter and forage beyond summer home range (Airola and Barrett 1985).

REFERENCES

- Airola, D. A., and R. H. Barrett. 1985. Foraging and habitat relationships of insect-gleaning birds in a Sierra Nevada mixed-conifer forest. *Condor* 87:205-216.
- Bent, A. C. 1946. Life histories of North American jays, crows, and titmice. U.S. Natl. Mus. Bull. 191. 495pp.
- Crane, F. T. 1976. Occurrence of the chestnut-backed chickadee in the Sierra Nevada Mountains, California. *Am. Birds* 30:673-675.
- Garrett, K., and J. Dunn. 1981. Birds of southern California. Los Angeles Audubon Soc. 408pp.
- Grinnell, J., and A. H. Miller. 1944. The distribution of the birds of California. Pac. Coast Avifauna No. 27. 608pp.
- Harrison, C. 1978. A field guide to the nests, eggs and nestlings of north American birds. W. Collins Sons and Co., Cleveland, OH. 416pp.
- Hertz, P. E., J. V. Remsen, Jr., and S. I. Zones. 1976. Ecological complementarity of three sympatric parids in a California oak woodland. *Condor* 78:307-316.
- McCaskie, G., P. De Benedictis, R. Erickson, and J. Morlan. 1979. Birds of northern California, an annotated field list. 2nd ed. Golden Gate Audubon Soc., Berkeley. 84pp.

B357

Life history accounts for species in the California Wildlife Habitat Relationships (CWHR) System were originally published in: Zeiner, D.C., W.F.Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. California's Wildlife. Vol. I-III. California Department of Fish and Game, Sacramento, California. Updates are noted in accounts that have been added or edited since original publication.