

California Wildlife Habitat Relationships System
California Department of Fish and Wildlife
California Interagency Wildlife Task Group

CRISSAL THRASHER

Toxostoma crissale

Family: MIMIDAE
B399

Order: PASSERIFORMES

Class: AVES

Written by: D. Dobkin, S. Granholm
Reviewed by: L. Mewaldt
Edited by: R. Duke

DISTRIBUTION, ABUNDANCE, AND SEASONALITY

A resident of southeastern deserts. Still fairly common in Colorado River Valley, but local and uncommon elsewhere. Occupies dense thickets of shrubs or low trees in desert riparian and desert wash habitats. In eastern Mojave Desert of San Bernardino and southeastern Inyo cos., also occurs in dense sagebrush and other shrubs in washes within juniper and pinyon-juniper habitats, up to 1800 m (5900 ft). Also resident in Imperial, Coachella, and Borrego valleys, but numbers have declined markedly in recent decades (Grinnell and Miller 1944, Remsen 1978, Garrett and Dunn 1981).

SPECIFIC HABITAT REQUIREMENTS

Feeding: Forages mostly on ground, especially between and under shrubs. Uses bill to dig in friable soil and to probe in litter. Diet poorly known, but includes insects, other invertebrates, berries and other small fruits, seeds, and occasionally small lizards (Bent 1948). Laudenslayer (1981) reported a diet of 69% Coleoptera, 10% Hymenoptera, 20% miscellaneous insects, and 1% plant material.

Cover: Cover provided by thickets of dense, shrubby vegetation along streams and in washes; frequently mesquite, screwbean mesquite, ironwood, catclaw acacia, arrowweed willow.

Reproduction: Builds nest within thicket of desert shrubs or on a forked branch of a small tree. Nest usually placed 0.6 to 3 m (2-10 ft) above ground.

Water: Bartholomew and Cade (1956) reported mean water consumption in captive individuals at 20-23°C of about 5% body weight per 24 hr.

Pattern: Frequents dense thickets in desert riparian and desert wash habitats.

SPECIES LIFE HISTORY

Activity Patterns: Yearlong, diurnal activity.

Seasonal Movements/Migration: Sedentary. Not migratory.

Home Range: In southern Nevada, mated pairs remained in only a few hundred ft of mesquite-bosque oak habitat, based on banding records (Gullion et al. 1959). Reported density per 40 ha (100 ac) include: 0.5 in Arizona desert grassland (Grue 1977); 0.2 to 0.6 pairs in New Mexico creosote scrub (Raitt and Maze 1968); 1.0 pair in Arizona oak woodland (Balda 1970); 2.0 to 3.5 pairs in dunes, and 1.0 to 2.0 pairs in bosque, in southern Nevada (Austin 1970). Ryder and Ryder (1976) found 2 on a 17 ha (44 ac) study site in cat claw-rabbit brush habitat in San Bernardino Co.

Territory: Laudenslayer (1981) reported territory varying from 1.52 ha (3.81 ac) to 3.71 ha (9.28 ac) in mesquite-salt cedar associations along the Colorado River.

Reproduction: Breeding season lasts from February into June with a peak in March and April. Clutch size 2-4 eggs, mostly 2 or 3. Frequently raises 2 broods per yr. Incubation lasts about 2 wk. Altricial young usually remain in nest 11-12 days; up to 16 days. Oberholser (1974) noted small groups, assumed to be families, congregating toward end of breeding season and in fall.

Niche: Of all arid-land thrashers, occurs in coolest, most shaded, and moist habitats. A California Species of Special Concern. Numbers have been reduced greatly by removal of mesquite brushland for agricultural development, and by introduction of tamarisk (Remsen 1978). Off-road vehicle activity also may degrade habitat and disturb these thrashers. Ejects cowbird eggs from nest.

REFERENCES

- Austin, G. T. 1970. Breeding birds of desert riparian habitat in southern Nevada. *Condor* 72:431-436.
- Balda, R. P. 1970. Effects of spring leaf-fall on composition and density of breeding birds in two southern Arizona woodlands. *Condor* 72:325-331.
- Bartholomew, G. A., and T. J. Cade. 1956. Water consumption of house finches. *Condor* 58:406-412.
- Bent, A. C. 1948. Life histories of North American nuthatches, wrens, thrashers, and their allies. U.S. Natl. Mus. Bull. 195. 475pp.
- Engels, W. L. 1940. Structural adaptations in thrashers (Mimidae: Genus *Toxostoma*) with comments on interspecific relationships. *Univ. Calif. Publ. Zool.* 42:341-400.
- Garrett, K., and J. Dunn. 1981. Birds of southern California. Los Angeles Audubon Soc. 408pp.
- Grinnell, J., and A. H. Miller. 1944. The distribution of the birds of California. *Pac. Coast Avifauna* No. 27. 608pp.
- Grue, C. E. 1977. The impact of powerline construction on birds in Arizona. M.S. Thesis, Northern Arizona Univ., Flagstaff. 265pp.
- Gullion, G. W., W. M. Pulich, and F. G. Evenden. 1959. Notes on the occurrence of birds in southern Nevada. *Condor* 61:278-297.
- Laudenslayer, W. F., Jr. 1981. Habitat utilization by birds of three desert riparian communities. Ph.D. Thesis, Arizona State Univ., Tempe. 148pp.
- Oberholser, H. C. 1974. The bird life of Texas. 2 vols. Univ. Texas Press. Austin. 1069pp.
- Raitt, E. J., and R. L. Maze. 1968. Densities and species composition of breeding birds of a creosote bush community in southern New Mexico. *Condor* 70:193-205.
- Remsen, J. V., Jr. 1978. Bird species of special concern in California. Calif. Dep. Fish and Game, Sacramento. Wildl. Manage. Admin. Rep. No. 78-1. 54pp.
- Ryder, R. A., and A. Ryder. 1976. Twenty-ninth winter bird population study. *Am. Birds* 30:1040-1075.