

State Water Project Incidental Take Permit Risk Assessment for Delta Smelt and Longfin Smelt

Section 1: Overview

Date: 05/02/2023

Life Stages Present:

Delta Smelt (DS): Larvae, Sub-adults, and Adults

Longfin Smelt (LFS): Larvae, Juveniles, Sub-adults, and Adults

Advice to Water Operations Management Team (WOMT):

No recommendation to WOMT this week.

Condition of Approval (COA) 8.5.2 is active with water temperature exceeding 12°C at Jersey Point and triggered due to 20mm Survey 4 Secchi reading being < 1m. However, this COA is not controlling because OMRI is highly positive and is predicted to remain highly positive throughout the week.

Risk Assessment:

Delta Smelt: Based on recent detection data and distribution patterns over the past decade, Delta Smelt are spawning and larval Delta Smelt are present. No adult Delta Smelt have been detected since 3/21/23. Twenty-two larval Delta Smelt (21 confirmed/1 preliminary) have been detected since 3/13/23. No Delta Smelt have been detected in salvage since 3/2/23. Three-day average water temperature at Jersey Point exceeded 12° C on 3/18/23, and the most recent Secchi depths in the South Delta were below 1m, triggering COA 8.5.2. However, these actions are not controlling OMRI. Due to highly positive QWEST and OMRI, overall risk for entrainment is low for all life stages of Delta Smelt throughout the Delta.

Longfin Smelt: Risk remains low for all life stages as a result of favorable hydrology. COA 8.4.3 was triggered on 3/3/23 and remains triggered with San Joaquin River flow at Vernalis continuing to exceed 5,000 cfs. No LFS were detected in the Central and South Delta stations by 20mm Survey 4. No salvage was detected since 3/2/23, and the cumulative adult and sub-adult seasonal salvage is 26. X2 is slightly east of Martinez (~59 km) and QWEST remains highly positive (~34,000 cfs). Many fish were detected by SKT, SLS, 20mm Survey, and EDSM in and westward of Suisun Bay, suggesting that LFS are distributed widely. Adult detections have been decreasing, suggesting that spawning is reaching the end. Four juvenile LFS was detected in Grizzly Bay and San Pablo Bay by EDSM Kodiak Trawl but no adult or sub-adult LFS were detected by EDSM or Chipps Island Trawl (4/24/23 – 4/28/23). Fish distribution remains downstream with high outflow.

Section 1-A: Sacramento River and Confluence

Table 1: Risk of entrainment into the Central Delta and export facilities for Delta Smelt in the Sacramento River and Confluence:

Species and life stage	Risk type	Risk level	Rationale (turbidity, exports, OMR level, X2, Q west, temperature, distribution etc.)
DS larvae and juveniles	Exposure Risk (Hydrology)	Low	Spawning is ongoing. Four larval DS were detected (QC completed) near the Confluence, Cache Slough, SDWSC, and Miner Slough by 20mm Survey 3. One larval DS was detected in the Confluence by EDSM 20mm trawl on 4/20/23. One larval DS was detected in the Confluence by 20mm Survey 4.
DS subadults and adults	Routing Risk (Behavior and life history)	Low	Spawning is ongoing. Distribution is widespread.
DS	Overall Entrainment Risk	Low	Same as above.

Table 2: Risk of entrainment into the Central Delta and export facilities for Longfin Smelt in the Sacramento River and Confluence:

Species and life stage	Risk type	Risk level	Rationale (turbidity, exports, OMR level, X2, Q west, temperature, distribution etc.)
LFS larvae and juveniles	Exposure Risk (Hydrology)	Low	Seven larval LFS were detected in the Lower Sacramento River and the Confluence by 20mm Survey 3 (processing is on-going). One larval LFS was detected in the Lower Sacramento River by EDSM from 4/10/23 to 4/14/23.
LFS sub-adults and adults	Routing Risk (Behavior and life history)	Low	Spawning is ending. X2 is ~59 km. Fish are widely distributed resulting in low risk. One adult LFS was detected in the Lower Sacramento River by EDSM on 4/10/23. No LFS were detected by Chipps Island Trawl from 4/24/23 to 4/28/23.
LFS	Overall Entrainment Risk	Low	Same as above.

Section 1-B: Central Delta

Table 3: Risk of entrainment into the export facilities for Delta Smelt in the Central Delta:

Species and life stage	Risk type	Risk level	Rationale (turbidity, exports, OMR level, X2, Q west, temperature, distribution etc.)
DS larvae and juveniles	Exposure Risk (Hydrology)	Low	Spawning is on-going, but no larvae have been detected in this region. No detections in salvage and qualitative larval samples over the last week.
DS subadults and adults	Exposure Risk (Hydrology)	Low	No adult or sub-adult DS were detected in salvage or in field surveys in this region in the last two weeks (4/17/23 – 5/1/23).

Table 4: Risk of entrainment into the export facilities for Longfin Smelt in the Central Delta:

Species and life stage	Risk type	Risk level	Rationale (turbidity, exports, OMR level, X2, Q west, temperature, distribution etc.)
LFS larvae	Exposure Risk (Hydrology)	Low	20mm Survey 4 detected no LFS in the Central and South Delta. No detections in salvage and qualitative larval samples over the last week.
LFS sub-adults and adults	Exposure Risk (Hydrology)	Low	No sub-adult and adult LFS were detected in salvage or in field surveys in this region in the last two weeks (4/17/23 – 5/1/23).

- Change in exposure from previous week: *(Note: The change in risk compared to previous weeks is not required by the Incidental Take Permit [ITP]).*
 - DS: Overall risk of entrainment remains low for all life stages of DS.
 - LFS: Overall risk of entrainment remains low for all life stages of LFS.
- Reporting Old and Middle River Index (OMRI) *(Number and range of OMRI bins will vary based on anticipated hydrology and operations)*
 - Expected daily OMRI range this week: +3,500 to +11,000 cfs

Section 2: Basis for Advice

The 2020 ITP ([Incidental Take Permit for Long-Term Operation of the State Water Project in the Sacramento-San Joaquin Delta 2081-2019-066-00](#)) states that advice to WOMT shall be based on the following Conditions of Approval:

List relevant Condition of Approval number and title based on species/life stage, time of year, etc.

8.3.1 Integrated Early Winter Pulse Protection. Between December 1 and January 31 each year Permittee shall reduce south Delta exports for 14 consecutive days to maintain a 14-day average OMRI no more negative than -2,000 cfs, and convene the SMT within one day of triggering the following criteria:

- Three-day running average daily flows at Freeport greater than, or equal to, 25,000 cfs, AND
- Three-day running average of daily turbidity at Freeport greater than, or equal to, 50 Nephelometric Turbidity Units (NTU), OR
- The SMT determines that real-time monitoring of abiotic and biotic factors indicates a high risk of DS migration and dispersal into areas at high risk of future entrainment.

After maintaining a 14-day average OMRI no more negative than -2,000 cfs for 14 days, Permittee shall maintain a 14-day average OMRI no more negative than -5,000 cfs, initiating the OMR Management season, until the OMR Management Season ends (Condition of Approval 8.8).

The Integrated Early Winter Pulse Protection Action may only be initiated once during the December 1 through January 31 time period each year.

8.3.2 Salmonid Presence. After January 1 each year, if Conditions of Approval 8.3.1 or 8.3.3 have not already been triggered, the OMR Management season shall begin when the Salmon Monitoring Team first estimates that 5% of the CHNWR or CHNSR population is in the Delta whichever is sooner. Upon initiation of the OMR Management season, Permittee shall reduce exports to achieve, and shall maintain a 14-day average OMR index no more negative than -5,000 cfs, until the OMR Management season ends (see Condition of Approval 8.8). In the event that a salmon daily or single-year loss threshold is exceeded (Conditions of Approval 8.6.1, 8.6.2, 8.6.3, or 8.6.4) prior to the start of OMR Management season the requirements in those Conditions shall control operations.

8.3.3 Adult Longfin Smelt Entrainment Protection. After December 1, if an Integrated Early Winter Pulse Protection (Condition of Approval 8.3.1) has not yet initiated, Permittee shall reduce south Delta exports to maintain a 14-day average OMRI no more negative than -5,000 cfs and initiate OMR Management (Condition of Approval 8.3) if:

- Cumulative combined LFS salvage (total estimated LFS counts at the CVP and SWP salvage facilities beginning December 1 through February 28 exceeds the most recent Fall Midwater Trawl (FMWT) LFS index¹ divided by 10, OR
- Real-time monitoring of abiotic and biotic factors indicates a high risk of LFS movement into areas at high risk of future entrainment, as determined by DWR and CDFW SMT staff.

When evaluating the possibility of LFS movement into areas that may be subject to an elevated risk of entrainment, the SMT shall evaluate catch of LFS with fork length ≥ 60 mm by the Chippis Island Trawl (conducted by USFWS) as an early warning indicator for LFS migration movement into the Delta, in addition to other available survey and abiotic data. The SMT shall communicate the results of these risk assessments and advice to the WOMT (Condition of Approval 8.1.3), and operational decisions shall be made as described in Condition of Approval 8.1.4 (Collaborative Approach to Real-time Risk Assessment).

8.4.1 OMR Management for Adult Longfin Smelt. From the onset of OMR Management (Condition of Approval 8.3) through February 28, the SMT shall conduct weekly, or more often as needed, risk assessments (see Condition of Approval 8.1.5.2) and decide whether to recommend an OMR flow requirement between -5,000 cfs and -1,250 cfs to minimize entrainment and take of adult LFS. The SMT may provide advice to restrict south Delta exports for seven consecutive days to achieve a seven-day average OMRI within three risk categories:

- Low risk: OMR between -4,000 cfs to -5,000 cfs
- Medium risk: OMR between -2,500 cfs to -4,000 cfs
- High risk: OMR between -1,250 cfs to -2,500 cfs

If a risk assessment conducted by the SMT determines that a more restrictive OMR flow requirement is needed to minimize take of adult LFS, the SMT shall provide its advice to WOMT (Condition of Approval 8.1.3) and operational decisions shall be made following the process described in Condition of Approval 8.1.4 (Collaborative Approach to Real-time Risk Assessment).

This Condition will terminate when a high-flow off-ramp occurs (Condition of Approval 8.4.3), or when LFS spawning has been detected in the system, as determined by the SMT, or, if there is disagreement and resolution is not reached within WOMT, as determined by CDFW. The SMT shall consider results from Additional LFS Larval Sampling (Condition of Approval 7.6.1) to inform its assessment of the start of LFS spawning. After LFS spawning has been observed,

¹ The Fall Midwater Trawl (FMWT) Survey annual abundance index for LFS is calculated as the sum of September through December monthly abundance indices and is typically reported at about the same date as adult salvage begins in December. The FMWT Index available beginning on December 1 each year shall be used to establish this threshold.

Permittee shall implement Condition of Approval 8.4.2 to minimize take of larval and juvenile LFS.

8.4.2 Larval and Juvenile Longfin Smelt Entrainment Protection. From January 1 through June 30, when a single Smelt Larva Survey (SLS) or 20 mm Survey (20 mm) sampling period exceeds one of the following thresholds:

- LFS larvae or juveniles found in four or more of the 12 SLS or 20 mm stations in the central Delta and south Delta (Stations 809, 812, 815, 901, 902, 906, 910, 912, 914, 915, 918, 919), or
- LFS catch per tow exceeds five LFS larvae or juveniles in two or more of the 12 stations in the central Delta and south Delta (Stations 809, 812, 815, 901, 902, 906, 910, 912, 914, 915, 918, 919).

Permittee shall restrict south Delta exports for seven consecutive days to maintain a seven-day average OMR index no more negative than -5,000 cfs. Permittee shall also immediately convene the SMT to conduct a risk assessment (see Condition of Approval 8.5.1.2) to assess the risk of larval and juvenile LFS entrainment into the South Delta Export Facilities, determine if an OMR flow restriction is warranted, and recommend an OMR flow limit between -1,250 and -5,000 cfs. The SMT risk assessment and operational advice shall be reviewed by the WOMT (Condition of Approval 8.1.3) via the Collaborative Real-time Decision-making process (Condition of Approval 8.1.4). Permittee shall operate to the export restriction and OMR flow target approved through Conditions of Approval 8.1.3 and 8.1.4. Each week the SMT shall convene to conduct a new risk assessment and determine whether to maintain, or off ramp from, export restrictions based on the risk to LFS, or until the DS and LFS off-ramp has been met as described in Condition of Approval 8.8 (End of OMR Management).

From January 1 through June 30, DWR and CDFW SMT staff shall conduct weekly, or more often as needed, risk assessments (see Condition of Approval 8.5.1.2) to assess the risk of larval and juvenile LFS entrainment into the South Delta Export Facilities. As a part of the risk assessment the SMT shall provide advice on the appropriate OMR flow targets to minimize LFS entrainment or entrainment risk, or both. The SMT shall provide its advice to WOMT (Condition of Approval 8.1.3) and use the Collaborative Approach to Real-time Risk Assessment process described in Condition of Approval 8.1.4 to determine if an OMR flow restriction is warranted and determine OMR flow limit between -1,250 and -5,000 cfs. The OMR flow limit shall be in place until the next risk assessment conducted by the SMT determines that it is no longer necessary to minimize take or related impacts to LFS, or until the DS and LFS off-ramp has been met as described in Condition of Approval 8.8 (End of OMR Management).

8.4.3 High Flow Off-Ramp from Longfin Smelt OMR Restrictions. OMR management for adult, juvenile, or larval LFS as described in Conditions of Approval 8.4.1 and 8.4.2 are not required, or would cease if previously required, when river flows are (a) greater than 55,000 cfs in the Sacramento River at Rio Vista or (b) greater than 8,000 cfs in the San Joaquin River at Vernalis. If flows subsequently drop below 40,000 cfs in the Sacramento River at Rio Vista or below 5,000

cfs in the San Joaquin River at Vernalis, the OMR limit previously required as a part of Conditions of Approval 8.4.1 and 8.4.2 shall resume.

8.5.1 Turbidity Bridge Avoidance. The purpose of this Condition is to minimize the risk of entrainment of adult DS in the corridors of the Old and Middle rivers into the south Delta export facilities. This Condition is intended to avoid the formation of a turbidity bridge from the San Joaquin River shipping channel to the south Delta export facilities, which historically has been associated with elevated salvage of pre-spawning adult DS.

After the Integrated Early Winter Pulse Protection (Condition of Approval 8.1.3) or February 1 (whichever comes first), until April 1, Permittee shall manage exports to maintain daily average turbidity in Old River at Bacon Island (OBI) at a level of less than 12 FNU. If the daily average turbidity at OBI is greater than 12 FNU, Permittee shall restrict south Delta exports to achieve an OMR flow that is no more negative than -2,000 cfs until the daily average turbidity at OBI is less than 12 FNU.

If, after five consecutive days of OMR flow that is less negative than -2,000 cfs, the daily average turbidity at OBI is not less than 12 FNU the Smelt Monitoring Team may convene to assess the risk of entrainment of DS (Condition of Approval 8.1.5.2). The Smelt Monitoring Team may provide advice to WOMT regarding changes in operations that could be conducted to minimize the risk of entrainment of DS (Condition of Approval 8.1.3). The Smelt Monitoring Team may also determine that OMR restrictions to manage turbidity are infeasible and may instead provide advice for a different OMR flow target that is between -2,000 and -5,000 cfs and is protective based on turbidity and adult DS distribution and salvage to the WOMT for consideration (Condition of Approval 8.1.3). Operational decisions shall be made following the process described in Condition of Approval 8.1.4 (Collaborative Real Time Risk Assessment).

Turbidity readings at individual sensors can generate spurious results in real time. Spurious results could be incorrectly interpreted as a turbidity bridge, when in fact the cause is a result of local conditions or sensor error. To assess whether turbidity readings at OBI are attributable to a sensor error or a localized turbidity spike, Permittee, in coordination with Reclamation, may consider and review data from other nearby locations and sources. Additional information that will be reviewed include regional visualizations of turbidity, alternative sensors, and boat-based turbidity mapping, particularly if there was evidence of a local sensor error. Permittee may bring data from these additional sources to the Smelt Monitoring Team for consideration during the development of a risk assessment to be provided to the WOMT for evaluation (Condition of Approval 8.1.3).

Permittee shall use the decision-making process described Condition of Approval 8.1.4 (Collaborative Real-time Risk Assessment) to determine if south Delta exports may increase after five-days of OMR no more negative than -2,000 cfs, or to determine that this action is not warranted due to a sensor error or localized turbidity

event. Permittee shall implement this action until CDFW is in agreement that the action may be ended or modified.

8.5.2 Larval and Juvenile Delta Smelt Protection. If the five-day cumulative salvage of juvenile DS at the CVP and SWP facilities is greater than or equal to one plus the average prior three years' FMWT index (rounded down), Permittee shall restrict south Delta exports for seven consecutive days to maintain a seven-day average OMR index no more negative than -5,000 cfs.

Additionally, if the five-day cumulative salvage threshold is met or exceeded, Permittee shall immediately convene the Smelt Monitoring Team to conduct a risk assessment (Condition of Approval 8.1.5.2) and determine the future risk of entrainment and take of larval and juvenile DS. The Smelt Monitoring Team may provide advice to further restrict south Delta exports to maintain a more positive OMR than -5,000 cfs. The Smelt Monitoring Team may provide advice for further restrictions within three risk categories:

- Low risk: Limit OMR between -4,000 cfs to -5,000 cfs
- Medium risk: Limit OMR between -2,500 cfs to -4,000 cfs
- High risk: Limit OMR between -1,250 cfs to -2,500 cfs

The duration and magnitude of operational advice shall be provided to the WOMT (Condition of Approval 8.1.3) and decisions shall be made following the process described in Condition of Approval 8.1.4 (Collaborative Real Time Risk Assessment). When conducting risk assessments to evaluate the risk of entrainment and take of juvenile DS the Smelt Monitoring Team shall evaluate the following information sources, in addition to any other models or surveys they deem appropriate and those listed in Condition of Approval 8.1.5.2:

- Results from a CDFW approved DS life cycle model.
- DS recruitment levels identified by the Smelt Monitoring Team using the CDFW approved life cycle model that links environmental conditions to recruitment, including factors related to loss as a result of entrainment such as OMR flows. In this context, recruitment is defined as the estimated number of post-larval DS in June per number of spawning adults in the prior February-March period.
- Hydrodynamic models and forecasts of entrainment informed by the EDSM or other relevant survey data to estimate the percentage of larval and juvenile DS that could be entrained.

When a larval or juvenile DS is detected in the SLS or 20 mm, or the 3-day average water temperature at Jersey Point is greater than or equal to 12°C, and Secchi depth from the most recent SLS or 20 mm survey is less than or equal to 1 meter, averaged across the 12 south Delta survey stations (809, 812, 815, 901, 902, 906, 910, 912, 914, 915, 918, and 919). Permittee shall restrict south Delta exports to maintain a seven-day average OMR index no more negative than -3,500 cfs until the average Secchi depth is greater than 1 meter in the south Delta stations in a subsequent SLS or 20mm survey. If average south Delta Secchi depth continues to be less than or equal to 1 meter in a subsequent SLS or 20mm survey then Permittee shall continue restrictions and request a risk assessment by the Smelt Monitoring Team to determine if

additional advice and subsequent restrictions are warranted and provide advice to WOMT (see Condition of Approval 8.1.3) and follow the decision-making process described in Condition of Approval 8.1.4.

8.12 Barker Slough Pumping Plant Longfin and Delta Smelt Protection. Permittee shall operate the BSPP to protect larval LFS from January 15 through March 31 of dry and critical water years. Permittee shall operate to protect larval DS from March 1 through June 30 of dry and critical years. If the water year type changes after January 1 to below normal, above normal or wet, this action will be suspended. If the water year type changes after January to dry or critical, Permittee shall operate according to this Condition of Approval.

From January 15 through March 31 of dry and critical water years, Permittee shall reduce the maximum seven-day average diversion rate at BSPP to less than 60 cfs when larval LFS are detected at Station 716. In addition, in its weekly meetings from January 15 through March 31, the Smelt Monitoring Team shall review LFS abundance and distribution survey data and other pertinent abiotic and biotic factors that influence the entrainment risk of larval LFS at the BSPP. When recommended by the Smelt Monitoring Team, and as approved through the decision making processes described in Conditions of Approval 8.1.3 and 8.1.4, Permittee shall reduce the maximum seven-day average diversion rate at BSPP according to the advice provided by the Smelt Monitoring Team.

From March 1 through June 30 of dry and critical water years, Permittee shall reduce the maximum seven-day average diversion rate at BSPP to less than 60 cfs when larval DS are detected at Station 716. In addition, in its weekly meetings from March 1 through June 30, the Smelt Monitoring Team shall review DS abundance and distribution survey data and other pertinent abiotic and biotic factors that influence the entrainment risk of larval DS at the BSPP (including temperature and turbidity). When recommended by the Smelt Monitoring Team, and as approved through the decision-making processes described in Conditions of Approval 8.1.3 and 8.1.4, Permittee shall reduce the maximum seven-day average diversion rate at BSPP to less than 60 cfs.

The DS requirements described in this condition may be adjusted to align with USFWS requirements to minimize take of DS through an amendment to this ITP.

8.13 Water Year Type Definition. All references to water year type in this ITP shall be defined based on the Sacramento Valley Index unless otherwise noted.

Discussion of Conditions of Approval

Provide discussion addressing criteria for each Condition of Approval listed in "Basis for Advice" section. Refer to data below where appropriate.

COAs relevant to OMR management went into effect December 1st. The Smelt Monitoring Team (SMT) conducted a Risk Assessment based on COA 8.1.5.2.

8.3.1: This COA was triggered by conditions measured on 12/31/22 when the three-day average of daily flow and turbidity was 26,552 cfs and 77 FNU and respectively. Operations were reduced on 1/3/23 targeting a 14-day average OMRI no more negative than -2,000 cfs for 14 consecutive days through 1/16/23. After 1/16/23, the 14-day average OMRI shall be no more negative than -5,000 cfs, initiating the OMR Management Season until the OMR Management Season ends (COA 8.8). This condition has been off-ramped as of 1/16/23.

8.3.2: This COA is no longer active due to the initiation of an Integrated Early Winter Pulse Protection (IEWPP- COA 8.3.1).

8.3.3: This COA is no longer active due to the initiation of an IEWPP (COA 8.3.1). One adult LFS was detected in salvage on 1/1/23, this was the first LFS salvage of WY 2023. The FMWT LFS index for September through December is 403, therefore the salvage threshold to trigger this COA is 40 LFS.

8.4.1: This COA is no longer active due to the detection of larval LFS by SLS.

8.4.2: This COA was triggered for the first time this season on 2/16/23 with the detection of LFS larvae at four (stations 809, 812, 901, and 902) of the 12 Central and South Delta stations by SLS 4. Exports were managed for seven consecutive days to maintain a seven-day average OMR index no more negative than -5,000 cfs. SMT convened for an off-cycle meeting on 2/17/23 and discussed the risk of larval LFS entrainment in the Central and South Delta while looking at the PTM run provided by DWR. There was non-consensus between CDFW and DWR, and the decision was elevated to WOMT. WOMT convened for an off-cycle meeting on 2/17/23 and decided that OMRI would be limited to -3,500 cfs on a seven-day average. On 2/21/23 SMT agreed to continue to recommend OMRI be limited to -3,500 cfs, however there was non-consensus on the duration of this protection between CDFW and DWR in SMT and WOMT on 2/21/23 and 2/22/23 respectively. This was elevated to the Directors, and they decided that DWR would manage exports to -3,500 cfs OMRI until SMT can reassess with further LFS data by SLS 5.

This COA was triggered for the second time this season on 2/28/23 with the detection of 14 LFS larvae in the Central and South Delta (Station 809, 812, 901, 902, and 915) by SLS 5. SMT did not come to consensus on a recommendation on 2/28/23. CDFW recommended OMRI be limited to -2,000 cfs on a 7-day average for the protection of larval LFS based on the PTM results and recent detections by SLS 5. DWR recommended limiting OMRI to -5,000 cfs due to a positive QWEST and improvement in hydrodynamic conditions relative to last week. The decision was elevated to WOMT. It was decided to keep the -3,500 cfs OMRI restrictions until COA 8.4.3 off-ramps COA 8.4.2 later in the week. COA 8.4.3 off-ramped this COA on 3/2/23.

This COA was not triggered by SLS 6. Twelve larvae were detected in three of the 12 Central and South Delta stations (Station 812, 815, and 901) by SLS 6 and one larvae (Station 901) was detected by 20mm Survey 1. Neither SLS 6 nor 20mm Survey 1 met the threshold for this COA, and thus this COA is no longer triggered as of 3/21/23.

This COA was not triggered by 20mm Surveys 2, 3, or 4. No LFS were detected at Central and South Delta stations by 20mm Surveys 2, 3, or 4.

8.4.3: This COA was triggered for the first time this season by the conditions measured on 1/3/23 when the flow of the San Joaquin River at Vernalis exceeded 8,000 cfs, but no longer triggered as of 2/11/23 due to flow in the San Joaquin River at Vernalis decreasing to under 5,000 cfs.

This COA was triggered for the second time this season by the conditions measured on 3/2/23 when the flow of San Joaquin River at Vernalis exceeded 8,000 cfs, and temporarily off-ramped COA 8.4.2. It is expected for this condition to remain triggered through the week of 5/1/23.

8.5.1: This condition was triggered for the first time this season on 1/18/23 by the conditions measured on 1/17/23 when the turbidity at OBI was 17 FNU. OMRI was limited to no more negative than -2,000 cfs. After the first five days (1/17/23 through 1/21/23), turbidity was still above 12 FNU at OBI, therefore the SMT reconvened to assess risk. The SMT reassessed risk for DS but was unable to reach consensus on a recommendation between -2,000 cfs and -5,000 cfs on 1/19/23. On 1/20/23 WOMT reached consensus to allow operational flexibility to maintain maximum exports until 1/24/23 when the SMT met again, which may have resulted in an OMRI as negative as -3,500 cfs. On 1/24/23 the OMRI had not reached -3,500 cfs (was -2,100 cfs as of 1/23/23) and proposed operations were to maintain maximum exports as long as possible and operate to an OMRI of -5,000 cfs for the week. The SMT reassessed risk for DS and determined that risk for DS in the South Delta was high and moderate outside the South Delta because of ongoing high turbidity. Additionally, the SMT agreed that risk of entrainment would increase if OMRI were to become more negative, however the SMT was unable to reach consensus on an OMRI recommendation. WOMT met on 1/25/23 and came to a consensus for -5,000 cfs OMRI for one week starting on 1/26/23. The SMT reassessed risk for DS on 1/31/23 and came to a consensus that no further restrictions were warranted, because turbidity was decreasing. On 2/9/23 daily turbidity at Old River at Bacon Island decreased to less than 12 FNU, therefore this COA was no longer triggered.

This COA was triggered for the second time this season by the conditions on 2/15/23. SMT agreed that a turbidity bridge had formed, and it was not a localized event nor a sensor error. Five-day average OMRI would have been restricted to -2,000 cfs for five days starting on 2/18/23, if turbidity didn't drop below 12 FNU on or before 2/18/23. On 2/17/23, the daily turbidity at OBI decreased to less than 12 FNU, therefore this COA was no longer triggered.

SMT met on 2/21/23 and agreed that turbidity will likely spike again in the afternoon, and DWR will have three days to comply to -2,000 cfs OMRI restriction if the daily average turbidity at OBI exceeds 12 FNU. The daily average turbidity at OBI was above 12 FNU on 2/21/23, triggering this COA for the third time this season. DWR exports were restricted to the SWP share of -2,000 OMRI from 2/24/23 through 2/26/23. On 2/26/23, the daily turbidity at OBI decreased to less than 12 FNU, therefore this COA was no longer triggered.

This COA was triggered for the fourth time this season by the conditions on 3/17/23. However, it was not controlling between 3/17/23 to 4/1/23.

This COA was off-ramped on 4/1/23.

8.5.2: This COA became active on 3/18/23 with Jersey Point exceeding 12°C and triggered due to 20mm Survey 4 recording Secchi depth < 1m. However, it has not been controlling and is unlikely to be controlling within the next seven days due to OMRI being highly positive.

As of 2/21/23, the federal agencies are following COA 8.5.2 per order 6(i) of the Interim Operations Plan (IOP).

8.12: This COA is not currently active due to water year type. The April water year type forecast is Wet. This COA is unlikely to become active this Water Year but will if the Water Year Type forecast is updated to dry or critical in May. As of 4/1/23, the LFS Protection portion has been off-ramped but DS Protection portion continues to be not active.

8.13: The Sacramento Valley Water Year Type Index (SVI) April forecast corresponding to the 50% probability of exceedance is 9.58 which is in the range for a Wet water year classification. The forecast was reported on the California Data Exchange Center (CDEC) [Water Supply Index Webpage](#), accessed on 04/11/23.

Section 3: Hydrology and Operations

Assessment of hydrologic, operational, and meteorological information. 8.1.5.2 A.

Section 3-A: Water operations conditions. 8.1.5.2.A. i

- Antecedent Actions: *(e.g. Delta Cross Channel [DCC] gate closure and actions such as integrated early winter pulse protection, etc.)*
 - DCC is closed as of 11/28/22.
- Controlling Factors: Limited real time demand
- Water Temperature:
 - Clifton Court Forebay (CCF) Daily Average Water Temperature = 18.16°C
 - 3 Station Average = 17.2°C
- Tidal Cycle: peak spring cycle with full moon on 5/5/23, heading into neap tide on 5/12/23.
- Turbidity:
 - 8.3.1 Freeport 3-day average = 16.94 formazin nephelometric units (FNU)
 - 8.5.1 Old River at Bacon Island (OBI) Turbidity = 6.7 FNU
- Salinity: X2 = ~ 59 km
- Hydrologic Footprint: Particle Tracking Model will be run when projections show San Joaquin River flow at Vernalis will drop below 5,000 cfs and off-ramp COA 8.4.3.

Section 3-B: Water operations outlook. 8.1.5.2.A. ii

- Outages
 - State Water Project (SWP): None
 - Central Valley Project (CVP): None
- Exports:
 - CCF: 3,500 cfs. Anticipated range: 2,500 to 6,680 cfs
 - Jones: 3,500 cfs. Anticipated range: 2,700 to 3,500 cfs
 - Combined: 5,200 to 10,180 cfs
- Meteorological Forecast: Cool and unsettled weather for the week with chances of showers, thunderstorms, and late season snow. High temperatures will remain 5-10 degrees below normal. Overall moisture beyond this weekend looks light.
- Six-day Storm Event Projection: n/a

Section 3-C: Projected conditions. 8.1.5.2.A. iii

- DCC Gates position: Scheduled to remain closed for season.
- Sacramento River flow at Freeport: 50,798 cfs as of 5/1/23. Anticipated range: 40,000 to 55,000 cfs
- San Joaquin River flow at Vernalis: 28,949 cfs as of 5/1/23. Anticipated range: 25,000 to 30,000 cfs
- Qwest: +35,462 cfs as of 4/30/23. Anticipated range: remain highly positive throughout the week.
- OBI Turbidity: 6.7 FNU
- NDOI: 76,246 cfs as of 4/30/23. Anticipated range: 65,000 to 80,000 cfs
- Upstream releases:
 - Keswick = 10,000 cfs. Anticipated range: 10,000 to 20,000 cfs.
 - Nimbus = 8,000 cfs. Anticipated range: 8,000 to 10,000 cfs.
 - Goodwin = 1,500 cfs. Anticipated range: 300 to 1,500 cfs.
 - Oroville = 20,000 cfs. Anticipated range: 10,000 to 20,000 cfs

Table 5: Comparison of OMR and OMR Index (5-day and 14-day averages for OMR Index and USGS gauge were reported on [SacPAS website](#), accessed 02 May 2023.

Date	Averaging Period	USGS gauges (cfs)	Index (cfs)
4/29/23	Daily	+9,740	+10,860
4/29/23	5-day	+8,910	+9,980
4/29/23	14-day	+7,860	+10,400

Section 4: Distribution and Biology.

8.1.5.2.B. Assessment of biological information for Delta Smelt and Longfin Smelt

Section 4-A: Delta Smelt population status 8.1.5.2.B. i

- EDSM: Two larval (FL: 6.5-6.8mm) DS were detected in the Confluence on 4/20/23 and Suisun Marsh on 4/19/23 (Table 2). Two larval (FL: 5.7-6.0mm) DS were detected in Suisun Bay on 4/12/23 and one larval (FL: 6.1mm) DS was detected in Grizzly Bay on 4/10/23. One larval (FL: 6.9mm) DS was detected in Suisun Bay on 4/5/23. One marked adult (FL: 83mm) was detected in the SDWSC on 3/21/23. Two marked adult (FL: 65-72mm) DS were detected in Suisun Marsh and SDWSC on 3/6/23 and 3/7/23 respectively. One marked adult (FL: 70mm) DS was detected in the Lower Sacramento River on 2/24/23. Three marked adult (FL: 76-79mm) DS were detected in the Confluence, Lower Sacramento River, and the SDWSC on 2/14/23, 2/15/23, and 2/17/23 respectively. One unmarked (FL: not measured) DS was detected in Suisun Marsh on 2/9/23. One unmarked adult (FL: 73mm) DS was detected in the Lower San Joaquin River on 1/31/23. Sixteen marked (FL: 47-80mm) DS were detected in Suisun Bay, Cache Slough, SDWSC, and Lower Sacramento from 1/24/23 to 2/7/23. One unmarked adult (FL: 71mm) DS was detected in the South Delta near Franks Tract on 1/17/23. One subadult DS (FL: 55mm) and one adult DS (FL: 62mm) were detected in Lower Sacramento River on 11/3/22 and 11/7/22 respectively.
- Chipps Island Trawl: One marked adult (FL: 84mm) DS was detected on 2/19/23. One marked DS (FL: 68mm) was detected on 1/19/23.
- Fall Mid-water Trawl (FMWT) Index for Delta Smelt: 0
- Delta Smelt life cycle model (LCM) discussion: n/a
- Biological Conditions: Turbidity remains below 12 FNU at Old River at Bacon Island. The average Secchi reading in 12 Central and South Delta stations from the last 20mm Survey is < 1m. X2 remains downstream, east of Martinez (~59 km). Spawning is on-going and larvae are present. There is a high degree of uncertainty regarding the response of cultured fish to environmental cues typically applied to wild DS. Distribution is widespread.
- % of population in Delta zones: n/a
- Smelt Larva Survey (SLS): SLS 6 detected (QC completed) three larval (FL: 5-12mm) DS near Chipps Island, Suisun Marsh, and Lower Sacramento River, and preliminarily detected (pending QC) one larval (FL: 10mm) DS in Honker Bay (Table 3).
- 20mm Survey:
 - Survey 1 detected (QC completed) six larval (FL: 6-12mm) DS near the Confluence, Suisun Bay, and Carquinez Strait (Table 4).
 - Survey 3 detected (QC completed) five larval (FL: 6-16mm) DS near the Confluence, Suisun Bay, Cache Slough, SDWSC, and Miner Slough, (Table 6).

- Survey 4 detected one larval (FL: 6mm) DS in the Confluence (Table 7). No DS were detected in the 12 Central and South Delta Stations, and the average Secchi depth there is 97cm.
- SKT: Survey 3 detected two marked DS (one ripe female, one unidentified sex) in SDWSC (station 719). Survey 2 detected two marked, ripe female DS in the Lower Sacramento River (station 704) and SDWSC (station 719).
- Salvage: No DS have been detected in salvage since 3/2/23. The cumulative seasonal salvage of adults is 52.
- FCCL lampara net sampling detected two adult DS (FL: ~60mm [estimated since fish were not directly handled]) in the Lower Sacramento River on 12/14/22. One fish was unmarked, and the other fish was tagged with red VIE tag (hard release) from the experimental release on 11/30/22.
- Experimental release: Approximately 13,000 cultured DS were released in the Sacramento Deepwater Shipping Channel on 1/25/23 and 1/26/23, 17,570 cultured DS were released in the Sacramento River near Rio Vista on 1/18/23 and 1/19/23, and 13,140 fish were released in the Sacramento River near Rio Vista on 11/30/22. No further experimental releases are scheduled for this water year.

Section 4-B: Longfin Smelt population status 8.1.5.2.B. ii.

- FMWT Index: 403
- EDSM: Four juvenile (FL: 39-41mm) LFS was detected in Grizzly Bay and San Pablo Bay during the 4/24/23 to 4/28/23 Kodiak Trawl (Table 1). 162 larval and juvenile (FL: 1-17.9mm) LFS were detected in Suisun Bay and Suisun Marsh during the 4/24/23 to 4/28/23 20mm Larval Surface Trawl (Table 2).
- Chipps Island Trawl: No LFS were detected during 4/24/23 to 4/28/23.
- 20-mm Survey:
 - Survey 2 preliminarily detected 102 new larval and juvenile (FL: 7-21mm) LFS in Carquinez Strait (Table 5).
 - Survey 3 detected one larval (FL: 6mm) LFS in the Lower Sacramento River, six larval (FL: 6-11mm) LFS in the Confluence, and 145 larval and juvenile (FL: 6-31mm) LFS in Suisun Bay and West Region (Table 6).
 - Survey 4 detected no LFS in the 12 Central and South Delta stations.
- SKT: Processing is ongoing. One juvenile (FL: 32mm) and one adult (FL: 90mm) LFS were detected in Suisun Marsh, and 52 juvenile (FL: 22-33mm) LFS were detected in Suisun Bay and West Region by SKT 4 (Table 8).
- Bay Study: In January, Bay Study detected six adults (FL: 87-109mm) and 44 sub-adult (FL: 58-84mm) LFS in stations ranging from the Lower Sacramento River to the South Bay. Distribution was widespread but overall, more downstream than in December.
- Salvage: No LFS have been detected in salvage since 3/2/23. The cumulative seasonal salvage of adults and sub-adults is 26.

Section 4-C: Additional data sources to assess sensitivity to entrainment Delta.8.1.5.2.C & D. i

Notes

- SKT 5 is on the water this week.
- Between SLS and 20mm Surveys thus far, there are 15 confirmed detections of larval DS, and one larval DS awaiting QC.
- EDSM Kodiak Trawl Phase 1 has concluded and will resume with the start of Phase 3 in July.

Attachments: Table 1: EDSM Kodiak Trawl Catch Table, Table 2: EDSM 20mm Larval Surface Trawl Summary Table, Table 3: SLS 6 Catch Table, Figure 1: Map of SLS, Table 4: 20mm Survey 1 Catch Table, Table 5: 20mm Survey 2 Catch Table, Table 6: 20mm Survey 3 Catch Table, Table 7: 20mm Survey 4 Catch Table, Figure 2: Map of 20mm Survey, Table 8: SKT 4 Catch Table, and Figure 3: Map of SKT.

Table 1: DS and LFS catch for EDSM 2023 Phase 1 Kodiak trawls (4/24/23 - 4/28/23). Only stations with catch of DS and LFS are reported here. These data are preliminary and subject to change. LFS that were mortalities upon capture were returned to the Lodi Fish and Wildlife Office (LFWO) to be frozen.

Date	Stratum	Subregion	Station Code	Species	Mark Type	Fork Length (mm)	Total Catch	Disposition
04/24/2023	Suisun Marsh	Grizzly Bay	23-39-SM01	LFS	None	39	1	LFWO
04/25/2023	Western Delta	East San Pablo Bay	23-39-WD02	LFS	None	41	1	LFWO
04/25/2023	Western Delta	East San Pablo Bay	23-39-WD02	LFS	None	38	2	Released

Table 2: All DS and LFS confirmed catch for EDSM 2023 Phase 2 20mm larval surface trawls by week and life stage or mark type (L=larvae, J=juvenile, A=adult, M=marked). Week 36 was April 3-7th, week 37 was April 10-14th, and week 38 was April 17-21st.

Survey Week	Stratum	# Sites	DSM Confirmed ID				LFS Confirmed ID		
			L	J	A	M	L	J	A
36	Suisun Bay	5	1	0	0	0	4	0	0
36	Suisun Marsh	5	0	0	0	0	0	3	0
36	Lower Sacramento	10	0	0	0	0	0	0	0
36	Cache Slough LI	5	0	0	0	0	0	0	0
36	Sac DW Ship Chan	5	0	0	0	0	1	0	0
36	Lower San Joaquin	10	0	0	0	0	0	0	0
37	Suisun Bay	10	2	0	0	0	60	1	0

Survey Week	Stratum	# Sites	DSM Confirmed ID				LFS Confirmed ID		
			L	J	A	M	L	J	A
37	Suisun Marsh	10	1	0	0	0	11	0	0
37	Lower Sacramento	5	0	0	0	0	1	0	0
37	Cache Slough LI	5	0	0	0	0	0	0	0
37	Sac DW Ship Chan	5	0	0	0	0	0	0	0
37	Lower San Joaquin	5	0	0	0	0	0	0	0
38	Suisun Bay	5	1	0	0	0	133	3	0
38	Suisun Marsh	5	1	0	0	0	26	0	0
38	Lower Sacramento	5	0	0	0	0	0	0	0
38	Cache Slough LI	10	0	0	0	0	0	0	0
38	Sac DW Ship Chan	5	0	0	0	0	0	0	0
38	Lower San Joaquin	10	0	0	0	0	0	0	0

Table 3: DS and LFS catch for SLS 6 (3/13/23 – 3/16/23). Only stations with catch of DS and LFS are reported here. These data are preliminary and subject to change. Samples that have been QC-ed are indicated with “Complete” ID Status whereas samples awaiting to be QC-ed are marked as “Preliminary”.

Year	Survey	SLS Station	Date	Turbidity (NTU)	Secchi (cm)	Sample Status	Species	Smelt Catch	ID Status	Min Length (mm)	Max Length (mm)	Mean Length (mm)	Yolk Sac (# of Individuals)
2023	6	306	3/16/2023	n/a	43	Processed	Longfin Smelt	2	Preliminary	8	9	8.5	0
2023	6	308	3/16/2023	16.6	47	Processed	Longfin Smelt	18	Preliminary	7	10	n/a	5
2023	6	322	3/16/2023	n/a	34	Processed	Longfin Smelt	8	Preliminary	8	10	8.8	4
2023	6	328	3/16/2023	35.8	36	Processed	Longfin Smelt	73	Preliminary	7	11	n/a	35
2023	6	336	3/16/2023	42.5	21	Processed	Longfin Smelt	168	Complete	6	19	7.9	43
2023	6	338	3/16/2023	15.8	49	Processed	Longfin Smelt	82	Preliminary	6	15	n/a	50
2023	6	342	3/17/2023	n/a	38	Processed	Longfin Smelt	87	Preliminary	7	19	n/a	23
2023	6	343	3/17/2023	n/a	19	Processed	Longfin Smelt	20	Preliminary	6	18	n/a	10
2023	6	344	3/17/2023	n/a	19	Processed	Longfin Smelt	4	Preliminary	6	18	7.8	2
2023	6	404	3/15/2023	30.7	28	Processed	Longfin Smelt	24	Preliminary	6	19	9.5	8
2023	6	405	3/15/2023	60.3	18	Processed	Longfin Smelt	18	Complete	6	9	7.4	6
2023	6	411	3/15/2023	39.3	41	Processed	Longfin Smelt	14	Complete	6	8	6.8	9
2023	6	418	3/15/2023	35.5	36	Processed	Longfin Smelt	12	Complete	6	10	7.8	4
2023	6	501	3/15/2023	21.1	38	Processed	Longfin Smelt	3	Complete	5	7	6.0	0
2023	6	504	3/15/2023	31.8	34	Processed	Longfin Smelt	31	Complete	6	7	6.4	16
2023	6	508	3/13/2023	36.0	38	Processed	Delta Smelt	1	Complete	5	5	5.0	0
2023	6	508	3/13/2023	36.0	38	Processed	Longfin Smelt	11	Complete	6	8	7.0	5
2023	6	513	3/13/2023	36.0	49	Processed	Longfin Smelt	5	Complete	6	7	6.6	2
2023	6	519	3/15/2023	43.9	33	Processed	Delta Smelt	1	Preliminary	10	10	10.0	0
2023	6	519	3/15/2023	43.9	33	Processed	Longfin Smelt	3	Complete	6	8	7.0	1
2023	6	520	3/13/2023	18.0	57	Processed	Longfin Smelt	5	Complete	6	7	6.4	2
2023	6	602	3/15/2023	38.9	40	Processed	Longfin Smelt	17	Complete	6	9	6.8	13

Year	Survey	SLS Station	Date	Turbidity (NTU)	Secchi (cm)	Sample Status	Species	Smelt Catch	ID Status	Min Length (mm)	Max Length (mm)	Mean Length (mm)	Yolk Sac (# of Individuals)
2023	6	606	3/15/2023	85.5	17	Processed	Longfin Smelt	14	Complete	7	18	11.8	0
2023	6	609	3/15/2023	35.3	38	Processed	Longfin Smelt	3	Complete	7	9	8.3	1
2023	6	610	3/15/2023	17.7	50	Processed	Delta Smelt	1	Complete	12	12	12.0	0
2023	6	610	3/15/2023	17.7	50	Processed	Longfin Smelt	3	Complete	6	7	6.7	3
2023	6	704	3/13/2023	15.7	44	Processed	Longfin Smelt	6	Complete	6	9	7.3	2
2023	6	705	3/13/2023	19.4	44	Processed	Longfin Smelt	2	Complete	7	8	7.5	0
2023	6	706	3/13/2023	19.8	42	Processed	Longfin Smelt	3	Complete	6	7	6.3	2
2023	6	706	3/13/2023	19.8	42	Processed	Delta Smelt	1	Complete	6	6	6.0	1
2023	6	707	3/13/2023	18.6	35	Processed	Longfin Smelt	5	Preliminary	6	10	7.0	2
2023	6	711	3/13/2023	19.9	25	Processed	Longfin Smelt	1	Complete	8	8	8.0	0
2023	6	723	3/13/2023	32.7	37	Processed	Longfin Smelt	1	Complete	8	8	8.0	0
2023	6	801	3/13/2023	15.8	53	Processed	Longfin Smelt	2	Complete	6	8	7.0	2
2023	6	804	3/13/2023	n/a	46	Processed	Longfin Smelt	1	Complete	6	6	6.0	1
2023	6	812	3/13/2023	26.6	42	Processed	Longfin Smelt	2	Complete	8	10	9.0	0
2023	6	815	3/13/2023	29.3	37	Processed	Longfin Smelt	7	Complete	6	8	7.1	3
2023	6	901	3/13/2023	18.5	54	Processed	Longfin Smelt	3	Complete	5	7	5.7	2




Figure 1: Map of SLS station

Table 4: DS and LFS catch for 20mm Survey 1 (3/13/23 – 3/16/23). Only stations with catch of DS and LFS are reported here. These data are preliminary and subject to change. Samples that have been QC-ed are indicated with “Complete” ID Status whereas samples awaiting to be QC-ed are marked as “Preliminary”.

Year	Survey	Station	Date	Secchi (cm)	# Tows Processed	Species	Total Catch	ID Status	Min Length (mm)	Max Length (mm)	Mean Length (mm)
2023	1	306	3/13/23	n/a	1	Longfin Smelt	1	Preliminary	6	6	6
2023	1	308	3/13/23	36	3	Longfin Smelt	6	Preliminary	9	18	n/a
2023	1	323	3/15/23	22	3	Longfin Smelt	31	Preliminary	8	19	n/a
2023	1	330	3/13/23	n/a	2	Longfin Smelt	10	Preliminary	7	8	n/a
2023	1	335	3/13/23	43	1	Longfin Smelt	1	Preliminary	19	19	19
2023	1	336	3/15/23	29	2	Longfin Smelt	10	Preliminary	7	21	12.9
2023	1	338	3/15/23	30	2	Longfin Smelt	3	Preliminary	9	87	n/a
2023	1	401	3/14/23	50	3	Longfin Smelt	6	Preliminary	7	18	11.5
2023	1	404	3/15/23	30	3	Longfin Smelt	59	Complete	7	22	17.4
2023	1	405*	3/15/23	24	3	Delta Smelt	2	Complete	9	12	10.5
2023	1	405*	3/15/23	24	3	Longfin Smelt	7	Complete	9	20	17.1
2023	1	411	3/17/23	33	3	Delta Smelt	2	Complete	6	6	6.0
2023	1	411	3/17/23	33	3	Longfin Smelt	2	Complete	6	7	6.5
2023	1	504	3/16/23	25	3	Longfin Smelt	2	Complete	7	22	14.5
2023	1	602	3/16/23	40	3	Longfin Smelt	12	Complete	9	22	16.3
2023	1	606	3/16/23	37	3	Longfin Smelt	19	Complete	15	23	18.6
2023	1	801	3/15/23	20	3	Longfin Smelt	2	Complete	7	20	13.5
2023	1	804	3/15/23	32	3	Delta Smelt	2	Complete	6	10	8.0
2023	1	804	3/15/23	32	3	Longfin Smelt	3	Complete	6	7	6.3
2023	1	706	3/17/23	21	3	Longfin Smelt	1	Complete	6	6	6.0
2023	1	901	3/13/23	50	3	Longfin Smelt	1	Complete	7	7	7.0

Table 5: DS and LFS catch for 20mm Survey 2 (3/27/23 – 3/30/23). Only stations with catch of DS and LFS are reported here. These data are preliminary and subject to change. Samples that have been QC-ed are indicated with “Complete” ID Status whereas

samples awaiting to be QC-ed are marked as “Preliminary”.

Year	Survey	Station	Date	Secchi (cm)	# Tows Processed	Species	Total Catch	ID Status	Min Length (mm)	Max Length (mm)	Mean Length (mm)
2023	2	401	3/30/23	38	2	Longfin Smelt	102	Preliminary	7	21	n/a
2023	2	404	3/28/23	42	3	Longfin Smelt	25	Preliminary	9	25	17.6
2023	2	405	3/28/23	38	3	Longfin Smelt	17	Preliminary	6	13	9.5
2023	2	411	3/28/23	39	3	Longfin Smelt	15	Preliminary	16	25	19.3

Table 6: DS and LFS catch for 20mm Survey 3 (4/10/23 – 4/14/23). Only stations with catch of DS and LFS are reported here. These data are preliminary and subject to change. Samples that have been QC-ed are indicated with “Complete” ID Status whereas samples awaiting to be QC-ed are marked as “Preliminary”.

Year	Survey	Station	Date	Secchi (cm)	# Tows Processed	Species	Total Catch	ID Status	Min Length (mm)	Max Length (mm)	Mean Length (mm)
2023	3	322	4/13/23	48	2	Longfin Smelt	1	Preliminary	27	27	27
2023	3	338	4/14/23	58	1	Longfin Smelt	3	Preliminary	12	13	12.5
2023	3	401	4/11/23	45	3	Longfin Smelt	135	Preliminary	8	31	n/a
2023	3	501	4/12/23	43	3	Delta Smelt	1	Complete	7	7	7.0
2023	3	501	4/12/23	43	3	Longfin Smelt	1	Complete	7	7	7.0
2023	3	504	4/13/23	54	3	Longfin Smelt	5	Complete	6	6	6.0
2023	3	508	4/13/23	50	3	Longfin Smelt	5	Complete	6	11	7.6
2023	3	801	4/13/23	78	2	Longfin Smelt	1	Complete	6	6	6.0
2023	3	704	4/10/23	49	3	Delta Smelt	1	Complete	7	7	7.0
2023	3	706	4/10/23	49	3	Longfin Smelt	1	Complete	7	7	7.0
2023	3	716	4/11/23	68	3	Delta Smelt	1	Complete	6	6	6.0
2023	3	719	4/11/23	29	2	Delta Smelt	1	Complete	16	16	16.0
2023	3	726	4/11/23	58	3	Delta Smelt	1	Complete	12	12	12.0

Table 7: DS and LFS catch for 20mm Survey 4 (4/24/23 – 4/28/23). Only stations with catch of DS and LFS are reported here. These data are preliminary and subject to change. Samples that have been QC-ed are indicated with “Complete” ID Status whereas samples awaiting to be QC-ed are marked as “Preliminary”.

Year	Survey	Station	Date	Secchi (cm)	# Tows Processed	Species	Total Catch	ID Status	Min Length (mm)	Max Length (mm)	Mean Length (mm)
2023	4	804	4/24/23	80	3	Delta Smelt	1	Complete	6	6	6.0




Figure 2: Map of 20mm Survey stations

Table 8: DS and LFS catch for SKT 4 (4/3/23 – 4/7/23). Only stations with catch of DS and LFS are reported here. These data are preliminary and subject to change.

Year	Sample Date	Survey Number	Station Code	Catch	Species	Fork Length (mm)
2023	4/7/2023	4	340	50	LFS	22
2023	4/7/2023	4	340	50	LFS	23
2023	4/7/2023	4	340	50	LFS	23
2023	4/7/2023	4	340	50	LFS	24
2023	4/7/2023	4	340	50	LFS	24
2023	4/7/2023	4	340	50	LFS	24
2023	4/7/2023	4	340	50	LFS	25
2023	4/7/2023	4	340	50	LFS	25
2023	4/7/2023	4	340	50	LFS	25
2023	4/7/2023	4	340	50	LFS	25
2023	4/7/2023	4	340	50	LFS	26
2023	4/7/2023	4	340	50	LFS	26
2023	4/7/2023	4	340	50	LFS	26
2023	4/7/2023	4	340	50	LFS	26
2023	4/7/2023	4	340	50	LFS	26
2023	4/7/2023	4	340	50	LFS	26
2023	4/7/2023	4	340	50	LFS	26
2023	4/7/2023	4	340	50	LFS	27
2023	4/7/2023	4	340	50	LFS	27
2023	4/7/2023	4	340	50	LFS	27
2023	4/7/2023	4	340	50	LFS	27
2023	4/7/2023	4	340	50	LFS	27
2023	4/7/2023	4	340	50	LFS	27
2023	4/7/2023	4	340	50	LFS	28
2023	4/7/2023	4	340	50	LFS	28
2023	4/7/2023	4	340	50	LFS	28

Year	Sample Date	Survey Number	Station Code	Catch	Species	Fork Length (mm)
2023	4/7/2023	4	340	50	LFS	28
2023	4/7/2023	4	340	50	LFS	28
2023	4/7/2023	4	340	50	LFS	28
2023	4/7/2023	4	340	50	LFS	28
2023	4/7/2023	4	340	50	LFS	28
2023	4/7/2023	4	340	50	LFS	28
2023	4/7/2023	4	340	50	LFS	28
2023	4/7/2023	4	340	50	LFS	28
2023	4/7/2023	4	340	50	LFS	28
2023	4/7/2023	4	340	50	LFS	28
2023	4/7/2023	4	340	50	LFS	29
2023	4/7/2023	4	340	50	LFS	29
2023	4/7/2023	4	340	50	LFS	29
2023	4/7/2023	4	340	50	LFS	30
2023	4/7/2023	4	340	50	LFS	30
2023	4/7/2023	4	340	50	LFS	30
2023	4/7/2023	4	340	50	LFS	30
2023	4/7/2023	4	340	50	LFS	30
2023	4/7/2023	4	340	50	LFS	30
2023	4/7/2023	4	340	50	LFS	30
2023	4/7/2023	4	340	50	LFS	31
2023	4/7/2023	4	340	50	LFS	31
2023	4/7/2023	4	340	50	LFS	32
2023	4/7/2023	4	340	50	LFS	33
2023	4/7/2023	4	340	50	LFS	33
2023	4/7/2023	4	405	1	LFS	28
2023	4/7/2023	4	411	1	LFS	26
2023	4/7/2023	4	606	1	LFS	32
2023	4/7/2023	4	610	1	LFS	90

Spring Kodiak Trawl Sampling Stations




Figure 3: Map of SKT stations