

Natural Resource Damage Assessment and Restoration Considerations: Ephemeral Data Collection Plan for a Northern California River Spill That Also Impacts Coastal Resources



Photo Credits: Smith River Alliance



Photo Credits: Del Norte County



Photo Credits: Del Norte County



Photo Credits: Smith River Alliance

September 2023

**California Department of Fish and Wildlife
Office of Spill Prevention and Response
West Sacramento, California**

**National Park Service
Fort Collins, Colorado**

**U.S. Department of the Interior
Office of Environmental Policy &
Compliance
Sacramento, California**

**Prepared with support from:
Industrial Economics, Incorporated and Research Planning, Inc.**

[this page intentionally left blank]

Table of Contents

1.0	Introduction	1
1.1	Purpose, Goals, and Objectives of this Plan	1
1.2	Safety	1
1.3	Communication Among Agency Representatives	2
1.4	Relationship and Communication with the Unified Command	2
1.5	Field Teams	4
1.6	Special Requirements for Response in Wild and Scenic Rivers and Federally Designated Wilderness Areas	4
2.0	Oil Spill Conceptual Models	5
2.1	Oil Fate and Behavior: Medium Crude Oil and Diesel Fuel Spills to Water	5
2.2	Conceptual Model: Northern California River Spill That Also Impacts Coastal Resources.....	7
3.0	Ephemeral Data Collection Procedures	10
3.1	Overview	10
3.2	Source Oil/Oil.....	11
3.3	Sheen	11
3.4	Water	11
3.5	Tissue	12
3.6	Sediment.....	12
3.7	Human Use/Recreational Use.....	12
3.8	Benthic Macroinvertebrates in Rivers	13
3.9	Field Documentation and Photography.....	13
3.10	Chain of Custody	14
3.11	Chemical Analysis Guidelines.....	15
4.0	Natural Resource Trustee Representatives Contact Information (current as of July 2023).....	16
4.1	CA Department of Fish and Wildlife, Office of Spill Prevention and Response.....	16
4.2	U.S. Fish and Wildlife Service.....	16
4.3	U.S. Department of the Interior.....	16
4.4	National Oceanic and Atmospheric Administration, Office of Response and Restoration	16
4.5	U.S. Department of Agriculture	16
5.0	Technical Support And Chemical Analysis Contact Information.....	17
5.1	U.S. Environmental Protection Agency.....	17

5.2	CA Department of Fish and Wildlife – Office of Spill Prevention and Response Bioassessment Team.....	17
5.3	Chemical Analysis/Technical Support	17
6.0	Other State and Local Agency Contacts.....	17
6.1	CA Department of Fish and Wildlife – Regional Office	17
6.2	CA Department of Parks and Recreation.....	18
6.3	CA Regional Water Quality Control Board.....	18
6.4	Office of Environmental Health Hazard Assessment.....	18
6.5	California Coastal Commission	19
6.6	California State Lands Commission.....	19
6.7	Public Utilities	19
6.8	Railroads	19
6.9	Tribal Contacts	19
7.0	References Cited	19

Appendices:

APPENDIX 1: Detailed Standard Operating Procedures: Source Oil and Oil

APPENDIX 2: Detailed Standard Operating Procedures: Sheen

APPENDIX 3: Detailed Standard Operating Procedures: Water

APPENDIX 4: Detailed Standard Operating Procedures: Tissue

APPENDIX 5: Detailed Standard Operating Procedures: Sediment

APPENDIX 6: Detailed Standard Operating Procedures: Human Use/Recreational Use

APPENDIX 7: Detailed Standard Operating Procedures: Benthic Macroinvertebrates in Rivers

APPENDIX 8: Detailed Standard Operating Procedures: Field Documentation Using Photography

APPENDIX 9: Detailed Standard Operating Procedures: Chain of Custody and Sample Collection Summary Form

APPENDIX 10: Example Coordination Agreements

APPENDIX 11: Resource-Specific Work Plan Examples.

APPENDIX 12: Links to Oil Spill Job Aids and Data Sets on Baseline Resource Condition

List of Figures

Figure 1. Spill response Unified Command Structure, including communication and coordination points with NRDAR.....	3
Figure 2. Plots of oil fate using GNOME for the release of 500 barrels of a medium crude oil (A) and a diesel fuel (B). Refer to the text for the model inputs. Note the high amounts of evaporation and dispersion for diesel fuel versus a medium crude oil.....	6

List of Tables

Table 1. Conceptual model for a northern California river spill that also impacts coastal resources (e.g., Smith River).....	8
Table 2. Timeline for ephemeral data collection for Northern California rivers and estuaries.	10
Table 3. Key sampling requirements for each sample type.	15
Table 4. Analytes and analysis methods by matrix.....	15

List of Abbreviations and Acronyms

AWB	Auto White Balance
BMI	benthic macroinvertebrates
BTEX	benzene, toluene, ethylbenzene, and xylene
CDFW-OSPR	California Department of Fish and Wildlife, Office of Spill Prevention and Response
COC	Chain of Custody
DOI	Department of the Interior
EDCC	Ephemeral Data Collection Coordinator
EPA	U.S. Environmental Protection Agency
GNOME	General NOAA Oil Modeling Environment
GPS	global positioning system
HAZWOPER	Hazardous Waste Operations & Emergency Response
L	liter
mL	milliliter
MU	mussel
NOAA	National Oceanic and Atmospheric Administration
NPFC	National Pollution Fund Center
NPFC	National Pollution Fund Center
NPS	National Park Service
NRDAR	Natural Resource Damage Assessment and Restoration
PAH	polycyclic aromatic hydrocarbons
PFD	personal floatation device
PIANO	paraffins, iso-paraffins, aromatics, naphthenes, and olefins
PPE	personal protective equipment
RI	Rocky Intertidal
RP	Responsible Party
SCAT	Shoreline Cleanup Assessment Technique
SLR	single lens reflex
SOP	Standard Operating Procedure
SWAMP	Surface Water Ambient Monitoring Program
the Plan	Ephemeral Data Collection Plan
TPH	total petroleum hydrocarbons
"Trustees"	Natural Resource Trustees

UAS	using uncrewed aerial systems
USDA	U.S. Department of Agriculture
USFWS	United States Fish and Wildlife Service
WCJAT	West Coast Joint Assessment Team
WSR	Wild and Scenic Rivers

Executive Summary

This Ephemeral Data Collection Plan (the Plan) was developed in collaboration between Natural Resource Damage Assessment Trustees with funding from the Department of the Interior's (DOI) Inland Oil Spill Preparedness Project. The Plan describes and prioritizes ephemeral data collection in the event of an accidental release of oil that affects a Northern California river and estuary habitats. Ephemeral data are information that can only be collected within a narrow timeframe after a spill occurs—before that information is lost. This effort is intended to be a “general template” applicable to a broader habitat setting.

The Natural Resources Damage Assessment and Restoration (NRDAR) team will collect ephemeral data. The team may consist of representatives of the responsible party (RP) and the natural resource trustees (hereafter referred to as the ‘Trustees’), including but not limited to the California Department of Fish and Wildlife, Office of Spill Prevention and Response (CDFW-OSPR); U.S. Department of the Interior Bureaus including the U.S. Fish and Wildlife Service (USFWS), the National Park Service (NPS), the Bureau of Land Management and/or other DOI Bureaus; the National Oceanic and Atmospheric Administration (NOAA); and the U.S. Department of Agriculture (U.S. Forest Service). State Trustee representatives from the California Department of Parks and Recreation and the California State Lands Commission may also contribute to the implementation of the Plan.

The purpose of this Plan is to enhance and increase preparedness for effective implementation of ephemeral data collection for NRDAR of a spill that affects Northern California river and estuary habitats. NRDAR is dependent on successful collection of ephemeral data during an accidental release. Thus, a well-designed ephemeral data collection plan that can be implemented in a timely way is key. Ephemeral data consist of many types of information needed to determine immediate effects of a release on the surrounding environment and resources, including petroleum hydrocarbon concentrations in environmental media, impacts to biota from released oil, or changes in recreational use. To this end, the Plan describes a conceptual model of the behavior, fate, impacts, and resources at risk from spills of a medium crude oil or a diesel fuel. This Plan also includes detailed procedures for collection of ephemeral data for different media (source oil, oil, surface water, sediment, and tissue samples) and resources (human/recreational use and benthic macroinvertebrates) in the event of an accidental release of petroleum into Northern California river and estuary habitats. Several components important to NRDAR, such as procedures for fish collections and bird necropsies, are not included because they are not relevant to the resources at risk or because they require specialized equipment and training. Therefore, this Plan has representative NRDAR ephemeral data considerations but may not be comprehensive for all NRDAR needs. Finally, this Plan provides guidance on working with the response organizations and co-trustees during a spill. Proper coordination with the multiple parties involved in a spill response and NRDAR will be critical for safe and effective collection of ephemeral data, without negatively affecting the incident response. Having this Plan in place prior to a release will facilitate the collection of critical environmental information during the early stages of a response effort.

The goals of the Plan are to collect ephemeral data that will (1) assist in documenting the source of oil(s); (2) document petroleum hydrocarbon concentrations in water, sediment, and selected freshwater organisms; and (3) document impacts to human/recreational users and benthic macroinvertebrates prior to and following an accidental release of oil in Northern California river and estuary habitats.

Before implementing this Plan, representatives of the NRDAR team should determine whether incident circumstances warrant implementation of the Plan, and, if so, any modifications to the Plan that may be required. Also, during an incident, the ephemeral data collection procedures and locations detailed in this Plan may be modified following lessons learned from future oil spill response drills or real-time

responses to accidental releases within Northern California river and estuary habitats. Any lessons learned should prompt periodic updates to this Plan and ideally, at a minimum, annual reviews should be conducted to maintain maximum value.

Finally, the Trustees recognize that other (non-petroleum) spills may occur in Northern California river and estuary habitats and may benefit from the sampling design and procedures described in this Plan. In the event of a release, the Trustees may choose to implement all or parts of this Plan or to modify it to assist them during NRDAR pre-assessment and assessment activities.

1.0 Introduction

1.1 Purpose, Goals, and Objectives of this Plan

Purpose: Enhance and increase preparedness for effective implementation of ephemeral data collection for natural resource damage assessment and restoration (NRDAR) of a spill that affects Northern California river and estuary habitats. The Plan includes: (1) A conceptual model of the behavior, fate, impacts, and resources at risk from spills of a medium crude oil and a diesel fuel that is used to guide the ephemeral data collection plan; (2) Detailed procedures for collection of ephemeral data for media (source oil, surface water, sediment, and tissue samples) and resources (human/recreational use and benthic macroinvertebrates) in the event of an accidental release of petroleum into Northern California river and estuary habitats; and (3) Guidance on working with the response organizations and co-trustees during a spill.

Goals: To obtain data that will assist in determining the source of oil(s); document petroleum hydrocarbon concentrations in water, sediment, and selected freshwater organisms; and document impacts to human/recreational users and benthic macroinvertebrates prior to and following an accidental release of oil in Northern California river and estuary habitats.

Objectives: (1) To collect source oil, water, sediment, and tissue samples within the first hours, days, and weeks after an accidental release of oil for petroleum hydrocarbon analysis; (2) Collect data on human/recreational use and benthic macroinvertebrate abundances and species composition. During an incident, there is a narrow window of opportunity for collection of these data, and, accordingly, they are referred to as “ephemeral” data (i.e., if the samples are not collected, the opportunity to collect them will be lost permanently). Ephemeral and baseline data can be critical in identifying the need for, and scope of, subsequent environmental sampling and injury assessment.

1.2 Safety

Safety is the most important consideration in plan implementation. Field teams may encounter oil during collection of water, sediment or tissue samples. Personnel collecting data in the field should be at least 24-hour Hazardous Waste Operations & Emergency Response (HAZWOPER) certified if oil is present (unless their employer policies require additional safety training) and have received permission from the Unified Command Incident Site Safety Officer to enter impacted areas.

Before sampling in impacted areas, all field team members must read and be familiar with and follow the procedures specified in the site safety plan prepared by the Unified Command for the incident unless the NRDAR team has specified a different safety plan or safety procedures. Generally, the safety plan should require that, before going into the field, all field team members will receive a daily safety briefing from the Ephemeral Data Collection Coordinator (EDCC; discussed in detail in Section 1.4). Field team members collecting samples by boat, if applicable, will receive a boat safety briefing by the boat operator prior to leaving. When on or near water, field team members will wear personal floatation devices (PFDs). Good judgment must be used at all times, particularly when considering fieldwork during inclement weather and collecting samples near cliffs and in high-flow situations. If sampling is to be conducted where high flows create dangerous swift water conditions, field team members must survey the river for features (e.g., access points, eddies) and hazard risks (e.g., strainers, rocks) and wear appropriate river safety gear (e.g., PFD, helmet, whistle, and knife) prior to entering the river. Shore-based field team members also should be equipped with the same safety gear, including a throw bag, should a rescue scenario arise. No sampling will be conducted in the dark. While working near the river, field team members should be mindful of slippery surfaces (e.g., rocks) and sharp objects. Field team

members should wear sunglasses, sunscreen, appropriate footwear, and other personal protective equipment (PPE) as required by the safety officer.

PPE will depend on the specific hazardous petroleum materials and their concentrations. Under no circumstances should the field team enter uncharacterized, freshly impacted riverbanks or coastal estuary shorelines without proper training (i.e., 40-hour HAZWOPER certification and current 8-hour refresher) and combustible gas/hydrogen sulfide meters. Likewise, sampling in the impacted area will not be conducted if respirators are required or the safety officer deems the area unsafe. When collecting water samples in the impacted area, field sampling team members will wear appropriate PPE (e.g., gloves, Tyvek, PFD). Nitrile gloves and/or cut-resistant gloves will be worn when sampling any medium of interest and will be changed between each sampling site.

1.3 Communication Among Agency Representatives

Communications should be as follows when possible, although will vary according to the specifics of the incident. Ideally, after receiving notification of a release of greater than *de minimis* quantities, the Trustees and the RP will notify, to the extent feasible, the key RP and Natural Resource Trustee Agency Lead(s) or their alternates (Section 4). If a cooperative assessment with the RP is possible, the RP and Trustee Agency representatives may coordinate to assess the circumstances and determine sampling priorities including what, if any, elements of the Plan should be implemented, modified, or if additional monitoring elements should be considered. Key contacts for mobilizing field teams are identified in Section 4. If the RP and Trustee Agency Lead(s) are unreachable in the early hours/days of the spill, CDFW-OSPR Agency Lead(s) will begin coordinating initial sampling priorities and efforts, including implementation of this Plan.

In the case of a cooperative assessment, the RP, Trustee Agency Lead(s), or their alternates (Section 4), may participate in an initial conference call to determine the specific plan elements to be implemented, to share needed contact and location information, and then to assign their respective staffs to participate in sample collection activities as appropriate. A Trustee EDCC, appointed by the NRDAR Agency Lead(s), will be assigned to provide project oversight and management. The EDCC will manage implementation of this Plan and coordinate with the Unified Command for the response (see Section 1.4), via the NRDAR Representative, for increased efficiencies in all aspects of data collection for the response and the damage assessment.


1.4 Relationship and Communication with the Unified Command

The NRDAR for an incident is done in parallel with the incident response but is separate from it. The goals of response and NRDAR are different. The goals of spill response are to stop and stabilize the source of the spill, remove oil from the environment, protect the safety and health of the responders and the public, and avoid or minimize harm to the environment. The goal of NRDAR is to identify the type and amount of restoration needed to restore injured natural resources.

Figure 1 details the response Incident Command Structure organization for a spill and the coordination points with NRDAR. Since NRDAR field assessment activities may overlap those of the response, close coordination and cooperation between the two efforts is necessary. The NRDAR Agency Lead(s) are responsible for establishing the communication link with the Incident Command (Unified Command if the incident is federalized) via the NRDAR Representative or Liaison as described in the West Coast Joint Assessment Team (WCJAT) guidance document (WCJAT, 2017) and the CDFW-OSPR Policy 603-1, Communication and Coordination between Natural Resource Damage Assessment and Restoration (NRDAR) and Incident Command Structure During Spill Response (2009). All communications with the Unified Command will be coordinated through the NRDAR Representative.

Indicates NRDAR – ICS Coordination Need

Indicates Primary NRDAR Contact Point



* Includes communication and coordination points with NRDAR

Figure 1. Spill response Unified Command Structure, including communication and coordination points with NRDAR.

The EDCC will prepare a General Message (*ICS 213*) to establish and document communications and resource requests between NRDAR and the Planning Section (Environmental Unit) and the Operations Section (Wildlife Branch) of the Unified Command. This will allow for the coordination of environmental sampling and field data collection activities between NRDAR and the response. For example, when NRDAR ephemeral data collection field teams are directed to sample in impacted areas, health and safety must be addressed and permission to enter impacted areas must be coordinated through the NRDAR Representative and the EDCC. A General Message (*ICS 213*) will be provided to the Unified Command daily throughout the duration of the field activities or as appropriate based on the incident.

1.5 Field Teams

The EDCC makes field assignments, reviews and disseminates health and safety procedures, monitors all field activities, and provides the field team leaders with information containing specific guidelines (e.g., health and safety, sampling locations and sites, sample collection procedures, etc.) for implementing the Plan. All personnel involved in implementation of the ephemeral data collection plan are responsible for reporting progress and results to the EDCC. The EDCC also will verify that all field sampling team members have read and signed a copy of the incident health and safety plan.

NRDAR field teams, including representatives of the RP and/or the Trustees, may collect samples both on water and land. To expedite sample collection, separate water (i.e., in wadable or non-wadable river reaches) and land field teams may be mobilized. Water-based sampling teams may sample from a boat in deeper waters and include an experienced boat operator and at least two other qualified people (i.e., people with experience or training for on-water sample collection): one to collect samples and the other to record notes and take photographs. Depending on the extent of planned collection and available personnel, on-water sampling teams may include up to four data collection personnel: one to collect water and sediment samples, one to collect tissue, one to record notes, and one to photo-document the riverbank. Source oil sample collection will usually be undertaken by a person designated by the Unified Command or Investigations. All communications with the Unified Command will be coordinated through the NRDAR Representative (Section 1.4). Key contacts for mobilizing field teams are identified in Section 4.

1.6 Special Requirements for Response in Wild and Scenic Rivers and Federally Designated Wilderness Areas

Wild and Scenic Rivers (WSR) are those designated as part of the National Wild and Scenic Rivers System, created by Congress in 1968 (PL 90-542;16 U.S.C. 1271 et seq.) to preserve certain rivers with outstanding natural, cultural, and recreational values in a free-flowing condition for the enjoyment of present and future generations. The Wild and Scenic Rivers Act charges administration of rivers in the National Wild and Scenic Rivers System to four federal land management agencies (Bureau of Land Management, National Park Service, U.S. Fish and Wildlife Service, and U.S. Forest Service). The Act allows for designation of segments of rivers, including tributaries. Boundaries of a WSR are typically one-quarter mile on either bank (lower 48) or one-half mile (Alaska) on either bank of the river.

The National Wilderness Preservation System is a network of more than 800 designated wilderness areas co-managed by the National Park Service, Bureau of Land Management, U.S. Fish and Wildlife Service, and U.S. Forest Service. The primary mandate of the Wilderness Act (16 U.S.C. 1131-1136) is to preserve wilderness character—the natural, untamed, undeveloped, and primitive aspects that make wilderness worthy of its name. The Act states that:

“...within any wilderness area designated by this Act and, except as necessary to meet *minimum requirements* for the administration of the area for the purpose of this Act (including measures required in emergencies involving the health and safety of persons within the area), there shall

be no temporary road, no use of motor vehicles, motorized equipment or motorboats, no landing of aircraft, no other form of mechanical transport, and no structure or installation within any such area.”

If a spill affects a WSR or federal wilderness area and its boundaries, the administrating agency must be contacted immediately. Special use permits may be required for emergency access to the WSR and its boundaries, including for Shoreline Cleanup Assessment Technique (SCAT)-type surveys, wildlife rescue, ephemeral data collection, and other response-related activities. Routine response and damage assessment activities may need to be adjusted for consistency with protections afforded WSRs or wilderness areas. The agency Point of Contact can assist with identifying the permitting entity and facilitate the process. Furthermore, response actions will need to have close monitoring and adherence to best management practices so that the character of the WSR or wilderness area is not degraded.

2.0 Oil Spill Conceptual Models

2.1 Oil Fate and Behavior: Medium Crude Oil and Diesel Fuel Spills to Water

The type of oil determines how the oil will behave once released to water, its fate and transport, and its likely impacts. For this Plan, information on two oil types is presented: a medium crude oil and a diesel fuel. The oil fate component of the General NOAA Operational Modeling Environment (GNOME; available at <https://gnome.orr.noaa.gov>) was used to generate plots of the fate of: 1) Dos Cuadras crude oil (a California crude oil from offshore in the Santa Barbara Channel) with an API of 25.6 and a pour point of -30°C, and 2) a diesel fuel with an API of 38.7.

The model was run using the following input:

- Spill volume: 500 barrels
- Water temperature: 65°F
- Wind speed: 5 knots
- Wave height: 1 foot
- Horizontal diffusion: 1×10^3 centimeter²/second (an order of magnitude lower than open water spills to account for less spreading in a river)

It should be noted that GNOME does not include inputs for turbulent flow in rivers, such as over rapids; therefore, the plots of the fate of each oil in Figure 2 are intended to show the relative behavior of these two types of oil under the modeled conditions. The actual fate of spills in turbulent water is likely to be characterized by higher natural dispersion and less loss by evaporation. The comparative fate, behaviors, and effects for each oil type are summarized below.




Figure 2. Plots of oil fate using GNOME for the release of 500 barrels of a medium crude oil (A) and a diesel fuel (B). Refer to the text for the model inputs. Note the high amounts of evaporation and dispersion for diesel fuel versus a medium crude oil.

Medium crude oil:

- Is a mixture of a wide range of compounds, from light to heavy and tends to increase in viscosity as it weathers;
- Has of loss by evaporation usually up to 30% in the first day after the release;
- Tends to spread into thick slicks that can coat surfaces and foul animals and vegetation;
- Fresher oil can penetrate porous sediments; and
- Poses a moderate risk to aquatic organisms from the dissolved fraction.

Diesel fuel

- Is a refined product with a narrow range of lighter compounds and very low viscosity;
- Can have high rates of loss by evaporation (up to 80%) in the first day after a release;
- Has high rates of natural dispersion because of its low viscosity;
- Poses a high risk to aquatic resources because of high amounts of dissolved and dispersed fractions that get mixed into the water column, especially as the oil flows over rapids and falls;
- Spreads into thin sheens;
- Coating of surfaces and vegetation tends to be thin and decreases over time; and
- Only thicker accumulations against the shoreline will penetrate into and persist in porous sediments.

These two oils are used in the conceptual model discussed in the following section.

2.2 Conceptual Model: Northern California River Spill That Also Impacts Coastal Resources

The conceptual model in Table 1 is based on a release of a medium crude oil or a diesel fuel into a northern California river that also impacts estuarine habitat, similar to the Smith River. The first column is the riverine habitat that could be affected. The second column are the potential resources at risk from a spill in that habitat. The third column describes the expected behavior of a diesel fuel or crude oil in that habitat. The fourth column describes the media samples to be collected, and the fifth column describes the impact assessment methods.

Table 1. Conceptual model for a northern California river spill that also impacts coastal resources (e.g., Smith River).

HABITAT	RESOURCES AT RISK	OIL BEHAVIOR	EXPOSURE ASSESSMENT	IMPACT ASSESSMENT
Tributary	<ul style="list-style-type: none"> • Riparian Vegetation • Fish spawning redds <ul style="list-style-type: none"> - Salmon - Trout • Outmigrating juvenile fish • Benthic macrofauna • Herpetofauna • Birds • Aquatic mammals • Hatchery water intakes • Recreational use <ul style="list-style-type: none"> - Fishing - Hiking 	<ul style="list-style-type: none"> • Moderate flow, riffles, pools, small falls so limited crude oil dispersion into the water column; diesel fuel is highly dispersible • Thick slicks pooling on surface; sheens extending far downstream • Oil coating of banks and mid-stream rocks, debris, vegetation • Oil penetration into sediments with falling water levels 	<ul style="list-style-type: none"> • Source oil sample • Sheen samples from pools near access points • Water column samples from larger pools below riffles near access points both upstream and downstream of the release site, and above hatchery water intakes • Tissue samples for bioavailability (e.g., crayfish and corbicula) • Sampling of oiled sediments 	<ul style="list-style-type: none"> • Oil/impact documentation with photography/field notes • Documentation of dead fish by counts by species and size • Documentation of presence (counts by species) and oiling/mortality of other fauna (e.g., birds/aquatic mammals/herpetofauna) during spill site visits • Benthic macro-invertebrate surveys in wadable reaches above/below release site • Comparison of water, tissue, and sediment chemistry to relevant toxicological benchmarks • Bioassays with spilled oil • Recreational user counts
Main Stem	<ul style="list-style-type: none"> • Riparian vegetation • Juvenile/adult fish <ul style="list-style-type: none"> - Salmon - Trout - Steelhead - Lamprey - Sturgeon • Benthic macrofauna • Birds • Aquatic mammals • Recreational use <ul style="list-style-type: none"> - Float fishing - Shore fishing - Kayak/canoe - Hiking - Swimming - Camping 	<ul style="list-style-type: none"> • Rapid flow with large rapids so high risk of oil dispersion into the water column, with potential for fish kills and impacts to benthic macrofauna and periphyton, especially for diesel fuel spills • Sheens re-surface in pools • Crude oil coating of banks, mid-stream rocks, debris, vegetation • Oil penetration into sediments with falling water levels 	<ul style="list-style-type: none"> • Source oil sample • Sheen samples from pools near access points • Water column samples at the head of pools below rapids near access points both upstream and downstream of the release site • Tissue samples for bioavailability (e.g., crayfish and corbicula), upstream and downstream of the release site • Sampling of oiled sediments 	<ul style="list-style-type: none"> • Oil/impact documentation with photography and field notes • Documentation of dead fish by counts by species and size • Documentation of presence (counts by species) and oiling/mortality of birds/aquatic mammals/herpetofauna during spill site visits • Benthic macro-invertebrate surveys in wadable reaches above/below release site • Comparison of water, tissue, and sediment chemistry to relevant toxicological benchmarks • Bioassays with spilled oil • Recreational user counts

HABITAT	RESOURCES AT RISK	OIL BEHAVIOR	EXPOSURE ASSESSMENT	IMPACT ASSESSMENT
Estuary	<ul style="list-style-type: none"> • Submerged aquatic vegetation • Fish spawning • Juvenile fish rearing habitat, particularly salmonids • Smelt • Sturgeon • Crabs • Marine and coastal birds • Marine mammals • High recreational use <ul style="list-style-type: none"> - Boat fishing - Shore fishing - Kayak/canoe • - Hiking 	<ul style="list-style-type: none"> • Thinner slicks on water surface from wider channels and resurfacing of dispersed oil • Risk of deposition of oil:particle agglomerations on the sea floor in the head of the estuary • Crude oil coats intertidal shoreline sediments and vegetation; diesel fuel forms a greasy band • Oil penetration into porous sediments intertidal 	<ul style="list-style-type: none"> • Source oil sample • Sheen/slick samples from the water surface • Water column samples where the river enters the estuary • Tissue samples for bioavailability (mussels/clams), from sites with visible oil and unoiled reference sites • Sampling of oiled sediments • 	<ul style="list-style-type: none"> • Oil/impact documentation with photography and field notes • Documentation of dead fish by counts by species and size • Documentation of presence (counts and species) and oiling/mortality of birds/crabs/marine mammals during spill site visits • Comparison of water, tissue, and sediment chemistry to relevant toxicological benchmarks • Bioassays with spilled oil • Counts of recreational users

Notes: Assume that wildlife rescue and rehab teams will be mobilized for birds and mammals as part of the response.

The timeline for ephemeral data collection is shown in Table 2. In the first 1-3 days of a spill, the first agency responders are likely local resource managers who are tasked with reconnaissance and documentation of the extent of oiling and immediate impacts to animals in oiled areas.

Table 2. Timeline for ephemeral data collection for Northern California rivers and estuaries.

VERY EARLY (DAYS 1-3)	EARLY (DAYS 3-6)	NEXT (DAYS 7-14)
<ul style="list-style-type: none"> • Source oil (best available) • Slicks/sheens from pools below the release site • Water samples from upstream and downstream of the release site • Reconnaissance <ul style="list-style-type: none"> ▪ Documentation of dead fish/crabs and other aquatic invertebrates ▪ Documentation of presence and oiling of birds/aquatic mammals/ herpetofauna during spill site visits ▪ Documentation of site location (e.g., oiled vegetation etc.) 	<ul style="list-style-type: none"> • Tissue samples from upstream and downstream of the release site • Sheen and water samples as the oil spreads downstream • Reconnaissance <ul style="list-style-type: none"> ▪ Continued documentation of dead fish/crabs and other aquatic invertebrates downstream ▪ Documentation of oiling in river on vegetation/debris/rocks ▪ Documentation of presence and oiling/mortality of birds/aquatic mammals/herpetofauna during spill site visits ▪ Documentation of site location (e.g., oiled vegetation etc.) 	<ul style="list-style-type: none"> • Continued documentation of site location (e.g., oiled vegetation, staining of rocks on shoreline) • Repeat water column sampling at established sites for time-series data • Continued documentation of dead fish/crabs and other aquatic invertebrates downstream • Continued documentation of counts and oiling/mortality of birds/aquatic mammals/herpetofauna during spill site visits • Sampling of oiled sediments • Kick-net surveys in wadable streams • Other quantitative biological surveys (e.g., benthic macroinvertebrates) • Tissue samples from downstream locations

Note: It is likely that wildlife rescue and rehabilitation teams will be mobilized to collect birds and mammals as part of the response.

3.0 Ephemeral Data Collection Procedures

3.1 Overview

This section describes procedures for collecting ephemeral data on source oil, weathered oil (i.e., tarballs), sheen, water, tissue, and sediment, as well as methods for assessing impacts to human/recreational use and benthic macroinvertebrates (BMI). The protocols described below are to be followed unless the NRDAR Agency Lead(s) decide otherwise and provide an alternative. This section provides a general overview of sample collection procedures. Appendices 1 to 9 provide more detailed standard operating procedures and guidelines for the different types of samples and data. **Appendix 10** includes example Coordination Agreements (e.g., for cooperative assessments with Responsible Party) and Initiate Funding Requests to the National Pollution Fund Center (NPFC). **Appendix 11** provides links

to examples of resource-specific injury assessment work plans. **Appendix 12** provides links to oil spill job aids and data sets on baseline resource condition.

3.2 Source Oil/Oil

It is critical that all sources of released oil be identified and sampled at the point of release. This is typically, and most appropriately, done as part of the response activities directed by the Unified Command or by state and federal incident investigators, not NRDAR personnel. The NRDAR Representative will coordinate with the Unified Command and incident investigators to confirm that sampling of the released material (e.g., oil) from its source will be conducted and can be split with or made available for the NRDAR efforts. The NRDAR Representative also will brief the incident investigators and Unified Command regarding NRDAR sampling activities and any special NRDAR concerns. Source sample collection at the point of release is overseen by the State incident investigator (i.e., warden) and collections are normally done by CDFW-OSPR Oil Spill Prevention Specialists. Source sampling by NRDAR personnel is usually not recommended because of the hazards and expertise needed to sample fuel tanks, pipelines, or vessels following a release.

Sampling of released oil in the environment is recommended for documenting the spatial extent of impacts and confirming the source of the oil. Provided safety considerations are met (Section 1.2), NRDAR field teams may collect these types of samples, such as tarballs. A detailed standard operating procedure for source oil sampling is included in **Appendix 1**.

3.3 Sheen

In cases where the released material is visible as a rainbow sheen either floating on water or coating structures, samples may be collected for petroleum fingerprinting purposes and to define the extent of the release. Sheen samples are collected by passing fiberglass sheets through the sheen floating on water or coating substrate, at approximately a 90-degree angle. The fiberglass sheets are placed into pre-cleaned glass jar and capped. See **Appendix 2** for a detailed standard operating procedure for collecting sheen samples.

3.4 Water

Water samples can be collected for analysis of benzene, toluene, ethylbenzene, and xylenes (BTEX), total petroleum hydrocarbons (TPH), and polycyclic aromatic hydrocarbons (PAH) to document exposure of water-column organisms to oil compounds, to determine the source of the oil in the water column via chemical fingerprinting the oil, and to support oil transport modeling. Oil in water is one of the most ephemeral types of data, so rapid collection after a release is important. Samples for TPH and PAH are collected in 1 L glass jars, so they can take up a lot of space and can be heavy, which can be challenging for areas without road access. Samples are collected at accessible locations from upstream, unoiled sites, and downstream oiled sites at various distances below the release site.

The goal is to collect samples of the dissolved oil fractions; therefore, water samples should be collected in streams and rivers from pools rather than areas of rapids and falls, to avoid inclusion of oil droplets. Sampling areas need also to consider important habitats for natural resources with trustee responsibility. When taking water samples, it is important to prevent contamination with surface oil. Visible oil should be moved aside with a water hose, paddle, or sorbent pad. The sampling container should always be opened only at the sampling depth. Usually, a water sample is collected at a depth of 0.5 ft below the surface. See **Appendix 3** for a detailed standard operating procedure for collecting water samples.

3.5 Tissue

Tissue samples from mollusks and other invertebrates are collected to document the concentrations and composition of oil compounds that are bioavailable and an exposure pathway to higher trophic organisms. Most invertebrates uptake both dissolved and dispersed oil quickly, whereas depuration takes weeks. Therefore, time-series tissue samples can record past exposure levels and then document when exposure starts to decrease. Invertebrates in Northern California river and estuarine habitats that may be collected for tissue analysis include bivalves, crayfish, snails, and caddisflies. Tissue samples are relatively easy to sample; whole animals are collected, placed in aluminum foil, and kept cold until transport to the laboratory. Tissue samples are collected at accessible locations from upstream, unoiled sites and oiled sites at various distances below the release site. See **Appendix 4** for a detailed standard operating procedure for collecting tissue samples.

3.6 Sediment

Samples of oiled sediment are collected for chemical analysis for fingerprinting to the source oil, to document exposure of sediment-dwelling organisms to the oil, and to track oil weathering over time. There are sediment quality criteria that can be compared with the concentrations of PAHs to determine initial toxicity and changes over time. Sampling locations for oiled sediment should be representative of potential exposure, such as from stream banks or the intertidal zone where the oil has stranded or accumulated, and/or from below the water surface. Samples are collected at accessible locations from upstream, unoiled sites, and oiled sites various distances below the release site. Sampling areas need also to consider important habitats for natural resources with trustee responsibility.

Composite samples (of at least three subsamples) are preferred for characterization of a sampling site. Sub-samples of sediments on the bank should be collected using a pre-cleaned stainless steel scoop or wooden spoon at random locations within a 5-meter radius and within the same water-level or tidal zone or sampling interval. The subsamples are homogenized in a clean aluminum tray before placement in a pre-cleaned 250 mL glass jar.

Ideal sediment collection sites are wadable depositional zones. These zones are often where there is low energy, shallow waters, in pools, in the inside of stream bends, or downstream of point bars. Specific areas of trustee concern (e.g., sediments near mussel beds, fish spawning areas) would need to be considered and prioritized. Starting from a downstream point, approach the selected sampling site by moving upstream. Using a pre-cleaned, hand-held stainless-steel scoop or wooden spoon, scoop the sediment from the bottom of the waterbody in the upstream direction. Collect sediment from 5 to 10 depositional zones that represent the left bank, right bank, center channel and different depths (if practical). Each sample should be described (grain size, depth, visual oiling conditions) and photographed. Sampling using cores and sediment grabs in deeper water are usually conducted at a later stage of the assessment. See **Appendix 5** for a detailed standard operating procedure for collecting sediment samples.

3.7 Human Use/Recreational Use

Human and recreational uses of natural resources may be impacted as a result of an oil spill and the ensuing incident response (e.g., closure of recreational areas). Many of these impacts may be relatively short-term and difficult or impossible to document later. The primary purpose of this initial data collection is to determine the type of uses and injuries that should be investigated in longer-term studies. Data that local land managers should consider collecting to assess potential human use impacts in the immediate aftermath of a spill include:

- Photo documentation of impacted uses (e.g., closures, advisory signs, etc.);
- Curation of relevant media such as news articles and social media posts;

- Identification of areas of heavy recreational use for further study; and
- As appropriate based on available time and staff experience, observations of recreational users (counts, activities) and interviews with recreational users in the impacted area or with site managers familiar with the normal regional recreational activities.

Early data collection, even if qualitative, is important and can guide the development of more quantitative studies if necessary. More detailed guidelines on the collection of ephemeral human use data can be found in **Appendix 6**.

3.8 Benthic Macroinvertebrates in Rivers

Collection of benthic macroinvertebrates is performed to document exposure to oil. This is a time-critical biological assessment method because affected fauna can be rapidly lost. At each stream reach, three samples are collected from evenly spaced across the stream width and composited into a single sample for that reach. Samples are collected at areas that have been or may be oiled, as well as unoiled reference locations. Samples should also be collected from locations known or suspected to be impacted by other natural or anthropogenic sources of contamination. Clean, unoiled 500- μm D-frame nets are used to collect samples. Collected samples are placed in a 500 mL wide-mouth plastic jar.

See **Appendix 7** for a detailed standard operating procedure for collecting benthic macroinvertebrate samples. The procedures described are based on the “Standard Operating Procedures for the Collection of Field Data for Bioassessments of California Wadable Streams: Benthic Macroinvertebrates, Algae, And Physical Habitat” (Ode et al. 2016) created by the Surface Water Ambient Monitoring Program (SWAMP). Following these procedures will allow comparison with historical data where available.

3.9 Field Documentation and Photography

Systematic field observations by the first responders in the days after a spill are key to documentation of potential injuries that may be lost before scientific surveys can be conducted. A dedicated field notebook should be used to record all observations, noting date and time, location, observations, and photographs. Do not erase or black out erroneous entries in the field notebook. Errors should be corrected by crossing out the entry with a single line and initialing and dating the strike-through.

The following is a partial list of items to always document:

- The general site conditions and location;
- How the oil spill happened – including oil source;
- Oil on the water surface, stranded on the shoreline, and in direct contact with or over sensitive areas such as shallow pools, in-stream debris, and dry stream beds
- Oiled wildlife;
- Oil recovery and cleanup operations (response);
- NRDAR staff working and sampling activities;
- Species presence and habitat use; and
- Site use by humans (kayaking, canoeing, fishing).

Field sketches showing the locations of these types of items are good to make, in addition to photographs. The field sketch is an important part of documentation because: 1) it provides a focused picture of the site conditions and affected resources and 2) it adds discipline to the field observation process because it forces the person doing the sketch to make detailed mental notes of all the relevant features at a site. Sketches, in combination with photographs, provide a powerful way to capture information.

Photographs and video are taken in the field to document the pre-oiling and oiling conditions and are key pieces of information that can be introduced as evidence. All the standard operating procedures

(SOPs) for ephemeral data collection include taking photographs. Each photograph should tell a specific part of the story. Before taking a photograph, you should consider what critical information you are trying to convey. Did the photograph capture the details you need? Are there key images (data) that you have missed? In the first case, you should take a better photograph (but without deleting any photographs). In the second case, you should look for photographs that will fill in gaps. Time series of photographs are helpful to document exposure or changes in oil degree and distribution over time. Time series should be taken by standing in the same spot and facing the same direction each time; it helps to have notes or prior photographs to be able to occupy the same location and perspective. See **Appendix 8** for a detailed standard operating procedure for field documentation using photography.

Imagery collected using uncrewed aerial systems (UAS) or drones can also be used to document extent of oiling, response activities, and user groups, among others. Any deployment of UAS must comply with requirements from the Federal Aviation Authority as well as airspace restrictions and other restrictions from organizations managing restricted areas. Examples include critical infrastructure; designated wilderness areas; Wildlife Sanctuaries; National Parks; and areas where wildlife disturbance may be a concern, such as marine mammal haul-outs and nesting bird colonies. The NOAA (2021) guide provides specific guidance for use of small UAS to collect data in support of an oil spill response.

3.10 Chain of Custody

For the purposes of litigation, agencies must be able to prove the legal integrity of all samples and data introduced as evidence. This means that it is necessary to have an accurate written record to track possession, handling, and location of samples and data from collection through reporting. Chain of custody (COC) procedures are followed to authenticate a sample from the time it is taken until the results are introduced as evidence and facilitate the verification process. Failure to follow chain of custody procedures does not necessarily render data unusable; however, any deviations from the chain of custody guidelines should be noted. Assuring that proper chain of custody guidelines are followed is vital to assuring the integrity of the samples, and the data generated by the analysis of those samples.

All samplers handling samples collected for NRDAR MUST follow chain of custody procedures when collecting, handling, and securing samples. All team leads and supervisors are responsible for ensuring that the designated custodian(s) understand this procedure and strictly adhere to it for all sampling events. The sample collector is responsible for care and custody of the samples until they are turned over to an assigned custodian or properly dispatched to the receiving laboratory. All custodians must ensure that each sample remains in their custody (as defined below) so that no one can tamper with it during the entire duration of their responsibility.

The chain of custody form is a document detailing who is legally responsible for samples at any point in time from collection until the sample is received by the laboratory. A sample is in your custody when:

- It is in your actual physical control and presence;
- It is in your view after being in your possession;
- It is not in your physical presence, but is secure in a place of storage to which only you have access; or
- It is not in your view or physical presence but is secured in a place of storage or secure area to which only you and identified others have access.

Before shipping samples, make sure that each chain of custody form is filled out completely and properly. Check that the sample identifications on sample bottles match the sample identifications on the chain of custody. Verify that the date, time, type, matrix and container types, and analyses requested are clearly indicated. Not all locations will ship samples with wet ice; however, when shipping with wet ice be sure that the cooler is properly sealed, contained from leaking, and that bottles are protected from breakage. Hazardous material shipping regulations must be followed if samples contain

large volumes of oil. Consult with the laboratory about the volume of oil that will trigger these regulations. The CDFW-OSPR chain of custody form and procedures are provided in **Appendix 9**.

3.11 Chemical Analysis Guidelines

Samples collected with the SOPs in Appendices 1 through 5 will be sent to analytical laboratories under chain of custody. This section outlines the types of analysis generally requested following oil spills.

The NRDAR Agency Lead(s) and the EDCC shall coordinate which chemical analyses are suitable for the collected samples. Table 3 presents the anticipated container and storage types by sample matrix. Table 4 presents the standard analyses and methods by sample matrix. Samples will be sent to a laboratory based on the decision of the NRDAR Agency Lead(s). The address and shipping instructions will be on the chain of custody form.

Table 3. Key sampling requirements for each sample type.

SAMPLE TYPE	COLLECTION CONTAINER(S)	STORAGE
Source Oil/Oil	<ul style="list-style-type: none"> • 1 L wide mouth glass jars, amber glass preferred • 250 mL wide mouth glass jars, amber glass for weathered oil such as tarballs 	Keep at 4°C. Do not freeze.
Sheen	<ul style="list-style-type: none"> • Four 12" fiberglass sheets or Teflon nets placed in 250 mL glass jar 	Keep at 4°C. Do not freeze.
Water	<ul style="list-style-type: none"> • 1 L glass jars, amber glass preferred • 40 mL septum-capped vials, HCl-preserved preferred, amber glass preferred, for volatiles analysis 	Keep at 4°C. Do not freeze.
Tissue	<ul style="list-style-type: none"> • Double layer of foil inside a double layer zipper-top bag 	Keep at 4°C. Do not freeze.
Sediment	<ul style="list-style-type: none"> • 250 mL certified organic-clean jars with Teflon-lined lids – for TPH/PAH/biomarkers; can use 1 L bottles if smaller-sized jars are not available • Zipper-top or Whirl-Pak bags – for grain size and TOC samples 	TOC - Keep at 4°C. Do not freeze. Grain Size – No refrigeration required.

Table 4. Analytes and analysis methods by matrix.

ANALYTES (SUGGESTED ANALYSIS METHOD)	SOURCE OIL/ SHEEN/TARBALL	WATER	SEDIMENT	TISSUE
PAHs and alkylated PAHs (EPA Method 8270-modified-GC/MS/SIM)	X	X	X	X
Total Petroleum or Extractable Hydrocarbons (TPH/TEH) (EPA Method 8015, GC/FID, extended range)	X	X	X	n/a
BTEX (EPA Method 8260 modified GC/MS/SIM)	TBD	TBD	TBD	n/a
% Moisture	n/a	n/a	X	X
% Lipids	n/a	n/a	n/a	X
Grain Size, Total Organic Carbon	n/a	n/a	X	n/a

The request for BTEX (benzene, toluene, ethylbenzene and xylene) analysis will depend upon the volatility of the source oil; BTEX is more prevalent in lighter weight fuels such as gasoline than in crude oils. Additional methods that may be requested for oil samples include SARA (saturates, aromatics, resins and asphaltenes), PIANO (paraffins, iso-paraffins, aromatics, naphthenes, and olefins; may include

BTEX), density, boiling curve (simulated distillation), metals, and sulfur content. Fingerprinting analyses vary between laboratories and may include standard terpane and sterane biomarkers as well as mono- and tri-aromatic steroids. Contact the EDCC or the laboratory that will be analyzing these samples to determine which analyses to request.

4.0 Natural Resource Trustee Representatives Contact Information (current as of July 2023)

4.1 CA Department of Fish and Wildlife, Office of Spill Prevention and Response

4.2 U.S. Fish and Wildlife Service

4.3 U.S. Department of the Interior

Office of Environmental Policy & Compliance

Bureau of Land Management, Northern California District Office

4.4 National Oceanic and Atmospheric Administration, Office of Response and Restoration

4.5 U.S. Department of Agriculture

National Response Team / Regional Spill Coordinator - Pacific Southwest Region

5.0 Technical Support And Chemical Analysis Contact Information

5.1 U.S. Environmental Protection Agency

Pacific Southwest (Region 9) Duty Officer for oil or chemical spills

5.2 CA Department of Fish and Wildlife – Office of Spill Prevention and Response Bioassessment Team

5.3 Chemical Analysis/Technical Support

California Department of Fish and Wildlife Petroleum Chemistry Laboratory

6.0 Other State and Local Agency Contacts

6.1 CA Department of Fish and Wildlife – Regional Office

Northern Region

North Central Region

Marine Region (All CA coastline)

6.2 CA Department of Parks and Recreation

North Coast Redwoods District

Northern Buttes District Sonoma-

Mendocino Coast District

Sierra District

Office of Historic Preservation

6.3 CA Regional Water Quality Control Board

North Coast (Region 1)

Central Valley (Region 5; Redding Office)

6.4 Office of Environmental Health Hazard Assessment

Pesticide and Environmental Toxicology Branch

6.5 California Coastal Commission

6.6 California State Lands Commission

6.7 Public Utilities

Pacific Gas & Electric

California Geologic Energy Management Division (CalGEM) - Northern District

6.8 Railroads

Union Pacific Railroad (UPRR)

Burlington Northern Santa Fe (BNSF)

6.9 Tribal Contacts

California Native American Heritage Commission

7.0 References Cited

Ode, P.R., A.E Fetscher, and L.B. Busse. 2016. Standard Operating Procedures for the Collection of Field Data for Bioassessments of California Wadable Streams: Benthic Macroinvertebrates, Algae, and Physical Habitat. SWAMP Bioassessment Procedures.
https://www.waterboards.ca.gov/water_issues/programs/swamp/bioassessment/sops.html

Soucek, D.J., A.M. Farag, J.M. Besser, and J.A. Steevens. 2023. Guide for benthic invertebrate studies in support of Natural Resource Damage Assessment and Restoration: U.S. Geological Survey Open-File Report 2022-1110, 11 p. <https://doi.org/10.3133/ofr20221110>.

West Coast Joint Assessment Team (WCJAT). 2017. Recommendations for Conducting Cooperative Natural Resource Damage Assessment.

<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=200743&inline>

APPENDIX 1: Detailed Standard Operating Procedures for: Source Oil and Oil

Standard Operating Procedure for Collecting Ephemeral Data: SOURCE OIL and OIL

June 2023

Sampling Objectives

Characterize Oil

- Obtain sample(s) of the original oil source(s) involved in the incident for characterization, fingerprinting, and predicting/quantifying oil fate and effects, (e.g., sediment toxicity testing).
- Obtain samples of oil released to the environment for fingerprinting to the source oil.

Quality Assurance/Quality Control

- Ensure the integrity of the oil sample(s) throughout sampling, transport, and storage.
- Ensure the reliability of chemical characterizations.

Before Field Sampling

- Ensure that all personnel have required safety training and protective equipment for field work.
- Source sample collection at the point of release is overseen by the State incident investigator (i.e., warden) and collections are normally done by CDFW-OSPR Oil Spill Prevention Specialists. Source sampling by NRDAR personnel is usually not recommended because of the hazards and expertise needed to sample fuel tanks, pipelines, or vessels following a release. Coordinate with the emergency response before carrying out source oil sampling.
- Sampling of released oil in the environment is **recommended for documenting the spatial extent of impacts and confirming the source of the oil in the environment**. Provided safety considerations are met, NRDAR field teams may collect these types of oil samples.

Sampling Areas and Timing

- Follow a sampling plan/work plan if one is available.
- The source oil sample should be collected from the freshest source possible (i.e., directly from the tank truck, wellhead, or pipeline); secondary options are from the water or ground as close as practical to the spill release site.

Field Sampling Methods

Sampling Equipment/Containers

Note: Analytical laboratories may provide required sampling and sample storage and transport materials.

- Coolers – for sample storage and transport
- Blue ice packs or wet ice – for storage temperature regulation
- Thermometer or temperature logger (1 per cooler)
- Disposable nitrile gloves
- Sampling jars – certified organic-clean glass jars (solvent rinsed) with Teflon-lined lids and labels:
 - 1 L wide mouth glass jars, amber glass preferred
 - 250 mL wide-mouth glass jars, with Teflon-lined lid, amber glass for weathered oil such as tarballs
 - If BTEX (volatile) samples are to be collected, 40 mL septum-capped vials, HCl-preserved preferred, amber glass preferred

- Samplers – Wheaton grab sampler, Volskom sampler (if available)
- Stainless steel spatulas or spoons for collection of viscous oil samples
- Sampler material – Teflon (preferred), glass, or PVC (less ideal)
- Funnels (for transferring; solvent rinsed) – metal/stainless steel (preferred), glass or plastic (less ideal)
- Field sample forms (template in Appendix 9) and field notebook
- Chain of Custody forms
- GPS, camera (with spare batteries), photo scales, evidence tape
- Packaging materials (bubble wrap, sorbent pads, and tape) for glass jars – may be provided by the analytic laboratory
- Suitable disposal bags for oiled PPE and disposable sampling materials

Sample Collection Methods

- Use field data forms included in the work plan if one is available. Otherwise, use form in Appendix 9. Coordinate data form development/modification with the NRDAR data management group.
- Make sure that all GPS units are using the same coordinate system, datum, reporting units, and correct time. Follow the recommended GPS datum of the study plan if one is available. Alternatively, set the default to WGS84.
- Record GPS coordinates for each sample site.
- Photograph the sampling site prior to sample collection to document the site conditions, as well as the sample collected. Make sure each photograph or series can be later associated with the corresponding sampling locations (e.g., through use of GPS Photo link software or by keeping a detailed photo log with waypoints and/or lat/long). Do not delete or alter any photographs. The numbering sequence of photographs uploaded from your camera must not have any gaps (see Field Photography guidelines).
- Safety is the highest priority. Be aware of physical and chemical hazards at the site. Get a safety briefing from the Safety Officer before entering any restricted area. Use recommended safety equipment and procedures.
- There may be different products loaded in different compartments or tanks. Get as much data as practical on the products from the RP. Slight differences in product properties, such as specific gravity, may indicate different products. Each unique product should be sampled.
- Collect a minimum of 1 L per source.
- Collect sample with gloved hands.
- When collecting source oil:
 - Source samples can be collected directly into the sample container, minimizing risks of contamination. Use a device that holds the container, such as a Wheaton grab sampler (holds a 1 L bottle strapped to a metal rod with the ability to unscrew the cap remotely) or a Volskom sampler (frame which holds a container lowered on a rope). For black oils, these methods are less ideal because of heavy oil coating of the device and container.
 - Sampler material, in order of preference, include Teflon (there are inexpensive, disposable models), glass, and PVC.
 - Leave 1 cm headspace in the neck of each jar. A larger headspace (e.g., 2.5 cm) is recommended to allow for sample expansion if water is present in the sample and freezing may occur.
 - Care should be taken not to contaminate the outsides of the lid or the sample jars. If contamination does occur, the containers should be cleaned with soap and water, or less preferably with sorbent material. Do not store containers that have external oil contamination with other samples until they have been cleaned.

- If the collection of samples directly from the source is not possible, 1 L samples of oil should be taken from the water or ground as close to the source as practical.
 - When collecting source oil from water surface, concentrate the source oil in the sample container by skimming the oil from the water, sampling from the thickest part of the freshest oil slick.
 - When collecting source oil from the ground, carefully scrape the thickest part of the stranded source oil from the surface into the sample container minimizing the transfer of sediment or other foreign material.
 - Be aware of sources of contamination or mixing of products on-scene, such as transferring of product between tanks, dilution with fire-fighting water, or application of a foam blanket. If contamination is suspected, it may be appropriate to take samples of the potential source of contaminant.

Sample Labeling and Record Keeping

- Verify that all oil samples are properly labeled and that field sample forms are properly filled out.
- Follow chain of custody procedures for securing samples and complete chain of custody forms (See Chain of Custody guidelines in Appendix 9).
- Make notation on the Chain of Custody form about any problems or observations during sampling, such as presence of water or sediment in the sample.
- Maintain strict chain of custody during sample storage and transportation.
- Record the sample number on both the sample jar label and lid. Fill out the data sheet.
- Keep a detailed photo log so that each photograph can be labeled. All sample numbers must be unique. The sample number should consist of a 2-letter Field Team name (e.g., ED for ephemeral data), 1-number field team number, 6-number date (MMDDYY), 2-letter sample type, and a 2-number sequential sample number. See the Sample Collection Summary Form for codes in Appendix 9. All sample numbers must be unique. Note any deviations from the SOPs in the field notebook.

Sample Preservation, Holding Times, and Shipping

- Tape lids on sample bottles so that they do not accidentally come off.
- Protect the samples from direct sun exposure (e.g., UV radiation).
- Immediately place all source oil samples in a refrigerator or cooler and keep at 4°C. DO NOT FREEZE. Use frozen gel packs or wet ice to maintain the temperature. A programmable temperature logger or thermometer should be placed in each cooler to maintain a record of storage temperatures.
- Keep source oil samples separate from environmental samples to reduce risk of cross contamination.
- Keep the samples under chain of custody until they are handed off for transport to the laboratory.
- Do not discard ANY oil samples.


APPENDIX 2: Detailed Standard Operating Procedures for: Sheen

Standard Operating Procedure for Collecting Ephemeral Data: SHEEN

June 2023

Definition

By definition, sheen is a very thin layer of oil on the water surface that is described as silver (S), rainbow (R), or metallic (M) as shown in the image below.



Sampling Objectives

Characterize Oil

- Determine the source of sheens via chemical fingerprinting analysis
- Document the presence of oil, and characterize oil weathering and fate

Study Exposure

- Document exposure of water-surface organisms to oil sheen compounds
- Support exposure modeling

Quality Assurance/Quality Control

- Ensure the integrity of the sample(s) throughout sampling, transport, and storage
- Ensure the reliability of chemical characterizations

Before Field Sampling

- Ensure that all personnel have required safety training and protective equipment for field work.

Study Design

- It is important to have a defined sampling strategy prior to conducting fieldwork; however, this plan may have to be modified based on actual conditions in the field because sheens are very

dynamic. Sampling sheens may be opportunistic if one is encountered during other sampling efforts.

- The following terminology is used to define general to specific sampling geographies:
 - Area = general area of uniform characteristics, such as degree of oil exposure, physical setting, habitat types present, etc.
 - Location = a specific location that is representative of the area and contains the type of habitat to be sampled, such as an eelgrass bed or lagoon
 - Site = a specific point at which samples are collected or observations are made
- Plan the number of sites and the number of samples to be collected at each site, accounting for level of effort, potential logistical limitations, weather conditions, and other unanticipated issues that may compromise sample integrity.

Equipment

- Review the list of sampling equipment/containers, adjust as needed, and ensure that all essential field materials are ready to be taken to the field.
- It may be necessary to coordinate with the laboratory that will receive the samples to ensure that acceptable materials and conditions are used for sampling, sample storage, and shipping.
- Make sure that all essential equipment is in working order and that spare equipment and materials are available.

Sampling Areas and Timing

- Follow a sampling plan/work plan if one is available.
- Sheen samples should be collected from specific locations to answer specific questions, such as what is the source of a reported sheen, what is the spatial extent of oiling, or how is the oil on the water surface weathering over time and distance?
- The number of samples collected need to be considered accordingly, making sure that there is enough space in the coolers to accommodate all samples without sacrificing their integrity.

Area Selection

- Sampling should focus on collecting samples of sheens from the spill and other natural or non-spill sources if they are observed.
- Opportunistic sheen samples should be collected if sheens are observed during other field data collection activities.
- The number of locations and number of samples per location should be defined in the study design. A minimum guideline for collecting sheen samples is one per sampling location for fingerprinting.

Field Sampling Methods

Sampling Equipment/Containers

- Coolers – for sample storage and transport
- Blue ice packs or wet ice – for storage temperature regulation
- Thermometer or temperature logger (1 per cooler)
- Disposable nitrile gloves
- 250 mL wide-mouth sampling jars – certified organic clean glass jars (solvent rinsed) with Teflon-lined lids and labels

- Teflon (PTFE-fluorocarbon polymer) nets/sheets (preferred) – 4-inch diameter with deployment gear (wand, pole, line). Alternate: Four 12-inch fiberglass sheets or sorbent pads
- Tweezers, hemostats, or pliers – one for each sample, pre-cleaned, wrapped in foil
- Field Sample Forms (template in Appendix 9) and field notebook
- Chain of Custody forms (see Chain of Custody guideline)
- GPS, camera (with spare batteries), and photo scales
- Packaging materials (bubble wrap, sorbent pads, and tape) for glass jars
- Suitable disposal bags for oiled PPE and disposable sampling materials

Sample Collection Methods

- Use field data forms included in the work plan if one is available. Otherwise, use forms in Appendix 9. Coordinate data form development/modification with the data management group.
- Because GPS units will be used to record locations and times, make sure that all units are using the same coordinate system, datum, reporting units, and correct time. Follow the recommended GPS datum of the study plan, if one is available. Alternatively, set the default to WGS84.
- Record GPS coordinates for each sample site.
- For each sampling site, record:
 - Date, time, weather conditions (e.g., wind direction and speed), and tide level
 - Presence of biological resources or other relevant information
- Photograph the sampling site prior to sample collection to document the site conditions, as well as the sample collected. Make sure each photograph or series can be later associated with the corresponding sampling locations (e.g., through use of GPS Photo link software or by keeping a detailed photo log with waypoints and/or lat/long). Do not delete or alter any photos. The numbering sequence of photos uploaded from your camera must not have any gaps (see Field Photography guidelines, Appendix 8).
- A description of the characteristics and spatial extent of the sheen that the samples are collected from should be recorded in the field data sheet or notebook.
- Sheen samples can be collected from boats, from the shoreline, or by wading in shallow water.
- Collect sample with gloved hands.
- Teflon (PTFE-fluorocarbon polymer) nets (50 to 70 micron-mesh screen, preferable) or fiberglass sheets, can be hand-held, attached to a sampling wand/pole, or attached to the line of a fishing pole. Slowly drag the net or sheet through the sheen at least five times or until the net or pads are visibly oiled, and transfer the material into a 250 mL glass container touching the material as little as possible. Use clean tweezers, hemostats, or pliers to handle the Teflon net or pad. Because light PAH fractions are extremely volatile, NEVER split sheen samples after sample collection.

Quality Assurance/Quality Control

- Obtaining an adequate number of quality control samples is essential. At a minimum, a trip blank (accounts for contamination introduced during shipping and handling) and field blank (account for contamination introduced during sampling) should be maintained for each sampling effort and generally be collected at a rate of 5% and 10%, respectively, of all samples.
- A trip blank is an unopened sampling jar and should be transported with the samples and remain sealed in the cooler during sampling activities.
- A field blank should be collected at approximately every third sampling site, or at least at an “un-oiled” and “oiled” site, by leaving the field blank sample jar open for the duration of the

sampling period at that site. Record the site where field blanks were taken on the field sample form.

- Ideally, trip and field blanks are a sampling jar with Teflon sampling material in them.
- If possible, store samples from field/trip blanks in one set of coolers, with oiled samples in a separate set of coolers. If possible, do not include other types of samples in the coolers for the sheen samples; otherwise, take precautions to prevent cross contamination.
- Rinsate blanks should be collected if there is a risk of cross contamination from reuse of sampling equipment. After cleaning the equipment in accordance with the procedures described in this method, rinse the clean equipment with solvent or cleaning solution and collect the rinsate in a sample jar. Note on the field sample form where and how rinsate blanks were collected.

Good Sampling Practices and Decontamination

- Good field practices and the development of a consistent sampling routine will ensure the integrity of the sheen samples and their validity in environmental assessments.
- Disposable nitrile gloves should be changed between samples to prevent cross contamination or if they become contaminated or damaged.
- Potential contamination while sampling from vessels (exhaust fumes, oily surfaces) is a very serious concern. Work up-wind of any exhausts, consider sampling sheens from non-motorized craft that is paddled upwind/current from the motorboat. Avoid sampling from the stern of motorboats.
- Take precautions to avoid cross-contamination of the site from oil on personal equipment. Sampling unoiled areas first, then lightly oiled areas and then heavily oiled areas can minimize cross-contamination. Personal equipment should be exchanged or cleaned between sites if it becomes oiled.

Sample Labeling and Record Keeping

- Verify that all samples are properly labeled and that field forms are properly filled out.
- Follow chain of custody procedures for securing samples and complete the Chain of Custody form, noting sample size, sampling device used and any other relevant information for the receiving laboratory in addition to the basic information about the sample indicated on the form. (See Chain of Custody guidelines).
- Make notation on the Chain of Custody form about any problems or observations during sampling.
- Maintain strict chain of custody during sample storage and transportation.
- The sample number should consist of a 2-letter Field Team name (e.g., ED for ephemeral data), 1-number field team number, 6-number date (MMDDYY), 2-letter sample type, and a 2-number sequential sample number. See the Sample Collection Summary Form for codes in Appendix 9. All sample numbers must be unique.
- Keep a detailed photo log so that each photograph can be labeled.
- Note any deviations from the recommended guidelines in the field book.

Sample Preservation, Holding Times, and Shipping

- Follow chain of custody procedures for sample storage and shipping.
- Immediately following collection, place all sheen samples in cooler and keep at approximately 4°C. Use frozen gel packs or wet ice to maintain the temperature if ambient temperatures are above freezing. In below freezing temperatures, collapsible water jugs filled with warm water can be used to maintain the temperature if heated storage space is not available. A

programmable temperature logger or thermometer should be placed in each cooler to maintain a record of storage temperatures.

- Protect the samples from direct sun exposure (e.g., UV radiation).
- Tape lids on sample jars in accordance with chain of custody guidelines so they do not accidentally come off.
- Sheen samples should be stored and transported in a separate set of coolers.
- Use packing material, such as bubble wrap or sorbent pads, around containers to prevent breakage during handling and shipping.
- Ship samples directly to the laboratory as soon as possible with complete chain of custody forms. Holding time for sheen samples is 7 days. If necessary, samples can be stored under specified conditions and with complete chain of custody until they can be shipped. Ensure that samples are packaged to protect them from breakage, shipping containers are sealed, and use ice packs or wet ice to maintain storage temperatures during shipment to the lab.
- Ship highly oil-contaminated samples separate from non-contaminated or low-contaminated samples to reduce risk of cross contamination.
- NEVER discard ANY samples even if these have exceeded their recommended holding times or storage temperatures.
- Refer to the Chemical Analysis Guidelines in Table 4 for analytical methods, sample volumes, minimum detection limits, and recommended holding time.

APPENDIX 3: Detailed Standard Operating Procedures for: Water

Standard Operating Procedure for Collecting Ephemeral Data: WATER

June 2023

Sampling Objectives

Characterize Oil in Water Media

- Determine the concentration and composition of oil compounds in the water column
- Determine the source of contamination via chemical fingerprinting analysis and characterize oil weathering and fate

Study Exposure

- Document exposure of water-column organisms to oil compounds
- Support exposure and environmental transport modeling

Quality Assurance/Quality Control

- Ensure the integrity of the sample(s) throughout sampling, transport, and storage
- Ensure the reliability of chemical characterizations

Before Field Sampling

- Ensure that all personnel have required safety training and protective equipment for field work.

Study Design

- It is important to have a defined sampling strategy prior to conducting fieldwork.
- The following terminology is used to define general to specific sampling geographies
 - Area = general area of uniform characteristics, such as degree of oil exposure, physical setting, habitat types present, etc.
 - Location = a specific location that is representative of the area and contains the type of habitat to be sampled, such as a stream reach
 - Site = a specific point at which samples are collected or observations are made
- Plan the number of areas and samples to be collected at each area, taking into account level of effort, potential logistical limitations, weather conditions, and other issues that may compromise sample integrity. Generate a list of alternate sites when unforeseen circumstances prevent sampling at the primary site(s).
- For water samples, sampling “areas” can be defined as: 1) waterbodies with defined boundaries (such as stream reach); 2) distances down current from the release site (such as 0-5 km, 5-10 km); and 3) waterbodies that are expected to have similar oil exposure based on observations.
- Depending on the water depth, water samples can be collected at three depths: near surface (0-1 m), mid-depth, and 1 m above the bottom. Generally, near surface samples should be prioritized if the sampling effort is limited by logistics or other factors. In shallower water, samples should be collected just below the surface, but without including any surface oil.
- Contact the laboratory that will be receiving field samples for analysis and ensure that they have the capacity to receive and analyze samples from the study. Follow relevant guidelines from the laboratory and consult with them about necessary modifications.

Equipment

- Review the list of sampling equipment/containers, adjust as needed, and ensure that all essential field materials are ready to be taken to the field.
- It may be necessary to coordinate with the laboratory that will receive the samples to ensure that acceptable materials and conditions are used for sampling and sample storage and shipping.
- Make sure that all equipment is in working order and spare equipment and materials are available.

Sampling Areas and Timing

- Follow a sampling plan/work plan if one is available.
- If a sampling plan is not available, data collection should focus on collecting samples from a range of unoiled, likely to be oiled, and already oiled areas.
- The number of locations and number of sites per location need to be considered accordingly, making sure that there is enough space in the coolers to accommodate all samples without sacrificing their integrity.
- Water samples can be collected from boats or by wading in shallow water.

Area Selection

- Use your own observations, trajectory models, conceptual models, overflight observations, SCAT data, etc. to determine what locations have been oiled and which ones are likely to be oiled.
- Samples should also be collected from locations known or suspected to be affected by other natural or anthropogenic sources of contamination (e.g., oil seeps, coal, peat, mining, combustion engines), as these will be important to differentiate background sources and levels of contamination.
- It may be necessary to prioritize sampling locations. In this case, highest priority samples are to be collected from waters adjacent to locations that are sensitive habitats, biologically productive, or highly relevant for human use. Collecting pre-oiling water samples from sensitive/productive locations that are likely to be oiled by the spill in the near future is also a priority. Sampling at unoiled “control” areas and sampling other sources of contamination should be prioritized based on the ephemerality of the data and relative importance to developing a NRDAR case.
- Water samples should be collected pre-oiling, if possible, as soon as practical after oiling, and periodically thereafter. Sampling frequency should be defined in the study design.
- The number of locations and number of sites per location should be defined in the study design. A minimum guideline for collecting water samples is at least three samples per waterbody location. If logistical limitations are a concern, prioritize sample collection by selecting a minimum of one reference/pre-oiling location and two heavily oiled locations.
- Sample along exposure gradients, starting in the cleanest zone and then at regular intervals proportional to the exposure area.

Field Sampling Methods

Sampling Equipment/Containers

- Coolers – for sample storage and transport
- Blue ice packs/wet ice – for storage temperature regulation (if ambient temperature exceeds 4°C)
- Thermometer or temperature logger (1 per cooler)

- Disposable nitrile gloves
- Sampling jars – certified organic-clean glass jars (solvent rinsed) with Teflon-lined lids and labels:
 - 1 L glass jars, amber glass preferred
 - 40 mL septum-capped vials, HCl-preserved preferred, amber glass preferred
- Trip and field blanks – 1 L and 40 mL sampling jars filled with distilled water
- Sorbent pads
- Field Sample Forms (template in Appendix 9) and field notebook
- Chain of Custody forms (Appendix 9)
- Evidence tape (see Chain of Custody guidelines), GPS, camera (with spare batteries), photo scales
- Packaging materials (bubble wrap, sorbent pads, and tape) for glass jars – may be provided by the analytic laboratory
- Suitable disposal bags for oiled PPE and disposable sampling materials
- Subsurface water sampler (e.g., Niskin bottle or other) – if needed

Sample Collection Methods

- Use field data forms included in the work plan if one is available. Otherwise, use forms in Appendix 9. Coordinate data form development/modification with the data management group.
- Because GPS units will be used to record locations and times, make sure that all units are using the same coordinate system, datum, reporting units, and correct time. Follow the recommended GPS datum of the study plan if one is available. Alternatively, set the default to WGS84.
- Record the sampling site location using a GPS.
- For each sampling site, record:
 - Date, time, weather conditions (e.g., wind direction and speed), and tide level
 - Water depth (in meters) for water samples
 - Presence of biological resources or other relevant information
- Photograph the sampling site prior to sample collection to document the site conditions, as well as the sample collected. Make sure each photograph or series can be later associated with the corresponding sampling locations (e.g., through use of GPS Photo link software or by keeping a detailed photo log with waypoints and/or lat/long). Do not delete or alter any photographs. The numbering sequence of photographs uploaded from your camera must not have any gaps (see Field Photography guidelines, Appendix 8).
- To minimize risks of cross-contamination, collect water samples directly into the sample container by hand (wearing clean, disposable Nitrile gloves); a less ideal alternative is to use samplers that can hold 1 L glass bottles. This may be necessary for the collection of subsurface water samples where sampling bottles need to be opened/closed at the targeted water depth.
- Clear surface slicks and sheens prior to deploying the equipment by sweeping the area with a sorbent pad or placing a barrier up-current to divert surface oil around the sampling area, avoiding physical dispersion of the oil into the water column.
- Collect BTEX samples in HCl-preserved, 40 mL septum-capped vials. Fill vials completely and cap at the sampling depth or, if using a water sampler, fill the vials to overflow and cap immediately. Vials should not have headspace or air bubbles. If BTEX sampling vials are not available, water samples for THC and PAH should still be collected.
- Collect water samples for THC and PAH in glass containers (organic clean). Leave headspace of about 2 cm for 1 L jars. If sampling directly into jars, fill completely and cap at the sampling

depth. Remove the cap only once the sampling jar is no longer in contact with the water and pour out the necessary volume to create headspace before recapping.

- Collect “near surface” water samples at a uniform depth (e.g., 30 cm, which would be up to your elbow if using your hands) below the water surface, taking care to avoid any surface slicks or sheens.
- If collecting samples by wading in shallow water, collect samples in waters that are at least 60 cm deep. Collect samples at a uniform depth (e.g., 30 cm, which would be up to your elbow if using your hands) below the surface. Avoid disturbing or suspending bottom material.
- When sampling by hand:
 - Stand facing the current, if any, and wait until any suspended material is flushed away
 - Plunge the bottle with the cap on, neck downward, under the water surface in front of you
 - Turn the bottle until the neck points slightly upwards with the mouth directed into the current
 - Uncap the sampling bottle and fill it. Do not touch the cap liner or the inside of the bottle
 - Cap the bottle under water immediately after filling
- When using a sampler:
 - Sampling equipment MUST be deployed and retrieved in the closed position, opening the sampler at the sampling depth
 - All field equipment that comes in contact with the sampling media MUST be thoroughly decontaminated after each sampling event to prevent inadvertent sample contamination
 - If possible, dedicate one set of sampling equipment per degree of oiling to minimize potential cross-contamination

Quality Assurance/Control

- Obtaining an adequate number of quality control samples is essential. At a minimum, a trip blank (accounts for contamination introduced during shipping and handling) and field blank (accounts for contamination introduced during sampling) should be maintained for each sampling effort and generally be collected at a rate of 5% and 10%, respectively, of all samples.
- Ideally, trip and field blanks are a sampling jar containing ultra-pure or distilled water. Blanks may be provided by the receiving laboratory.
- A trip blank is an unopened sampling jar and should be transported with the samples and remain sealed in the cooler during sampling activities.
- A field blank should be collected at approximately every third sampling site, or at least at an “un-oiled” and “oiled” site, by leaving the field blank sample jar open for the duration of the sampling period at that site. Record the site where field blanks were taken on the field sample form.
- If possible, store samples from field/trip blanks in one set of coolers, with oiled samples in a separate set of coolers. Otherwise, take precautions to prevent cross contamination.
- Do not split samples unless specified in the work plan.
- Rinsate blanks should be collected if there is a risk of cross contamination from reuse of sampling equipment. After cleaning the equipment in accordance with the procedures described in this method, rinse the clean equipment with solvent or cleaning solution and collect the rinsate in a sample jar. Note on the field sample form where and how rinsate blanks were collected.

Good Sampling Practices and Decontamination

- Good field practices and the development of a consistent sampling routine will ensure the integrity of the sheen samples and their validity in environmental assessments.
- Disposable nitrile gloves should be changed between samples to prevent cross contamination or if they become contaminated or damaged.
- Work upwind of any motor exhausts, consider sampling from non-motorized craft that is paddled upwind/current from the motorboat, and designate clean areas for sampling. Sampling on the windward side of the vessel is preferred. Avoid sampling from the stern of motorboats.
- Take precautions to avoid cross-contamination of the site from oil on personal equipment (e.g., boots, shovels, etc.). Sampling unoiled areas first, then lightly oiled areas and then heavily oiled areas can minimize cross-contamination. Personal equipment should be exchanged or cleaned between sites if it becomes oiled.

Sample Labeling and Record Keeping

- Verify that all samples are properly labeled and that field sample forms are properly filled out.
- Follow chain of custody procedures for securing samples and complete the Chain of Custody form, noting sample size, sampling device used and any other relevant information for the receiving laboratory in addition to the basic information about the sample indicated on the form. (See Chain of Custody guidelines).
- Make notation on the Chain of Custody form about any problems or observations during sampling.
- Maintain strict chain of custody during sample storage and transportation.
- Record the sample number on both the sample jar label and lid. All sample numbers must be unique. The sample number should consist of a 2-letter Field Team name (e.g., ED for ephemeral data), 1-number field team number, 6-number date (MMDDYY), 2-letter sample type, and a 2-number sequential sample number. See the Sample Collection Summary Form for codes in Appendix 9.
- Keep a detailed photo log so that each photograph can be labeled.
- Note any deviations from the recommended guidelines in the field book.

Sample Preservation, Holding Times, and Shipping

- Follow chain of custody procedures for sample storage and shipping.
- Immediately place all water samples in a cooler and keep at approximately 4°C. Use frozen gel packs or wet ice to maintain the temperature if ambient temperatures are above freezing. In below freezing temperatures, collapsible water jugs filled with warm water can be used to maintain the temperature if heated storage space is not available. A programmable temperature logger or thermometer should be placed in each cooler to maintain a record of storage temperatures.
- Protect the samples from direct sun exposure (e.g., UV radiation).
- Tape lids on sample jars in accordance with chain of custody guidelines so they do not accidentally come off.
- Store samples from unoiled areas in one set of coolers, with oiled samples in a separate set of coolers.
- Use packing material, such as bubble wrap or sorbent pads, around glass jars to prevent breakage during transport and shipping.
- Water samples can be held at 4°C in the dark for up to 7 days (includes recommended holding time in the field and receiving laboratory) without loss of sample integrity. Samples should not be frozen.

- Ship samples directly to the laboratory as soon as practical with complete chain of custody forms. If necessary, samples can be stored under specified conditions and with complete chain of custody until they can be shipped. Ensure that samples are packaged to protect them from breakage, shipping containers are sealed, and use ice packs or wet ice to maintain storage temperatures during shipment.
- Ship highly oil-contaminated samples separate from non-contaminated or low-contaminated samples to reduce risk of cross contamination.
- NEVER discard any samples even if these have exceeded their recommended holding times or storage temperatures.
- Refer to the Chemical Analysis Guidelines in Table 4 for analytical methods, sample volumes, minimum detection limits, and recommended holding time.

APPENDIX 4: Detailed Standard Operating Procedures for: Tissue

Standard Operating Procedure for Collecting Ephemeral Data: TISSUE

June 2023

Sampling Objectives

Characterize Oil in Tissues

- Determine the concentration and composition of oil compounds in biological tissues compared to background concentrations
- Determine the source of contamination via chemical fingerprinting analysis and characterize oil weathering and fate

Study Exposure and Injury

- Document the extent and duration of exposure to the spilled oil
- Document the bioavailability and exposure pathways of the spilled oil
- Quantify oil chemicals in tissues
- Document routes of exposure for higher trophic level organisms

Quality Assurance/Quality Control

- Ensure the integrity of the sample(s) throughout sampling, transport, and storage
- Ensure the reliability of biological characterizations

Before Field Sampling

- Ensure that all personnel have required safety training and protective equipment for field work.

Study Design

- The following terminology is used to define general to specific sampling geographies
 - Area = general area of uniform characteristics, such as degree of oil exposure, physical setting, habitats present, etc.
 - Location = a specific location that is representative of the area and contains the type of habitat to be sampled, such as a stream reach or tidal marsh
 - Site = a specific point at which samples are collected or observations are made
- Plan the number of locations and number of sites per location, accounting for level of effort, access, weather conditions, and other issues that may determine the actual sampling effort. Generate a list of prioritized alternate sites when unforeseen circumstances prevent sampling at the primary site(s).
- A stratified random sampling approach, which divides the sampling location into non-overlapping zones from which random samples are collected, is recommended if no other sampling strategy has been developed. This type of sampling improves the representative quality of samples by reducing sampling error (variability).
- Soucek et al. (2022) provide additional guidance on sample collection.
- Contact the laboratory that will be receiving field samples for analysis and ensure that they have the capacity to receive and analyze samples from the study. Follow relevant guidelines from the laboratory and consult with them about necessary modifications.
- The sampling strategy should have enough flexibility that it can be adjusted based on conditions in the field.

- Consult appropriate guidelines for the collection of other environmental media and biota concurrent with tissue sampling. Tarballs, sheens, or other oil residues can be collected opportunistically for chemical analysis and fingerprinting.

Equipment

- Review the list of sampling equipment/containers, adjust as needed and ensure that all essential field materials are ready to be taken to the field.
- It may be necessary to coordinate with the laboratory that will receive the samples to ensure that acceptable materials and conditions are used for sampling and sample storage and shipping.
- Do as much material preparation as possible prior to going into the field, including: labeling sample jars using permanent markers or laboratory labels, pre-cleaning sampling tools, etc.
- Make sure that all equipment is in working order and spare equipment and materials are available.

Sampling Areas and Timing

- Follow a sampling plan/work plan if one is available.
- If a sampling plan is not available, data collection should focus on collecting samples from a range of unoiled, likely to be oiled, and already oiled areas.
- The number of locations and number of sites per location need to be considered, making sure that there is enough space in the coolers to accommodate all samples without sacrificing their integrity.

Area Selection

- Sampling locations should be representative of areas that have been or may be oiled and unoiled reference locations, and include sensitive/productive sites.
- Use your own observations, trajectory models, conceptual models, overflight observations, SCAT data, etc. to determine what locations have been oiled and which ones are likely to be oiled.
- Samples should also be collected from locations known or suspected to be impacted by other natural or anthropogenic sources of contamination (e.g., oil seeps, coal, mining, combustion engines), as these will be important to differentiate background sources and levels of contamination.
- It may be necessary to prioritize site selection. In this case, highest priority samples are to be collected from oiled areas that are sensitive habitats, biologically productive, or highly relevant for human use. Collecting pre-oiled tissue samples from sensitive/productive sites that are likely to be oiled is also a priority. Sampling at unoiled “control” (upstream) sites is also a priority.
- Tissue samples should be collected pre-oiling, if possible, as soon as practical after oiling and periodically thereafter. Sampling frequency should be defined in the study design.
- Guidelines for the number of locations and sites to be sampled:
 - Minimum guidelines are at least 3-5 samples per location of relatively uniform exposure or distinct waterbody
 - If logistical limitations are a concern, prioritize sample collection by selecting a minimum of one reference/pre-oiled location, and two heavily oiled locations
- Bivalves that are considered special status species or species at risk should not be collected. [e.g. Western pearlshell (*Unionidae*), California floater (*Unionidae*), Oregon floater (*Unionidae*), Western ridged mussel (*Unionidae*), and the Montane peaclam (*Sphaeriidae*)]. To differentiate between these species and obtain additional information, use the NatureServe Explorer website

(<http://explorer.naturereserve.org>). In freshwater habitats, other bivalves, crayfish, or streambed benthic macroinvertebrates (e.g., New Zealand mud snail, caddisflies) may be collected.

- For exposure characterization, consider targeting species that are readily available across all locations and in quantities high enough for repeated sampling over time.
- Temperature can have a large impact on shellfish physiology. Some animals stop feeding or even passing water over their gills at low or high temperatures. Be aware of these differences when selecting species for monitoring and comparing results among species and over time.
- Uptake and depuration rates vary widely among species. Depuration usually takes weeks; thus, tissue sampling should be initiated within days after maximum exposure.
- Sample along exposure gradients, starting in the cleanest zone and then at regular intervals proportional to the exposure area.

Field Sampling Methods

Sampling Equipment/Containers

- Coolers – for sample storage and transport
- Blue ice packs/wet ice – for storage temperature regulation (if ambient temperature exceeds 4°C)
- Thermometer or temperature logger (1 per cooler)
- Disposable nitrile gloves
- Aluminum foil
- Ziploc or Whirl-Pak bags
- Dredges, knives, tongs, crayfish traps – washed and solvent cleaned, wrapped in foil and sealed for transport
- Shovel
- Rake
- Field Sample Forms (template in Appendix 9) and field notebook
- Chain of Custody forms (see Chain of Custody guideline in Appendix 9)
- Evidence tape (see Chain of Custody guidelines)
- GPS, camera (with spare batteries), photo scales
- Suitable disposal bags for oiled PPE and disposable sampling materials

Sample Collection Methods

- Use field data forms included in the work plan if one is available. Otherwise, use forms in Appendix 9. Coordinate data form development/modification with the NRDAR data management group.
- Because GPS units will be used to record locations and times, make sure that all GPS units are using the same coordinate system, datum, reporting units, and correct time. Follow the recommended GPS datum of the study plan. Alternatively, set the default to WGS84.
- Record the sampling site location using a GPS.
- Photograph the sampling site prior to sample collection to document the site conditions, as well as the sample collected. Make sure each photograph or series can be later associated with the corresponding sampling locations (e.g., through use of GPS Photo link software or by keeping a detailed photo log with waypoints and/or lat/long). Do not delete or alter any photographs. The numbering sequence of photographs uploaded from your camera must not have any gaps (see Field Photography guidelines, Appendix 8).
- Collect only live animals (e.g., for mussels, shells should be intact and tightly closed). Note the presence of dead animals on the field sample forms.

- If many dead or dying fish and/or invertebrates are observed, initiate the CDFW protocol for Shoreline Fish and Invertebrate Mortality Survey (**Appendix 11**) – collection and documentation without chemistry.
- If time is limited, note the location, species, life stage, visible oiling and approximate number or extent of the mortality event. Take pictures to document the die-off.
- Familiarize yourself with the target species and ensure that accurate taxonomic identification has been made during sample collection such that there is no mixing of closely related species. To meet the target of >10 g composite of soft tissue, a sample would need to approximately include 20 blue mussels 3-5 cm in length, 3 crayfish 6 cm in length, or 10 *Corbicula* clams 3 cm in length.
- Samples should be collected along the same vertical horizon or water depth, and from the same oiling exposure.
- Collect sample with gloved hands.
- If time is a concern, the primary target size range should include the larger individuals harvested at each sampling site.
- Record the presence of oil, weather conditions, water temperature, etc. in field notes.
- Bivalve sample collection:
 - Attached bivalves are pried away from the substrate with a knife, trowel, etc.
 - *Corbicula* can be sampled by digging into the stream sediment by hand or rake
 - Do not open shellfish in the field; collect the entire animal and shell
 - Wipe oiled shells with sorbent pads, wipes, etc. If heavily oiled, use a solvent damp wipe
- Subtidal/infaunal sample collection:
 - Deploy dredge from the side of the vessel that offers more space for safe deployment and retrieval, and that is further away from sources of contamination (e.g., fumes)
 - Record the GPS coordinates of the start/stop positions (when the dredge enters and before the dredge is brought onboard)
 - Dredge the sediment surface for 3 minutes at 2 knots in a circular pattern. Repeat this step at least 3 times per site (3 replicates), or until the desired target number of individuals is obtained
 - All individuals obtained via dredging MUST be collected during the same sampling event (2 hour sampling maximum)
 - Rinse with site water (clean water, if possible) to remove sediments except when oil sheens and slicks are present
- Crayfish sample collection:
 - Place 2 or more traps (see photograph) into the river, stream, or creek where flow energy is low and/or water is pooled. Traps must be placed in different locations, include bait (e.g., cat food), and be retrieved 24-hours after setting them
 - After retrieving the traps, remove collected crayfish. If missing limbs or cracked shells are visible, the crayfish are not useable and can be returned back to the water
 - Measure and record the carapace length of each crayfish (to the nearest millimeter). To do so, measure from the tip of the rostrum (point between the eyes) to the end of the carapace (end of main shell); do not include the tail segments in this measurement



- If necessary, rinse debris and sediment from them using (in order of preference) distilled water, clean tap water, or clean surface water
- Wrap composited tissue samples on a double layer of heavy-duty foil and placed directly into a double Ziploc-bag. Label the inner bag with marker pen and a waterproof sample label placed between the two bags. Use clear tape to protect the paper label. Make sure all organisms within the composite sample are stored together.
- Immediately place samples in coolers on ice and keep at 4°C.

Good Sampling Practices and Decontamination

- Good field practices and the development of a consistent sampling routine will ensure the integrity of the sheen samples and their validity in environmental assessments.
- Disposable nitrile gloves should be changed between samples to prevent cross contamination or if they become contaminated or damaged.
- To reduce the need for field decontamination, use pre-cleaned and/or disposable equipment and tools.
- The only equipment to be used between sites are a shovel and a dredge, which should be cleaned with soap and clean water. Repeated digging in clean sediments can be a last resort for cleaning the shovel if soap or clean water are not available. Alternatively, use a clean dry towel or other dry material to clean the shovel before its next use. Additional cleaning may be required when working at oiled sites (see below).
- Take precautions to avoid cross-contamination of the site from oil on personal equipment (e.g., boots, shovels). Sampling unoiled areas first, then lightly oiled areas and finally heavily oiled areas can minimize cross contamination. Personal equipment should be exchanged or cleaned between sites if it becomes contaminated.
- Potential sources of contamination while sampling from vessels (exhaust fumes, oily surfaces) are a concern. Work up-wind of any exhausts and designate clean areas for handling samples. Segregate dirty/clean areas. Layout clean surfaces to work on and replace frequently.

Sample Labeling and Record Keeping

- Verify that all samples are properly labeled and that field sample forms are properly filled out.
- Follow chain of custody procedures for securing samples and complete Chain of Custody forms (See Chain of Custody guidelines), noting where each tissue sample was collected, sampling equipment used, time/date of collection, size and container type, and sampler name.
- Make notations on the Chain of Custody form about any problems or observations during sampling.
- Maintain strict chain of custody during sample storage and transportation.
- Record the following on the field sample form:
 - Sample collection site (NRDAR sample grid ID and GPS coordinates)
 - Sample matrix (tissue)
 - Sample #, date/time, site location, tidal elevation, water depth
 - Species collected, number of individuals, size range, sample type (whole, tissue only)
 - Describe the oiling conditions (using standard shoreline assessment terminology), weather conditions (e.g., wind direction and speed), sediment characteristics, presence of biota, vegetation or debris, odors and other relevant information on the field data sheet
 - Sediment characteristics: grain size, texture, color, biota, vegetation, debris, odor, etc.; vertical changes in sediment characteristics
 - Record observations of any external evidence of contamination

- All sample numbers must be unique. The sample number should consist of a 2-letter Field Team name (e.g., ED for ephemeral data), 1-number field team number, 6-number date (MMDDYY), 2-letter sample type, and a 2-number sequential sample number. See the Sample Collection Summary Form for codes in Appendix 9.
- Keep a detailed photo log so that each photograph can be labeled.
- Note any deviations from the recommended guidelines in the field notebook.

Sample Preservation, Holding Times, and Shipping

- Follow chain of custody procedures for sample storage and shipping.
- Ship highly oil-contaminated samples separate from non-contaminated or low-contaminated samples to reduce risk of cross contamination. Ship samples directly to the laboratory as soon as practical, overnight (preferred), with completed Chain of Custody forms. If necessary, samples can be stored under specified conditions and with complete chain of custody until they can be shipped. Ensure that samples are packaged to protect them from breakage, shipping containers are sealed, and use ice packs or wet ice (if samples have been frozen after discussion with laboratory) to maintain storage temperatures during shipment to the lab.
- NEVER discard any samples even if these have exceeded their recommended holding times or storage temperatures.
- Refer to the Chemical Analysis Guidelines in Table 4 for analytical methods, sample volumes, minimum detection limits, and recommended holding time.

APPENDIX 5: Detailed Standard Operating Procedures for: Sediment

Standard Operating Procedure for Collecting Ephemeral Data: SEDIMENT

June 2023

Sampling Objectives

Characterize Oil in Sediment

- Determine the concentration and composition of oil compounds in sediments compared to background concentrations
- Determine the source of sediment contamination via chemical fingerprinting analysis and characterize oil weathering and fate (e.g., sediment toxicity testing)
- Estimate the areal extent and degree of sediment oiling (both along the shoreline and on the bottom of waterbodies)

Describe Habitat

- Measure sediment characteristics for interpreting chemical and biological results

Study Exposure

- Document exposure of sediment-dwelling organisms to oil compounds
- Support exposure modeling

Quality Assurance/Quality Control

- Ensure the integrity of the sample(s) throughout sampling, transport, and storage
- Ensure the reliability of chemical characterizations

Before Field Sampling

- Assure that all personnel have required safety training and protective equipment for field work.

Study Design

- It is important to have a defined sampling strategy prior to conducting fieldwork. Representative sediments are difficult to sample for oil contamination because of the inherent heterogeneity of oil distribution over space, depth, and time.
- The following terminology is used to define general to specific sampling geographies:
 - Area = general area of uniform characteristics, such as degree of oil exposure, physical setting, habitat types present, etc.
 - Location = a specific location that is representative of the area and contains the type of habitat to be sampled, such as a stream pool or shoreline of uniform oiling
 - Site = a specific point at which samples are collected or observations are made
- Plan the number of locations and number of sites per location, taking into account level of effort, potential logistical limitations, weather conditions, and other issues that may compromise sample integrity.

Equipment

- Review the list of sampling equipment/containers, make adjustments as needed, and ensure that all essential field materials are ready to be taken to the field.

- It may be necessary to coordinate with the laboratory that will receive the samples to ensure that acceptable materials and conditions are used for sampling and sample storage and shipping.
- Make sure that all essential equipment is in working order and operational and that spare equipment and materials are available.

Sampling Areas and Timing

- Follow a sampling plan/work plan if one is available.
- If a sampling plan is not available for ephemeral data collection immediately after a spill, data collection should focus on collecting samples from a range of unoiled, likely to be oiled, and already oiled areas.
- The number of locations and number of sites per location need to be considered, making sure that there is enough space in the coolers to accommodate all samples without sacrificing their integrity.
- Sediment samples can be collected from boats, the shoreline/stream bank, or wading in shallow water.

Area Selection

- Sampling locations should be representative of areas that have been or may be oiled by the spill and unoiled reference locations.
- Use trajectory models, overflight information, SCAT data, field observations, or other information to determine what areas have been oiled and which ones are likely to be oiled.
- Samples should also be collected from locations known or suspected to be impacted by other natural or anthropogenic sources of contamination (e.g., oil seeps, coal, peat, mining, combustion engines), as these will be important to differentiate background sources and levels of contamination.
- It may be necessary to prioritize sampling locations. In this case, highest priority samples are to be collected from oiled locations that are sensitive habitats, biologically productive, or highly relevant for human use. Collecting pre-oiling sediments from sensitive/productive locations that are likely to be oiled by the spill in the near future is also a priority. Sampling at unoiled “control” locations and sampling other sources of contamination should be prioritized based on the ephemerality of the data and relative importance to developing a NRDAR case.
- The number of locations and number of sites per location should be defined in the study design. A minimum guideline for collecting sediment samples is at least three sites per location of relatively uniform oiling exposure.
- Sample along exposure gradients, starting in the cleanest zone and then at regular intervals proportional to the exposure area.

Field Sampling Methods

Sampling Equipment/Containers

Note: The amount of equipment required depends on the sampling plan, desired sample volumes, and logistics. Analytical laboratories may provide required sampling and sample storage and transport materials – contact the receiving lab before preparing to collect samples in the field.

- Coolers – for sample storage and transport
- Blue ice packs/wet ice – for storage temperature regulation
- Thermometer or temperature logger (1 per cooler)
- Disposable nitrile gloves

- Sampling jars – 8 oz certified organic-clean jars with Teflon-lined lids and labels – for TPH/PAH/biomarkers
- Zipper-top (up to 1 gallon) or Whirl-Pak bags – for grain size and TOC samples
- Underwater sediment sampling devices. e.g., modified van Veen grab, Ekman grab, or ox dredge
- Coring tubes – single-use, disposable preferred
- Flat scoops (stainless steel or plastic) for shallow water, banks, and shorelines
- Stainless steel spatulas and spoons – cleaned, wrapped in foil, and sealed for transport
- Disposable aluminum pans – for composite samples
- Aluminum foil
- Sorbent pads
- Field Sample Forms (template in Appendix 9) and field notebook
- Chain of Custody forms (Appendix 9)
- Evidence tape (see Chain of Custody guidelines)
- GPS, camera (with spare batteries), and photo scales
- Packaging materials for glass jars (e.g., bubble wrap, sorbent pads, tape) – may be provided by the analytic laboratory
- Suitable disposal bags for oiled PPE and disposable sampling materials

Sample Collection Methods

- Use field data forms included in the work plan if one is available. Otherwise, use forms in Appendix 9. Coordinate data form development/modification with the data management group.
- Because GPS units will be used to record locations and times, make sure that all units are using the same coordinate system, datum, reporting units, and correct time. Follow the recommended GPS datum of the study plan. Alternatively, set the default to WGS84.
- Record the sampling site location using a GPS.
- Photograph the sampling site prior to sample collection to document the site conditions, as well as the sample collected. Make sure each photograph or series can be later associated with the corresponding sampling locations (e.g., through use of GPS Photo link software or by keeping a detailed photo log with waypoints and/or lat/long). Do not delete or alter any photographs. The numbering sequence of photographs uploaded from your camera must not have any gaps (see Field Photography guidelines, Appendix 8). If taking photographs of the sampling site is impractical, only take photographs of the sample collected.
- Surface sediment samples can be collected by using a pre-cleaned stainless steel scoop, wooden spoon, or by hand wearing clean gloves to collect the selected interval (e.g., 0-5 cm) from 3 locations and composite them using the aluminum tray before placing in a pre-cleaned 8-oz glass jar.
- Any sediment sampling device for collecting samples under water which meets the following requirements can be used:
 - Creates a minimum bow wake when descending
 - Penetrates the sediments to below the desired sampling depth
 - Closes to form a leak-proof seal after the device is triggered to close
 - Prevents sediment washout and disturbance when ascending
- Clear surface slicks prior to deploying the device by sweeping the area with a sorbent pad or placing a barrier up-current to divert surface oil around the sampler deployment area.
- Collect sample with gloved hands.
- When deploying and retrieving the sampling device under water:

- Lower and retrieve the sampling device at a controlled speed of ~30 cm per second to minimize potential bow wake activity and bottom disturbance as the sampler contacts the bottom, and loss and disturbance of the subtidal sediment sample during retrieval
- The device should contact the bottom gently, making sure it settles flat; only its weight or piston mechanism should be used to penetrate the sediment. It is important to minimize disturbance to the surface floc, which is likely to contain oil contaminants if they are present
- Secure the sampler on board and examine the sample for acceptability as follows:
 - The sampler is not overfilled; the sediment surface is not pressed against the sampler top
 - Overlying water is present, indicating minimal leakage
 - Sediment surface is undisturbed, indicating lack of channeling or sample washout
 - The desired penetration depth is achieved (e.g., 4-5 cm for a 2 cm sample)
- Siphon off the overlying water near one side of the sampler
- Using a flat scoop, accurately collect the top 2 cm from the center of each grab, avoiding sediments in contact with the sides of the sampler. Collect other intervals, as needed, using a new scoop for each sampling interval
- Make a composite sample of at least three subsamples within the same site and sampling interval, using a disposable aluminum pan or aluminum foil-lined container to homogenize the samples
- If time, equipment, or logistics do not allow collection of composite samples, collect grab sediment by following steps above, omitting the collection and mixing of subsamples. Place grab samples directly into jars for mixing, homogenizing and splitting in the laboratory
- In addition to collecting samples for chemical analysis, take samples for TOC and grain size (placed in Ziploc or Whirl-Pak bags)

Quality Assurance/Control

- Obtaining an adequate number of quality control samples is essential. At a minimum, a trip blank (accounts for contamination introduced during shipping and handling) and field blank (accounts for contamination introduced during sampling) should be maintained for each sampling effort and generally be collected at a rate of 5% and 10%, respectively, of all samples.
- A trip blank is an unopened sampling jar and should be transported with the samples and remain sealed in the cooler during sampling activities.
- A field blank should be collected at approximately every third sampling site, or at least at an “unoiled” and “oiled” site, by leaving the field blank sample jar open for the duration of the sampling period at that site. Record the site where field blanks were taken on the field sample form.
- Ideally, kiln-fired sand supplied by the laboratory can be transferred (poured or scooped) from one jar to another and returned to the lab as a field blank, but if this is not possible, use the open-jar technique (empty jar).
- Duplicate samples should be collected at every third sampling site or following the specifications of the work plan. A duplicate sample is collected from the same location and following the same steps as the preceding sample. This is not the same as collecting replicates from each site/depth. Duplicates should account for 10% of all samples, but consideration should be given to sample storage capacity. Do not split samples unless specified in the work plan.

Good Sampling Practices and Decontamination

- Good field practices and the development of a consistent sampling routine will ensure the integrity of the samples and their validity in damage assessments.

- Disposable nitrile gloves should be worn when sampling and changed between each sample collected or as necessary to prevent cross contamination.
- To reduce the need for field decontamination, use pre-cleaned and/or disposable equipment and tools.
- Sediment samples for THC and PAH analysis should be placed in certified organic-clean (solvent rinsed) glass containers with Teflon- or aluminum foil-lined lids. Samples for grain size and TOC can be placed in Ziploc or Whirl-Pak bags.
- Sediment sampling devices (e.g., dredges, coring tubes) must be decontaminated between samples:
 - Wipe off as much bulk oil as possible using sorbent pads
 - Wash device with laboratory-grade detergent and clean with a triple clean-water rinse. Cleaning with laboratory-grade water is preferred, though store-bought distilled water is a less ideal alternative and, as a last resort, “background” water from an up-current clean area can be used
 - Rinse with methanol or acetone, followed by hexane (Capillary GC Pesticide Residue Grade or equivalent). Collect solvent rinsate for proper disposal. Allow solvents to evaporate from equipment before use. If solvents are not available, use a diluted detergent solution and fresh water, followed by a distilled water rinse
- Take precautions to avoid cross-contamination of the site from oil on personal equipment (e.g., boots, shovels, etc.). Sampling unoiled areas first, then lightly oiled areas and finally heavily oiled areas can minimize cross-contamination. Personal equipment should be exchanged or cleaned between sites if it becomes contaminated.
- Potential sources of contamination while sampling from vessels (exhaust fumes, oily surfaces) are a concern. Work up-wind of any exhausts and designate clean areas for handling samples. Segregate dirty/clean areas. Layout clean surfaces to work on and replace frequently.

Sample Labeling and Record Keeping

- Verify that all samples are properly labeled and that field sample forms are properly filled out.
- Follow chain of custody procedures for securing samples and complete chain of custody forms (See Chain of Custody guidelines, Appendix 9), noting where each subtidal sample was collected, sampling equipment used, time/date of collection, size and container type, and sampler name.
- Make notation on the Chain of Custody form about any problems or observations during sampling.
- Maintain strict chain of custody during sample storage and transportation.
- Record the sample number on both the sample jar label and lid. All sample numbers must be unique. The sample number should consist of a 2-letter Field Team name (e.g., ED for ephemeral data), 1-number field team number, 6-number date (MMDDYY), 2-letter sample type, and a 2-number sequential sample number. See the Sample Collection Summary Form for codes in Appendix 9.
- If sample volume is split between two jars, both jars should receive the same sample ID and be recorded on a single line of the Chain of Custody form.
- Keep a detailed photo log so that each photograph can be labeled.
- Note any deviations from the recommended guidelines in the field notebook.

Sample Preservation, Recommended Holding Times and Shipping

- Follow chain of custody procedures for sample storage and shipping.

- Immediately place all sediment samples for chemical analysis in a cooler and keep at approximately 4°C. Use frozen gel packs or wet ice to maintain the temperature if ambient temperatures are above freezing. A programmable temperature logger or thermometer should be placed in each cooler to maintain a record of storage temperatures.
- Refrigerate (do not freeze) samples for TOC. Samples for grain size do not require refrigeration.
- Protect samples for chemical analysis from direct sun exposure (e.g., UV radiation).
- Samples for TOC and grain size can be stored separately from samples for chemical analysis.
- Tape lids on sample bottles in accordance with chain of custody guidelines so they do not accidentally come off.
- Store samples for chemical analysis from unoiled areas in one set of coolers, with oiled samples in a separate set of coolers.
- Use packing material, such as bubble wrap or sorbent pads, around glass jars to prevent breakage during transport and shipping. Take special care with gravel sediments because individual pieces can rattle around during shipping and break the glass jars.
- Freeze samples for chemical analysis as soon as practical or by the end of each day if samples are not going to be analyzed within 7 days of collection.
- Ship samples directly to the laboratory as soon as practical with complete Chain of Custody forms. If necessary, samples can be stored under specified conditions and with complete chain of custody until they can be shipped. Ensure that samples are packaged to protect them from breakage, shipping containers are sealed, and use ice packs to maintain storage temperatures during shipment to the lab.
- Ship highly oil-contaminated samples separate from non-contaminated or low-contaminated samples to reduce risk of cross contamination.
- NEVER discard any samples even if these have exceeded their recommended holding times or storage temperatures.

APPENDIX 6: Detailed Standard Operating Procedures for: Human/Recreational Use

Standard Operating Procedure for Collecting Ephemeral Data: HUMAN AND RECREATIONAL USE

June 2023

Guideline Objectives

The primary objective of this document is to provide guidelines for collecting ephemeral data relating to human and recreational use of natural resources during the early stages of an oil spill to support Natural Resource Damage Assessment and Restoration (NRDAR).

Background

Human and recreational uses of natural resources may be impacted as a result of an oil spill and the ensuing incident response (e.g., closure of recreational areas). Many of these impacts may be relatively short-term and difficult or impossible to document later. The following sections outline ephemeral data that local land managers can consider collecting to assess potential human use impacts in the immediate aftermath of a spill. What types of data can be collected will depend on the available time and experience of land managers or other local staff. Early data collection, even if qualitative, is important and can guide the development of more quantitative studies if appropriate.

Document Pathways

- Photo documentation:
 - Take pictures of oiled beaches/sites, advisory signs (“Warning: Do not contact water or eat fish”), closure signs/cordoned off areas, presence of response workers, and similar factors that may limit use of the area.
 - Follow the “Field Documentation Using Photography” guidelines found in Appendix 8 where possible.
 - At a minimum, include landmarks in pictures to assist with geolocation and with time-series images.
- Media documentation:
 - Save news articles and social media posts about the event made by health officials, towns, recreational users themselves, etc.
 - These may not stay online forever, so save all media on an electronic resource that is regularly backed up.
 - Online sources can be saved as screenshots (social media posts), saved locally as a static webpage (online news articles and the like), printed to PDF, or otherwise electronically archived.
 - Physical news articles or other documentation (e.g., informational flyers, community meeting posters) should be scanned or photographed and saved electronically with other media documentation.

Collect Basic Information about Recreational Use

These observation and interview activities are intended to provide information to guide development of comprehensive surveys of recreational and other use at affected sites. These activities are optional and are applicable depending on the available time and experience or background of staff.

- Observations of recreational users:
 - Count recreational users present at sites and define the boundaries of the count using permanent landmarks (e.g., between two jetties).
 - Observe and record what recreational users are doing and not doing, as applicable (e.g., playing volleyball, birdwatching, and sunbathing; nobody fishing or wading/swimming).
 - Conduct counts and record observations once or twice a day after the incident, targeting different times of day.
 - If feasible, implement data collection that can easily be repeated on the anniversary of the spill and document the process in a protocol document.
- Informational/qualitative interviews with recreational users
 - Talk to recreational users at sites about what they usually do at the site and how the spill may be affecting them.
 - Are they doing something else than they had planned?
 - Are they enjoying their experience less than usual?
 - Do they know of people who are avoiding the site?
- Information gathering from site managers (beach lifeguards, parks staff, etc.)
 - Ask for their impressions about how recreation use levels and activities are being affected, if at all, by the spill.
 - Ask them to compare that to the days leading up to the spill or during the same time of year without the spill.
 - If they have data on recreation use levels leading up to the spill, make sure they keep those data and continue collecting the data using the same methods in the days following the spill.
 - Note: resource managers themselves such as parks staff may be the ones documenting this information. It's important to still document the information and store with other electronic records, even if it is a "self" interview.

More Detailed Guidance and Templates

NOAA developed the current “Best Practices for Collecting Onsite Data to Assess Recreational Use Impacts from an Oil Spill” (2017). Appendix D provides example datasheets that resource managers may find useful for more systematic data collection (pp. 74-89). For incidents that may need a more stringent process and higher level of documentation than described above, there is a helpful summary, “Putting it All Together,” which covers the five basic steps (pp. 49-50):

1. Create a Sampling Plan
2. Develop Data Collection Materials
3. Hire and Train Personnel
4. Implement Data Collection
5. Prepare Data for Analysis

Resource managers should consult with agency economists about whether and how to implement a more robust onsite data collection effort following the best practices for these steps.

APPENDIX 7: Detailed Standard Operating Procedures for: Benthic Macroinvertebrates

Standard Operating Procedure for Collecting Ephemeral Data: BENTHIC MACROINVERTEBRATES

June 2023

Sampling Objectives

Study Injury

- Document the stream benthic macroinvertebrate community exposed to the oil.

Quality Assurance/Quality Control

- Ensure the integrity of the sample(s) throughout sampling, transport, and storage.
- Ensure the reliability of biological characterizations.

Before Field Sampling

- Ensure that all personnel have required safety training and protective equipment for field work.

Study Design

- It is important to have a defined sampling strategy prior to conducting fieldwork.
- The following terminology is used to define general to specific sampling geographies:
 - Area = general area of uniform characteristics, such as degree of oil exposure, physical setting, habitats present, etc.
 - Location = a specific location that is representative of the area and contains the type of habitat to be sampled, such as a stream reach or tidal marsh
 - Site = a specific point at which samples are collected or observations are made
- Plan the number of locations and number of sites per location, accounting for level of effort, potential logistical limitations, weather conditions, and other issues that may determine the actual sampling effort. Generate a list of alternate sites when unforeseen circumstances prevent sampling at the primary site(s).
- The sampling strategy should have enough flexibility that it can be adjusted based on conditions in the field.
- Consult appropriate guidelines for the collection of other environmental media and biota concurrent with benthic macroinvertebrate sampling. Tarballs, sheens, shellfish tissues, or sediments can be collected opportunistically for chemical analysis and fingerprinting.

Equipment

- Review the list of sampling equipment/containers, adjust as needed and ensure that all essential field materials are ready to be taken to the field.
- Do as much material preparation as possible prior to going into the field, including: labeling sample jars using permanent markers or laboratory labels, pre-cleaning sampling tools, etc.
- Make sure that all equipment is in working order and spare equipment and materials are available.

Sampling Areas and Timing

- Follow a sampling plan/work plan if one is available.

- If a sampling plan is not available, data collection should focus on collecting samples from a range of unoiled, likely to be oiled, and already oiled areas.
- The number of locations and number of sites per location need to be considered, making sure that there is enough space in the coolers to accommodate all samples without sacrificing their integrity.

Area Selection

- Sampling locations should be representative of areas that have been or may be oiled as well as unoiled reference locations.
- Use your own observations, trajectory models, overflight observations, SCAT data, etc. to determine what locations have been oiled and which ones are likely to be oiled.
- Samples should also be collected from locations known or suspected to be impacted by other natural or anthropogenic sources of contamination (e.g., oil seeps, coal, mining, combustion engines), as these will be important to differentiate background sources and levels of contamination.
- It may be necessary to prioritize site selection. Highest priority samples are to be collected from oiled areas that are sensitive habitats, biologically productive, highly relevant for human use, or have historical data. Collecting pre-oiled tissue samples from sensitive/productive sites that are likely to be oiled is also a priority. Sampling at unoiled “control” (upstream) sites is also a priority.
- Sample along exposure gradients, starting in the cleanest zone and then at regular intervals proportional to the exposure area.

Field Sampling Methods

Sampling Equipment/Containers

- Coolers – for sample storage and transport
- Disposable nitrile gloves
- D-frame net with clean replacement nets
- 500 mL wide-mouth plastic sample jars
- Digging tool: rebar or an abalone iron
- 95% ethanol solution
- White tray
- Forceps
- Field Sample Forms (template in Appendix 9) and field notebook
- Chain of Custody forms (see Chain of Custody guideline in Appendix 9)
- Evidence tape (see Chain of Custody guidelines)
- GPS, camera (with spare batteries), photo scales
- Suitable disposal bags for oiled PPE and disposable sampling materials

Sample Collection Methods

- Use field data forms included in the work plan if one is available. Otherwise, use forms in Appendix 9. Coordinate data form development/modification with the NRDAR data management group.
- Because GPS units will be used to record locations and times, make sure that all GPS units are using the same coordinate system, datum, reporting units, and correct time. Follow the recommended GPS datum of the study plan. Alternatively, set the default to WGS84.
- Record the sampling site location using a GPS.

- Photograph the sampling site prior to sample collection to document the site conditions, as well as the sample collected. Make sure each photograph or series can be later associated with the corresponding sampling locations (e.g., through use of GPS Photo link software or by keeping a detailed photo log with waypoints and/or lat/long). Do not delete or alter any photographs. The numbering sequence of photographs uploaded from your camera must not have any gaps (see Field Photography guideline).
- At each stream reach, three samples will be collected at 25%, 50%, and 75% of the stream width and composited into a single sample for that reach.
 - At the selected stream reach to be sampled, identify a point that is 25% of the stream width from the left bank. If it is not possible to collect a sample at the designated point because of deep water, obstacles, or unsafe conditions, adjust the sampling spot while keeping the point as close as possible to the designated position. Always be as objective as possible when identifying the sampling spot; resist the urge to sample the “best looking” or most convenient area of the streambed.
 - Once the sampling spot is identified, place a 500- μm D-frame net (see adjacent photo) in the water 1 meter downstream of the target spot. Position the net so its mouth is perpendicular to, and facing into, the flow of the water.
 - Holding the net in position on the substrate, visually define a square shape (a “sampling plot”) on the stream bottom upstream of the net opening, approximately one net-width wide and one net-width long. Because standard D-nets are 12 inches wide, the area within this plot is 1ft^2 (0.09 meter^2). Restrict sampling to within that area.
 - Working backward from the upstream edge of the sampling plot, check the sampling plot for heavy organisms such as mussels, caddis cases, and snails. Remove these organisms from the substrate by hand and place them into the net. Carefully pick up and rub stones directly in front of the net to remove attached animals. Pick up and clean all the rocks larger than a golf ball within the sampling plot such that all the organisms attached to them are washed downstream into the net. Set these rocks outside the sampling plot after they have been cleaned. Large rocks that protrude less than halfway into the sampling area should be pushed aside. If the substrate is consolidated, bedrock, or comprised of large, heavy rocks, kick and dislodge the substrate (with the feet) to displace benthic macroinvertebrates (BMIs) into the net. If a rock cannot be removed from the stream bottom, rub it with your hands or feet (concentrating on cracks or indentations), thereby loosening any attached insects. While disturbing the plot, let the water current carry all loosened material into the net. Do not use a brush to dislodge organisms from substrates.
 - Once the coarser substrates have been removed from the sampling plot, dig through the remaining underlying material with fingers or a digging tool (e.g., rebar or an abalone iron) to a depth of about 10 centimeters (less in sandy streams), where gravels and finer particles are often dominant. Thoroughly manipulate the substrates in the plot to encourage flow to dislodge any resistant organisms. Note: the sampler may spend as much time as necessary to inspect and clean larger substrates, but should take a standard time of 30 seconds for the digging portion of this step. To the extent practical, reduce the amount of sand particles in the net, as they damage organisms and degrade taxonomic data quality.



- For slack-water habitats, vigorously kick the remaining finer substrate within the plot using the feet while dragging the net repeatedly through the disturbed area just above the bottom. Keep moving the net so that the organisms trapped in the net will not escape. Continue kicking the substrate and moving the net for 30 seconds. For vegetation-choked sampling points, sweep the net through the vegetation within a 1-ft² (0.09 meter²) plot for 30 seconds. After 30 seconds, remove the net from the water with a quick, upward motion to wash the organisms to the bottom of the net.
- Dip the net into the water several times to remove fine sediment without loss of any animals. Place the collected material in a 500 mL wide-mouth plastic sample jar. Perform this operation over a large, white tray to avoid loss of any sampled material and make recovery of spilled organisms easier. If possible, remove the larger twigs and rocks by hand after carefully inspecting for clinging organisms. Use forceps to remove any organisms clinging to the net and place these in the sample jar. All samples should be completely transferred to the sample jar without elutriation. Then repeat at the mid-channel (50% of stream width) location and the adjacent to the right bank (75% of stream width). Place all BMI samples into the same jar (though never fill a jar more than halfway) and label.
- When the net becomes contaminated with oil, it will be difficult to clean it between sampling sites. Therefore, it is important to have replacement nets available. Oiled nets may be cleaned at the end of the day, but cleaning will require solvents and proper disposal of waste solvents.

Good Sampling Practices and Decontamination

- Disposable nitrile gloves should be changed between samples to prevent cross contamination or if they become contaminated or damaged.
- To reduce the need for field decontamination, use pre-cleaned and/or disposable equipment and tools.
- Take precautions to avoid cross-contamination of the site from oil on boots, digging tools, or other equipment. Personal protective equipment should be exchanged or cleaned between sites if it becomes contaminated.

Sample Labeling and Record Keeping

- Place a date/locality label, filled out in pencil, on the inside of the jar, and completely fill the jar with 95% ethanol. Note that the target concentration of ethanol is 70%, but 95% ethanol is used in the field to compensate for dilution from water in the sample. Final concentration of ethanol can be confirmed in the laboratory upon receipt of samples. To ensure proper preservation of BMIs, gently rotate jars that contain mostly mud or sand so that the ethanol is well distributed. Affix a second waterproof label on the outside of the jar. Fill the label out in pencil. Tape the label with transparent tape. Make sure all samples have both internal and external labels.
- Samples with an abundance of sand or organic material should be processed expeditiously at the lab, as specimens in these samples can degrade quickly. Therefore, the presence of these kinds of samples should be communicated to the taxonomy lab as soon as possible and they should not be stored for a long time before delivering to the taxonomy lab for processing.
- Verify that all samples are properly labeled and that field sample forms are properly filled out.
- Follow chain of custody procedures for securing samples and complete chain of custody forms (See Chain of Custody guidelines), noting where each sample was collected, sampling equipment used, time/date of collection, size and container type, and sampler name.
- Samples with a lot of organic material (e.g., plants, algae, leaf litter) tend to contain a lot of water that may inhibit sample preservation. Gently squeeze out as much water as possible (through the mesh of the D-frame net) before placing the sample in the jar, to prevent diluting

the alcohol too much. Approach this task gingerly, so as not to damage invertebrates during this process.

- Make notation on the Chain of Custody form about any problems or observations during sampling.
- Maintain strict chain of custody during sample storage and transportation.
- Record the following on the field sample form:
 - Sample collection site (NRDAR sample grid ID and GPS coordinates)
 - Sample matrix (BM)
 - Sample #, date/time, site location, tidal elevation, water depth
 - Describe the oiling conditions (using standard shoreline assessment terminology), weather conditions (e.g., wind direction and speed), sediment characteristics, presence of dead biota, vegetation or debris, odors, and other relevant information on the field data sheet
 - Record observations of any external evidence of contamination on the biota
- All sample numbers must be unique. The sample number should consist of a 2-letter Field Team name (e.g., ED for ephemeral data), 1-number field team number, 6-number date (MMDDYY), 2-letter sample type, and a 2-number sequential sample number. See the Sample Collection Summary Form for codes in Appendix 9.
- Keep a detailed photo log so that each photograph can be labeled.
- Note any deviations from the recommended guidelines in the field notebook.
- Ship samples directly to the laboratory as soon as practical, overnight (preferred), with completed Chain of Custody forms.

APPENDIX 8: Detailed Standard Operating Procedures for: Field Documentation Using Photography

Standard Operating Procedure for Collecting Ephemeral Data: FIELD DOCUMENTATION USING PHOTOGRAPHY

June 2023

Guideline Objectives

The primary objective of this document is to provide guidelines on procedures for taking photographs and recording videos for ephemeral data and samples collected in the field during the early stages of an oil spill to support Natural Resource Damage Assessment and Restoration (NRDAR) exposure and injury evaluations.

Background

Photographs and video are taken in the field to document the pre-oiling and oiling conditions and are key pieces of information that can be introduced as evidence. Each photograph should tell a specific part of the story. Before taking a photograph, you should consider what critical information you are trying to convey. Did the photograph capture the details you need? Are there key images (data) that you have missed? In the first case, you should take a better photograph. In the second case, you should look for photographs that will fill in gaps.

Document the Incident, the Location, and Use

- Photographs are taken to visually communicate what happened at a specific location or sampling site.
- Because photographs can be later viewed by various audiences (e.g., upper-level management, Congressional hearings, courts, the USCG National Pollution Fund Center, public hearings, training talks, outreach events, etc.) try to capture photographs for all types of audiences.
- Take as many photographs as needed. You may not get a second chance.
- The following is a partial list of subjects to always document:
 - The general site conditions and location
 - How the oil spill happened – including oil source
 - Oil on the water surface, stranded on the shoreline, and in direct contact with or over sensitive areas such as shallow pools, in-stream debris, and dry stream beds
 - Oiled wildlife
 - Oil recovery and cleanup operations (Response)
 - NRDAR staff working/sampling
 - Species presence and habitat use
 - Site use by humans (kayaking, canoeing, fishing)
- Time series of photographs are helpful to document exposure or changes in oiling over time. Repeat photographs for a time series by standing in the same spot and direction.

Document the Injury and Cause of the Injury

Directly Observable Injury

- Photographing and video recording direct injury can be very effective. Take photographs and make videos that clearly show conditions that are or may be caused by oil exposure and response actions, including but not limited to:
 - Oil on biota
 - Dead or injured animals and plants
 - Aberrant animal behavior (best capture with a video recording)
 - Impacts of cleanup operations

Cause of Injury

- Photographs and video are good for documenting visible oil exposure and impacts to recreation and human use:
 - When taking photographs of oiled areas, include perspective shots that show the degree of oiling as oiling occurs. Repeat day-to-day, if possible. Do not rely solely on SCAT to record the presence of oil.
 - It is important to document response actions that impact biota (e.g., removing, crushing, re-oiling, hazing) and other resources (e.g., sediment disturbances, etc.).
 - Also document closures of beaches, waterways, access points for fishing or recreation, including but not limited to photographs of official closures (e.g., posted closure signs), congestion effects (e.g., response taking over boat ramps), and popular use areas showing little or no recreational use.



Qualitative and Quantitative Approach

- Using a systematic photographic process to document oiled areas, reference areas, and the transitions between them can be an effective approach for documenting direct exposure.
- Rigorous photo transect and photo quadrat techniques may be appropriate depending on the assessment.
 - Use the sampling designs used for manual transect and quadrat surveys
 - Include oiled and non-oiled sites, or gradients in a continuum from most heavily oiled to non-oiled

Before Going to the Field

- Use field data forms included in the work plan if one is available. Otherwise, use forms in Appendix 8, Attachment A. Coordinate data form development/modification with the data management group.
- Make sure you have assembled a full photography kit appropriate for NRDAR field work (See Appendix 8, Attachment A: Full Gear and Field Gear checklists).
- Make sure all photographic gear is ready and complete before going to the field, including:

- Fully charged batteries
 - Memory cards (SD cards) (see below)
 - Clean lenses
- It is extremely important than the designated photographer and all personnel taking a camera into the field are knowledgeable of key camera settings. The recommended camera settings are as follows:
 - Resolution – MAX
 - ISO – Auto (avoid higher than 400 unless you are an experienced photographer)
 - Mode – Program (P)
 - Time stamp – off, especially if you are using GPS-Photo Link. Unlike film, there is no need to clutter photographs and use up pixel space with a time stamp. That information is automatically recorded in “EXIF” data—which is part of the image file
 - Time – local time
 - Continuous picture numbering – Set to use a running count for file names even after changing or formatting memory cards
 - Daily folders – Set camera to create a new folder each day
 - Advanced settings (e.g., spot metering, custom white balance, etc.) – reset. It is a good idea to return these advance settings to auto or a general setting before you go into the field
 - Camera reset – Most cameras have a way to return all settings to the factory default values. This is useful if images are poor or you have been experimenting with different camera settings and you cannot determine what setting may be causing the problem
- Make sure there are sufficient memory cards. Prior to going into the field, keep in mind the following recommendations:
 - Use high-quality memory cards with large storage capacity. Get enough capacity for a whole day’s photography.
 - Be sure all memory cards are working properly and are compatible with the camera being used.
 - Changing cards in the field risks getting moisture, salt, and dirt on the memory card contacts and inside the camera.
 - Format memory card after downloading photographs, at the end of day. It’s better to format than to delete all photographs.
 - At all times protect the electric contacts of memory cards from dirt and mechanical deformation.
 - NEVER take out a memory card when the camera is still writing to it. Turn off the camera before changing memory cards.
 - Use a quality memory card reader.

Learn Basic Camera Functions

- Remember that each digital camera is different. It is critical that all camera users know how to use their assigned camera before going into the field.
- For most field purposes and weather conditions, compact (aka “point-and-shoot”) cameras are cheaper, easier to use, more portable, and more resistant to salt, moisture, sand, and other factors. If you are an experienced photographer, can wait for favorable weather conditions, or require photographs for quantitative analyses, single lens reflex (SLR) cameras can provide higher resolution. and better quality photographs. SLR cameras generally perform well under freezing temperatures.

- Under some circumstances, and when taking photographs or video of underwater habitats and resources, small drop cameras (e.g., GoPro) or other underwater cameras may be necessary to document these areas.
- Cameras with 7-10 megapixels are recommended.
- The following are some basic functions that everyone should know. Many cameras require you to be in “P” (program) mode (not “A” [auto]) to use these:
 - **Light metering:** Spot. At this setting the camera meters the exposure at a designated spot in the photo frame. Most cameras show the “spot” as a box or circle in the center of the viewfinder. Spot metering is helpful when photographing a subject is much darker or lighter than the rest of the frame.
 - **Light metering:** Exposure compensation (+/-) adjustment. This feature tells the camera to make the photograph lighter or darker. It works like the lighter-darker adjustment on most copy machines.
 - **White balance adjustments:** White balance settings help the camera adjust the colors in the photographs based on the type of light (fluorescent, incandescent, sunny, cloudy, etc.). Most of the time Auto White Balance (AWB) works fine, but sometimes the camera does not adjust correctly. Manually choosing the type of light can fix the problem.
 - **Review photographs:** Know how to use the camera display to review a photograph. Know how to zoom in on the photograph in the display screen to check focus, exposure, and other key details.
 - **Forced flash:** In dim light or harsh shadows you may need to force the camera to use the flash to avoid losing details.
 - **Continuous shooting:** Most cameras will shoot consecutive photographs while you hold down the shutter. This is sometimes helpful when trying to capture moving wildlife.
- Some cameras may have GPS capabilities. The use of these cameras reduce location errors when labeling photographs as the location information is attached to the photo data. Basic GPS capability is essential for all field work, including photography. There are several key functions you need to set including (see Appendix A for detail):
 - Local time zone
 - Datum
 - Track (wrap, interval)
 - WAAS (on), etc.

Note: When the GPS recording is enabled, the camera battery life is shortened considerably.

While in the Field

- At each sampling location or site where photographs are taken, use the GPS to record waypoints. This will help with GPS-photo synchronization and processing.
- It is important to take photographs of a sampling site using labeled photo scales (e.g., 15 cm, 6 inches). The photo scale should be in one of the corners, preferably the lower right (see photograph).
- Scales should have intermediate reflectance, not bright white. A bright scale object can cause the camera to underexpose the rest of the photograph.



- Use spot metering or camera flash to eliminate harsh shadows that can obscure details. Use one of each if you're not sure which is better. Remember that setting the exposure for shadows may wash out and lose detail in bright areas of the photograph.
- Every close-up should be followed by one or more wider-angle shots that will show the close up in the context of the rest of the area. The closer the initial shot the more perspective shots are needed.
- Use the following distances as a guideline:
 - Macro (field of view ≤12"), useful for species identification, fine detail, or injury documentation
 - Close-Up (<1 m), useful for general documentation of oiled biota and resources
 - Mid-Level (1-2 m – at an angle), useful when documenting groups of biota and oiling
 - Distant/landscape (>10 m), useful when documenting habitats and spatial patterns of oiling; it is best to have a person in the photograph for scale (at least two photographs, facing forward and to the side of the observed area)
- Change batteries as soon as they are low so the camera does not lose power just as you are taking a critical photograph.
- Use the review feature to ensure that photographs show what you need.
- Use the zoom in function to see if you captured necessary details.
- Note key photographs and important details in the field notebook.
- Record basic information - locations, times, photographer, team members, including descriptions of GPS locations or waypoints.

Taking Photographs of Quadrats

- You may take high-resolution photographs of sampling quadrats for quantitative analysis.
- Quadrats should not be bright white. Make quadrats out of grey PVC (see photograph) or wrap white quadrats in grey colored duct tape.
- All photographs of quadrats must include a label containing the location name, transect, quadrat number.
- Take high-resolution, directly overhead photographs of each quadrat and record GPS coordinates. When taking photographs:
 - High-resolution photographs must include all four sides of the quadrat as these will be used to digitally count individuals and measure their coverage on a computer screen
 - Photographs need to be relatively flat so that the entire quadrat falls within a similar focal plane, with minimal shadowing from crevices or projections.

Taking Panoramic Photographs

- Panoramas are often unnecessary but if you need a wide, detailed photograph do the following:
 - Keep photo edges parallel
 - Do not change "zoom" factor
 - Overlap photographs by about 30%
 - Place a scale or natural distinctive feature in each overlap area for accurate alignment



- Lock your elbows against your sides for stability and pan as close to horizontally as you can. Use a tripod or monopod if you have one

Taking Video

- A short video synopsis of a location can be very helpful later for relaying or reviewing the general layout of a location.
- It is important to take video recordings documenting oil mixing into the water column over rapids as these can have impacts on oil fate.
- Take 30-45 seconds to slowly pan through a site while narrating key features.

Taking Photographs While Flying

- Taking photographs from a plane or helicopter can be difficult and requires additional skills. Point and shoot cameras and phones can take good photographs from the air but SLRs typically perform better.
- When taking photographs from a plane or helicopter:
 - Do not wear bright clothing as these may reflect in the windows of the aircraft
 - Use manual focus to set cameras to infinity (∞). This avoids accidentally focusing on the window
 - Using image-stabilized cameras or lenses will help take good quality photographs
 - To prevent transmitting aircraft vibration to the camera, do not rest the camera on an aircraft window frame or other part of the aircraft. Instead, hold the camera with your arms braced against your legs or torso, or the camera held against your face
 - Avoid shooting through a bubble window
 - Smaller aircraft often have sliding windows, or easily removable windows or doors; Make arrangements with your pilot before take-off
 - Avoid taking photographs towards the sun
 - Record on the field notebook the basic flight plan including altitude and track line, aircraft type, pilot and passenger names, port or starboard



Taking Underwater Photographs

- When taking underwater photographs:
 - Ensure that the camera is set on the underwater mode, which is designed to filter some wavelengths
 - If possible, include a scale with each photograph
- Underwater cameras may be the only way to document the extent and degree of oiling (sunken oil mats) and impacts (dead fish that sank to the bottom) below the surface. A GoPro camera with waterproof housing can be deployed by wading in shallow, quiet water.

Upon Returning from the Field

Legally Defensible Photographs

- Creating a legally defensible photograph record requires:
 - Maintaining a complete photograph record. DO NOT delete photographs from the camera or from your computer before the official archive is created
 - Keep one set of photographs that are never opened. In practice this means transferring one copy of the photographs from the camera memory card to a computer and then to a DVD-R or CD-R (non-editable) without ever opening them. The resulting continuous set of photograph files that have not been opened will demonstrate that you have a full, un-edited, photograph record for the court
- When you return from the field, download all photographs to a computer. Before reviewing photographs on the computer (review = open):
 - Create a copy in the “Working” directory and one copy in the “Archive” directory
 - The “Working” directory is used to process photographs through GPS-linking software and to log all photographs. DO NOT rename files in the “Working” directory
 - The “Archive” directory MUST include unopened, un-editable copy of all photographs. Burn the “Archive” directory to DVD-R. Do this when you have enough to fill a CD/DVD or at some set interval (every 2-3 days), and make a copy of the CD/DVD
 - NEVER open the files stored in the “Archive” directory
 - Additional backup copies can be made to portable hard drives

Locate Photographs – GPS Linking

- Field photographers should always collect a GPS track while in the field.
- Be sure to take a clear photograph of the operating GPS screen showing the date and time to synchronize the photographs with the GPS track (see photograph). The ideal GPS photograph should clearly show the GPS clock in Hours, Minutes, and Seconds.
- With the sync photo and a track file, all photographs can be linked to a specific Lat/Long/Time using special software. However, a different team would likely be responsible for processing GPS-camera information. They will use the sync photo and downloaded track file (using the software that came with the GPS) to GPS-link these photographs.



Appendix 8 Attachment A

1. Supporting Documentation – Photography Checklist and Forms

Photography forms may be provided by the NRDAR lead; otherwise, use the attached form.

- Print the form on weather-resistant paper (if available). Make more than enough copies of the form before going into the field.
- Fill out forms with waterproof pen or permanent marker. Do not use pencil or biro (erasable) ink.
- Fill in blanks with “N/A” if data are not applicable or not available. Avoid leaving blank values on data forms.
- Do not erase or black out erroneous entries on the field data forms. Errors should be corrected by crossing out the entry with a single line and signing and dating the strike-through.

2. Photography Gear Checklist

Photography - Complete Gear List; F Indicates Field Gear

<input type="checkbox"/>	Camera <i>F</i>	With neck strap
<input type="checkbox"/>	Camera case <i>F</i>	Sized to hold all camera gear; plus polypropylene freezer bag, if appropriate
<input type="checkbox"/>	Memory cards <i>F</i>	1-2 extra depending on size – (e.g., 200-300 high resolution photographs, each)
<input type="checkbox"/>	Rechargeable batteries <i>F</i>	Camera batteries: 2 is OK, 3 is better. AA's two sets of rechargeable are OK – extra alkaline or lithium for GPS etc.
<input type="checkbox"/>	Battery charger	Appropriate for each type of rechargeable batteries
<input type="checkbox"/>	Lens cleaning kit <i>F</i>	(e.g., soft cloth)
<input type="checkbox"/>	Card reader	One that accepts many types of cards is preferred
<input type="checkbox"/>	Cable – Camera to PC	
<input type="checkbox"/>	Camera manual	Paper and pdf
<input type="checkbox"/>	Underwater housing/kit <i>F</i>	Optional - useful in rough weather and small boat ops.
<input type="checkbox"/>	Photo scale <i>F</i>	15 cm waterproof, 15 cm disposable. Avoid white or light colors.
<input type="checkbox"/>	Image viewing software	All PC's and many cameras have software for reviewing photographs
<input type="checkbox"/>	Image editing software	Optional. Good for processing photographs for presentations etc.
<input type="checkbox"/>	External hard drives	
<input type="checkbox"/>	PhotoLogger database	
<input type="checkbox"/>	GPS-Photo Link software	
<input type="checkbox"/>	DVD-Rs – NOT RWs	
<input type="checkbox"/>	Waterproof bag <i>F</i>	Dry sack or heavy duty zip-lock bags
<input type="checkbox"/>	Polarizing lens <i>F</i>	Optional – reduces glare and reflections
<input type="checkbox"/>	GPS <i>F</i>	
<input type="checkbox"/>	Field notebook <i>F</i>	

3. Photo Log

PHOTO LOG

Page ____A of ____



Incident/Case: _____

Field Team Name and #: _____

Date (MMDDYY): _____

This form must be filled out for photos taken in the field.

Field Team Lead/ Photographer Name: Phone # <i>(Indicate if a personal number):</i> Agency/Affiliation:	Camera Make/Model/ID: Time zone: GPS Make/Model/ID: Time zone:
--	---

Photos must include: GPS unit (displaying time and date), up coast, down coast, landward, seaward, tissue sample (with photo scale ruler), sample label(s) and misc. sampling photos

¹ GPS Waypoint; ² Sample ID: Field Team Name (2 letters); Field Team # (1 number); Date (MMDDYY); Sample Type (2 letters); Sample # (sequential numbering begin with 01)

Team Member Initials: _____ Date: _____

PHOTO LOG

Page ___B of ___



Incident/Case: _____

Field Team Name and #: _____

Date (MMDDYY): _____

Suggested Keywords – These are suggested keywords to describe your photos. Keywords are used when importing the photos, where they will be queried by field staff, management, or outreach staff. Please select keywords that are general enough to represent the photos. Specific details should be entered into the Description/Notes/Comments section above.

Response/NRDA Related	Response/NRDA Related	Habitats	Habitats	Organisms	Conditions
Barge	Oil-Stain/Coat	Agricultural land	Marine Water	Amphibian	Cold/ Cool
Barrel	Outreach	Barren land	Marine Protected Area	Birds	Drought
Boat	Overflight	Beach-Sandy	Marsh	Coral	Dry
Boom	Pipeline	Beach-Gravel	Riverine	Crab	Fire
Cleanup Operations	Pits/Trenches	Cave	Riparian	Dead Wildlife	Flood
Dispersant	Produced water	Chaparral	Rip-Rap	Fish	High elevation
Dredging	Railroad	Desert land	Rocky Shoreline	Insect	High tide
Drilling Platforms	Quadrat	Desert shrub	Sediment/Soil	Invertebrate -Aquatic	Hot
Ephemeral data	Response Vessel	Dry creek	Stream	Invertebrate -Terrestrial	Ice
GPS Unit	Sample Container	Dune	Streambed	Mammal-Marine	Leak
Grounding	Sampling	Ephemeral Wetland	Subtidal	Mammal-Terrestrial	Low tide
Human Uses	SCAT	Estuary	Tree dominated	Native/ Non-native	Rain/wet
In-Situ Burn	Skimmer	Forest	Vernal pool	Reptile	Snow
Marine Debris	Source Oil	Grassland	Wetland	Shellfish	Storm
NRDA	Sunken Vessel	Herbaceous		Threatened/ Endangered	Wind
Oil	Tank	High Sierra		Tree	
Oil-Emulsified	Tanker Ship	Intertidal		Vegetation-Aquatic	
Oil-Sheen	Transect	Kelp Bed		Vegetation-Riparian	
Oil-Surface Residue	Waste Site	Lake			

Photos & GPS Data Relinquished By	Photos & GPS Data Received By
Name Signature:	Name Signature:
Name Printed:	Name Printed:
Agency/Affiliation Name Printed:	Agency/Affiliation Name Printed:
Date/Time:	Date/Time:

I, _____ [*Data Intake Manager print name*], without modification, downloaded the photographs referenced on this form in accordance with the NOAA OR&R Data Intake Protocols and uploaded without modification to DIVER in the File Collection ID number _____ with the following Photo Zip file named _____ and GPS File named _____.

Signature

Date/Time

Team Member Initials: _____ Date: _____

**APPENDIX 9: Detailed Standard Operating Procedures
for: Chain of Custody and Sample Collection
Summary Form**

Standard Operating Procedure for Collecting Ephemeral Data: CHAIN OF CUSTODY

June 2023

Guideline Objectives

The primary objective of this document is to provide guidelines on chain of custody procedures for ephemeral data and samples collected in the field during the early stages of an oil spill to support NRDAR exposure and injury evaluations.

Background

Chain of custody procedures are followed to authenticate a sample from the time it is taken until the results are introduced as evidence. For the purposes of litigation, agencies must be able to prove the legal integrity of all samples and data introduced as evidence. This means that it is necessary to have an accurate written record to track possession, handling, and location of samples and data from collection through reporting. Chain of custody facilitates this verification process. Failure to follow chain of custody procedures in this guideline does not necessarily render data unusable; however, any deviations from the chain of custody guidelines should be noted. Assuring that proper chain of custody guidelines is followed is vital to assuring the integrity of the samples, and the data generated by the analysis of those samples.

Responsibilities

All samplers handling samples collected for NRDAR MUST follow this procedure when collecting, handling, and securing samples. All team leads and supervisors are responsible for ensuring that the designated custodian(s) understand this procedure and strictly adhere to it for all sampling events.

Important Definition

- **Chain of Custody Form:** A document detailing who is legally responsible for samples at any point in time from collection until the sample is received by the laboratory.
- **Custody:** A sample is in your custody when
 - It is in your actual physical control and presence
 - It is in your view after being in your possession
 - It is not in your physical presence, but is in a secure place of storage to which only you have access
 - It is not in your view or physical presence, but is secured in a place of storage or secure area to which only you and identified others have access
- **Secure Area:** An area in which entry is restricted by keyed lock or similar to a designated custodian

Equipment for Chain of Custody

- Sample labels
- Tamper resistant evidence tape (for small sample jars and large shipping containers)
- Permanent markers
- Chain of custody and field data forms
- Secure storage area

Sample Collection – Chain of Custody

Note: As few people as practical should handle the sample from when it is collected through laboratory analysis.

- Sample custody begins immediately after a sample is collected. The sampler who collected the sample is responsible for the preservation and integrity of the sample(s) until that responsibility is transferred to someone else and documented with the chain of custody form. This chain of custody form then travels with the sample(s) and is used to document any other transfers of custody.
- When a sample is taken, the sampler must:
 - Complete a sample tag or label that identifies each sample. Use waterproof ink and attach the label to the sample jar or container at the time the sample is collected. Labels should contain the following legible information:
 - Sample number
 - Sample type (e.g., sediment, water)
 - Sample containing hazardous goods (such as formalin used as a preservative) (if applicable)
 - Time/date of collection
 - Location
 - Sampler name(s)
 - Seal each sample jar or container with tamper resistant evidence tape. When sealing jars, the tape should connect the jar to the lid. The sample collector should sign and date evidence tape so that the signature is partially on the tape of both the lid and the jar
 - If the sample is collected in a container that is not tamper proof (such as a plastic bag) then the container should be sealed with tamper resistant tape, a serial numbered zip-tie, or other means of verifying that the container has not been opened. The sample should also be stored in a container that is appropriate for chain of custody, such as a box or cooler that can be sealed with tamper resistant tape and signed by the collector
 - If tamper resistant evidence tape is not available, use masking or duct tape and sign across the end of the tape
 - Enter each sample on the chain of custody form
 - Document the sample in the field data sheet, noting details about the sample that may be pertinent later during sample analysis and injury determination
- The sample collector is responsible for care and custody of the samples until they are turned over to an assigned custodian or properly dispatched to the receiving laboratory. All custodians must assure that each sample remains in his/her custody (as defined above) so that no one can tamper with it during the entire duration of their responsibility.
- When samples are turned over to a new custodian:
 - The current custodian must officially relinquish the samples by signing the chain of custody form
 - The new custodian must review the samples, ensuring that they are in good condition and that the sample IDs matches the chain of custody form. Any damage or deviation must be noted on the chain of custody form before the new custodian accepts the samples by signing the form
 - The former custodian must retain a copy of the full set of forms. The original chain of custody forms always stay with the samples

- Pack and seal samples in suitable containers to avoid damage. A sample seal should be attached across the lid of each shipping container in such a manner that the container cannot be opened without breaking the seal. This lock and/or seal are not to be removed until the shipping container is opened by the laboratory custodian or designee.
- Ship samples by registered courier. Other certified shipping services may also be used. Keep all shipping receipts as part of the permanent chain of custody documentation regardless of how samples are shipped. The shipper does not need to sign the chain of custody form.
- Couriers picking up samples at the airport, post office etc. should sign the shipping documents to acknowledge receipt of the samples.

Photographs

- Digital photographs can be used as evidence. Like with physical samples, the objective is to be able to ensure that the photographs are an accurate representation of what was seen in the field (see Field Photography guideline).
- It is important to protect the legal integrity of digital photographs stored on digital memory cards (SD cards), as well as the legal integrity of the SD card itself.
- Digital photographs taken in the field and information stored on a digital memory card or camera internal memory MUST NOT be deleted, no matter the quality or other issues that may arise.
- All digital photographs files MUST be stored sequentially on the SD card and not renamed.
- Photographs should be downloaded to a secured PC and copied to an un-editable format (e.g., CD/DVD).
- Always keep a back-up copy of all photographs.
- See the Field Photography guideline for more details.

Other Important Considerations

- **Custodian list:** Maintain a list of people who are custodians on samples that includes how each person is related to the assessment.
- **Multiple custodians:** If multiple qualified custodians have access to a secure sample storage area then it is not necessary to document change of custody between custodians. All custodians should be identified on the custodian list.
- **Chain on multiple sheets:** Starting chain of custody documentation on one form and continuing it on a second form for the same samples is not a break in the chain. Care must be taken to keep the forms together.
- **Broken chain:** If the chain is broken for any reason or if you foresee deviations from the procedures in this guideline, contact the legal team for guidance. Changes to the procedures can be made but should be informed by a lawyer. Do not discard ANY samples even if the chain of custody has been broken.
- **Samples on the same sheet are split:** If samples that were recorded on a single chain of custody form are split for shipping, clearly mark the original chain of custody form to show which samples were removed, and when and where they are going. Create a new chain of custody form for the samples that are shipped and include a copy of the original chain of custody form. Keep the original form and a copy of the new form with the samples that are not shipped.

CA OSPR CoC Form

The information below describes how to fill out a Chain of Custody (COC) form with specific information applicable to NRDAR sampling during marine oil spills using the standard FG 1000 (Rev. 9/01) COC form.

The boxes across the top of the form are needed to identify the sampler, how the samples are being transported, the person who should receive the results, and the general information about the spill.

- Sampler and Ph #, address, city, zip: Print the name, affiliation, and contact information (phone and address) of the sampler.
- Date Required/Reason: Need not be completed unless desired by a specific date. ASAP indicates to the lab that the case is to be analyzed, but not rushed or held.
- Shipped Via: Usually completed by stating “Delivered in person” or “Hand carried”. However, if transported by a third party carrier (e.g., UPS), indicate the name of the transporter in that space.
- Send Results to, Address, City, Zip: The name and contact information of the person to be sent the lab results, often the NRDAR case lead (e.g., Michael Anderson, Resource Restoration Program Manager).
- Copies to, Address, City, Zip: The name and contact information of additional people, if any, to receive
- copies of the lab results, such as sampler.
- Lab Number: Leave blank. This is for lab use only.
- Field Number: Leave blank. This is for lab use only.
- Lab Storage: Leave blank. This is for lab use only.
- Spill Title: Name of Spill (e.g., Alameda drill).
- Suspect: Suspect source (e.g., T/V Moon).
- Index-PCA: Spill-specific billing code for use by OSPR and/or CDFW laboratories.

The next level of boxes is more applicable to sampling done by CDFW during enforcement actions - Fish & Wildlife (date and region); Fish & Game Code Violation; Suspected or Potential Problem; and Routine Analysis. None of these fields must be completed during NRDAR field sampling. In addition, the boxes for water quality measurements (temperature, pH, and conductivity), would be left blank unless measured.

The main portion of the COC in the middle is needed to identify the sample information and the specific analyses that should be run by the laboratory.

- Sample Identification: The general sample labeling scheme is: [FieldTeamName # Date SampleType Sample#]. For example, the sample collected by the Rocky Intertidal (RI) team 1 (1) on February 26, 2011 (022611) that was the second (02) mussel (MU) sample taken that day would be RI1022611MU02.
- Collection: Date and Time: Record the date and time (military) when the sample was collected (e.g., 2/26/11, 1310).
- Analysis Requested: Put an “X” in the box for the appropriate analysis. Write in analysis if not already included on the COC (e.g., PAH, BTEX, TPH, TOC, grain size, % lipid, % moisture).
- Sample Type: Place an “X” in the box under the sample type (e.g., water, filtered water, soil (or sediment), or tissue. If the sample type is not listed, cross through one option and write in the appropriate type.
- Number of Containers: write in the number of containers (e.g., 1), below the type of container used for a given sample (e.g., Plastic, Glass, VOA Vial, Foil, other). If more than one, describe in

the Sample Identification and/or Comment/Special Instruction boxes how the lab should handle these samples.

- Preservation: In the blank column, identify the type of preservation used. Typically, this would be a cooler with cube ice; if so, write in "Wet Ice," and place an "X" in the box below.

Note: See the attached three examples of completed COCs for details on appropriate labeling for sediment, water, sheen, tarball, and tissue.

See below for recommendations on how to distinguish the following situations:

- Field Duplicates - use a different sample ID, but same location and analysis
- Multiple chemical analyses of a sample stored in one container - use the same sample ID (e.g., WC3041411TB01) and put an "X" for each analysis needed
- Multiple chemical analyses of a sample stored in multiple containers (same location/same time) - use the same sample ID for each, specify the number of containers on the label (e.g., 1 of 4). Put an X for each analysis needed and write in the number of containers below the type of container used for a given sample
- Backup samples - use different sample ID and note "hold unless needed" in the comments/special instructions (e.g., WC3041411WT04).

Next, in the bottom right, leave blank the boxes for Pollution Action Kit, Glove Size, and Hazmat Shipper Requested.

In the bottom right, include:

- Problem Description: write in the type of incident, such as "Oil Spill."
- Suspect/Incident Location: write in the appropriate suspect source and incident location (e.g., T/V Moon/Anchorage 9).
- Comments/Special Instructions: Examples might include: "Mussel Samples – Hold Until Further Notice; or "Homogenize sediment samples in the laboratory prior to analysis".

The bottom of the COC is completed by both parties when the samples are transferred, the person relinquishing and the person receiving the samples. The original COC stays with the samples and the person relinquishing the samples keeps a copy.

- Samples Relinquished By (Signature): Signature of sampler releasing the sample.
- Print Name: Sampler to print name.
- Date: Date relinquished.
- Samples Received By (Signature): Signature of person receiving the sample.
- Print Name: Printed name of person receiving the sample.

Date: Date relinquished



DFG REQUEST FOR ANALYSIS AND CHAIN OF CUSTODY RECORD

Page _____ of _____

Sampler <i>Dennis Ryan, CDFW-OSPR</i>	Ph # <i>(916) 323-6288</i>	Send Results To <i>Michael Anderson, CDFW-OSPR</i>				Lab Number													
Address <i>1700 K Street</i>		Address <i>1700 K Street, Suite 250</i>				Field Number													
City <i>Sacramento</i>	Zip <i>CA</i>	City <i>Sacramento</i>		Zip <i>95811</i>		Lab Storage													
		Copies To <i>N/A</i>				Spill Title <i>T/V Moon</i>													
Date Required/Reason		Address <i>N/A</i>				Suspect <i>T/V Moon</i>													
Shipped Via <i>Hand Delivered</i>		City <i>N/A</i>		Zip		Index-PCA <i>e.g., 6121-54303</i>													
<input type="checkbox"/> Fish & Wildlife Loss Date: _____ Region: _____ <input type="checkbox"/> DFG Code Violation: _____ <input type="checkbox"/> Suspected or Potential Problem <input type="checkbox"/> Routine Analysis		Analysis Requested >>>		Water Temp:	F or C	pH:	DO:	mg/L Conductivity:	μmhos/cm										
Sample Identification/Location <small>(Draw map on separate sheet if necessary)</small>		Collection		Petroleum Fingerprint	Trace Elements [Specify Below]	Pesticides [Specify Below]	PAH [Include alkylated PAHs]	BTEX		Sample Type	Number of Containers	Preservation							
		Date	Time							Water	Filter-Clean-Sheen	Soil Tarball	Tissue	Plastic	Glass	VOA Vial	Temp	Acid	Wet Ice
<i>WC3041411TB01 [tarball]</i> <i>WC3041411SH01 [sheen]</i> <i>WC3041411WT01 [water]</i> <i>WC3041411SH02 [sheen]</i> <i>WC3041411WT02 [VOA]</i> <i>WC3041411WT03 [trip/field blank]</i>		4/14/11	1130	X			X				X			1				X	
		4/14/11	1321	X							X			1				X	
		4/14/11	1340			X				X			1				X		
		4/14/11	1510	X						X			1				X		
		4/14/11	1523				X			X			2				X		
		4/14/11	1534			X			X			1				X			
Problem Description <i>T/V Moon Oil Spill NRDA</i> Suspect/Incident Location <i>T/V Moon at Anchorage 9, San Francisco Bay</i> Comments/Special Instructions <i>Water Samples - Extract: Hold Extracts Until Further Notice</i> <i>Hold WC3041411WT02 unless needed as backup</i>												Pollution Action Kit: Yeso Noo Glove Size: Large o Medium o Hazmat Shipper Requested: Yeso Noo							
Samples Relinquished By (Signature)		Print Name		Date		Received By (Signature)		Print Name		Date									
<i>Dennis Ryan</i>		<i>Dennis Ryan</i>		<i>4/14/2011</i>		<i>[person receiving samples]</i>		<i>[person receiving samples]</i>		<i>4/14/2011</i>									

LAB COPIES: WHITE, CANARY, PINK

SUBMITTER: GOLDENROD

FG 1000 (Rev. 9/01)

Pesticide Investigations Lab
1701 Nimbus Road
Rancho Cordova, CA 95670

Petroleum Chemistry Lab
1995 Nimbus Road
Rancho Cordova, CA 95670

Water Pollution Control Lab
2005 Nimbus Road
Rancho Cordova, CA 95670



DFG REQUEST FOR ANALYSIS AND CHAIN OF CUSTODY RECORD

Page _____ of _____

111701 Nimbus Road

1995 Nimbus Road
Poncha Cordova, CA 95670

2005 Nimbus Road
Brentwood, California 94513

LAB COPIES: WHITE, CANARY, PINK

SUBMITTER: GOLDENROD

FG 1000 (Rev. 9/01)



DFG REQUEST FOR ANALYSIS AND CHAIN OF CUSTODY RECORD

Page _____ of _____

LAB COPIES: WHITE, CANARY, PINK

SUBMITTER: GOLDENROD

FG 1000 (Rev. 9/01)

9-9



CDFW REQUEST FOR ANALYSIS AND CHAIN OF CUSTODY RECORD

Page _____ of _____

Pesticide Investigations Lab
1701 Nimbus Road
Rancho Cordova, CA 95670

Petroleum Chemistry Lab
1995 Nimbus Road
Rancho Cordova, CA 95670

Water Pollution Control Lab
2005 Nimbus Road
Rancho Cordova, CA 95670

LAB COPIES: WHITE, CANARY, PINK

SUBMITTER: GOLDENROD

FG 1000 (Rev. 9/01)

APPENDIX 10: Example Coordination Agreements

**KEYDET ENERGY
CORPORATION
123 Petroleum Road
Los Angeles, CA**

April 22, 2004

Charles McKinley
Field Solicitor
Office of the Solicitor
U.S. Dept. of Interior
1111 Jackson St., Ste. 735
Oakland, CA 94607

LCDR Reismer, JACG, USN Assistant
Office of the Staff of Judge Advocate
Commander Navy Region Southwest
937 North Harbor Drive
San Diego, CA 92131-0058

Wendy Johnson
Office of Spill Prevention and Response
California Dept. of Fish and Game
1700 K St., Ste 250
Sacramento, CA 95814

Katherine Pease
Office of General Counsel
NOAA
501 W. Ocean Blvd., Ste. 4470
Long Beach, CA 90802

Re: Funding Commitment for Joint Pre-assessment/Assessment Activities

This is to confirm that Keydet Energy Corporation (Keydet) wishes to participate with the Natural Resource trustees (trustees) who are in receipt of this letter in their pre-assessment and assessment of injuries to natural resources related to the oil spill which occurred on or about April 20, 2004 in the waters approximately 10 miles northwest of San Diego, California. In consideration of the trustees' agreement to allow Keydet to participate cooperatively in these activities, Keydet hereby agrees to pay the reasonable costs previously incurred or to be incurred by the Department of the Interior (including the U.S. Fish and Wildlife Service, Bureau of Land Management, National Park Service, Office of Environmental Policy and Compliance, and Office of the Solicitor), the State of California (including the Department of Fish and Wildlife and Office of Oil Spill Prevention and Response), the Department of Commerce (including the National Oceanic and Atmospheric Administration), and the Department of Defense (including the U.S. Navy), or their designees, (collectively known as the "agencies"), for such activities.

So as to avoid any potential for violation of the Anti-Deficiency Act, Keydet agrees to provide within fifteen (15) days an initial payment of \$100,000 to the Department of the Interior for its costs, via electronic funds transfer, pursuant to instructions to be provided by the Department of the Interior. Additionally, Keydet agrees to provide within fifteen (15) days an initial payment of \$400,000 to the Department of Defense for its costs via electronic funds transfer, pursuant to instructions to be provided by the Department of Defense. Expenses incurred by the Department of Commerce will be reimbursed within 15 days of receipt of invoices. All requests for reimbursement for these activities should

The Agencies
April 22, 2004
Page 2

be provided, along with supporting documentation, to Jay Jones, Keydet Energy, at the above address.

The trustees and Keydet expect to negotiate and enter into a Cooperative Agreement for further specific cooperative assessment activities.

The costs of the cooperative assessment covered under this Agreement will be limited to the reasonable costs to implement the activities outlined in the attached "Exhibit A". The commitment contained in this letter will also cover all other costs incurred by the Agencies until five (5) days after Keydet provides the Agencies with written notice terminating the Funding Commitment, provided that Keydet's liability for such costs under this commitment shall not exceed \$1,000,000 without prior written agreement between Keydet and the Agencies.

Jay Jones
Incident Commander
Keydet Energy
Corporation

West Coast Joint Assessment Team (WCJAT). 2017. Recommendations for Conducting Cooperative Natural Resource Damage Assessment.

<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=200743&inline>

APPENDIX 11: Resource-Specific Work Plan Examples

CDFW-OSPR Ephemeral Data Collection Plans

Marine – <https://wildlife.ca.gov/OSPR/NRDA/Marine-EDCPs>

Inland – <https://wildlife.ca.gov/OSPR/NRDA/Inland-EDCPs>

Conceptual Model

- Ricker, R. 2014. Guidance for Developing Conceptual Models for Oil Spills. (PDF)

Amphibians

Arroyo Toad

- Survey Protocol for The Arroyo Toad.
https://rctlma.org/Portals/3/EPD/consultant/arroyo_toad_survey_protocol.pdf

California Red-legged Frog

- USFWS. 2005. Revised Guidance on Site Assessment and Field Surveys for the California Red-legged Frog. <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=83914&inline>

California Tiger Salamander

- USFWS. 2003. Interim Guidance on Site Assessment and field surveys for determining presence or a negative finding of the CA tiger Salamander.
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=83915&inline>

Foothill Yellow-legged Frog

- PGE. 2002. Standardized approach for habitat assessment and visual encounter surveys for the foothill yellow legged frog. <https://relicensing.pcwa.net/documents/Library/PCWA-L-270.pdf>

Aquatic Bioassessment

- California Rapid Assessment Method (CRAM) Resource and Document Website
<https://www.cramwetlands.org/documents#field+books+and+sops>
- California Water Quality Monitoring Council. 2013. California Rapid Assessment Method for Wetlands. **Bar-Built Estuarine Wetlands** Field Book ver. 6.1.
https://www.cramwetlands.org/sites/default/files/May2013_CRAM%20Field%20Book%20BBE_v61.pdf
- California Water Quality Monitoring Council. 2013. California Rapid Assessment Method for Wetlands. **Depressional Wetlands** Field Book ver. 6.1.
https://www.cramwetlands.org/sites/default/files/2013.03.19_CRAM_Fieldbook_Depressional_final_0.pdf
- California Water Quality Monitoring Council. 2020. California Rapid Assessment Method. **Episodic Riverine** Field Book ver. 6.2.
https://www.cramwetlands.org/sites/default/files/Episodic%20Riverine%20CRAM%20Field%20Book_v6.2.pdf
- California Water Quality Monitoring Council. 2013. California Rapid Assessment Method for Wetlands. **Perennial Estuarine Wetlands** Field Book ver. 6.1.
https://www.cramwetlands.org/sites/default/files/2013.03.19_CRAM%20Field%20Book%20Estuarine%206.1_0.pdf

- California Water Quality Monitoring Council. 2013. California Rapid Assessment Method for Wetlands. **Riverine Wetlands** Field Book ver. 6.1.
https://www.cramwetlands.org/sites/default/files/2013.03.19_CRAM%20Field%20Book%20Riverine%206.1_0.pdf
- California Water Quality Monitoring Council. 2017. California Rapid Assessment Method for Wetlands. **Slope Wetlands** Field Book ver. 6.2.
https://www.cramwetlands.org/sites/default/files/CRAM%20Slope%20Wetland%20Field%20Book%20v6.2_2018-09-05.pdf
- California Water Quality Monitoring Council. 2020. California Rapid Assessment Method. **Individual Vernal Pools** Field Book ver. 6.2.
<https://www.cramwetlands.org/sites/default/files/Individual%20Vernal%20Pools%20CRAM%206.2%20Field%20Book%202020.3.31.pdf>
- California Water Boards. 2016. Standard Operating Procedures (SOP) for the Collection of Field Data for Bioassessments of California Wadeable Streams: Benthic Macroinvertebrates, algae, and physical habitat.
https://www.waterboards.ca.gov/water_issues/programs/swamp/bioassessment/docs/01-combined-sop-final-v4-11mar2016.pdf
- California Water Boards. 2016. Supplemental Guidance for the SWAMP Bioassessment Field Protocol.
https://www.waterboards.ca.gov/water_issues/programs/swamp/bioassessment/docs/02-guidance-document-v311mar2016-update.pdf

Birds

Document links below are from - CDFW Resource and Document Website:

<https://wildlife.ca.gov/Conservation/Survey-Protocols>

Bald Eagle

- USFWS. 2004. Protocol for Evaluating Bald Eagle Habitat And Populations In California.
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=83707&inline>
- CDFW. 2017. Bald Eagle Breeding Survey Instructions.
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=83706&inline>

Burrowing Owl

- Burrowing Owl Survey Protocol and Mitigation Guidelines.
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=83842&inline>

Coastal CA Gnatcatcher

- Coastal California Gnatcatcher (*Polioptila californica californica*) Presence/Absence Survey Protocol. <https://www.fws.gov/sites/default/files/documents/survey-protocol-for-coastal-california-gnatcatcher.pdf>

Golden Eagle

- Driscoll, D.E. 2010. Protocol for golden eagle occupancy, reproduction, and prey population assessment. American Eagle Research Institute.
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=83955&inline>
- USFWS. 2010. Interim Golden Eagle Inventory and Monitoring Protocols (PDF).

Great Grey Owl

- USDA. 2000. Survey Protocol for The Great Gray Owl In The Sierra Nevada Of California.
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=83965&inline>

Land Birds

- USDA - Handbook of Field Methods for Monitoring Landbirds
https://www.fs.usda.gov/psw/publications/documents/psw_gtr144/psw_gtr144.pdf

Least Bell's Vireo

- DOI. 2001. Least Bells Vireo Survey Guidelines/.
<https://www.fws.gov/sites/default/files/documents/survey-protocol-for-least-bells-vireo.pdf>

Marbled Murrelets

- Pacific Seabird Group. 2003. Methods for Surveying Marbled Murrelets In Forests:
- A Revised Protocol For Land Management And Research
http://www.pacificseabirdgroup.org/publications/PSG_TechPub2_MAMU_ISP.pdf

Northern Goshawk

- USDA. 2006. Northern goshawk inventory and monitoring technical guide.
https://www.fs.usda.gov/rm/pubs_series/wo/wo_gtr071.pdf

Northern Spotted Owl

- USFWS. 2012. Protocol for Surveying Proposed Management Activities that may impact Northern Spotted Owls, <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=83977&inline>

Swainson's Hawk

- CA Energy Commission. 2010. Swainson's Hawk Survey Protocols, Impact Avoidance, and Minimization Measures for Renewable Energy Projects in the Antelope Valley of Los Angeles and Kern Counties, California <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=83990&inline>
- Swainson's Hawk Technical Advisory Committee. 2000. Recommended Timing and Methodology for Sainsons Hawk Nesting Surveys in California's Central Valley
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=83991&inline>

Tricolored Blackbird

- CDFW. 2015. Guidance for avoidance of impacts to Tricolored blackbird breeding colonies on Agricultural fields in 2015. <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=99310&inline>

Willow Flycatcher

- Bombay, H.L., Benson, T.M., Valentine, B.E., Stefani, R.A. 2003. Willow flycatcher survey protocol for CA. <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=84019&inline>

Dry Washes

- Pena, S.J., Mazor, R., and Olson, J.R. 2019. Bioassessment of Non-perennial Streams affected by oil and gas extraction (PDF).
- VIDEO: Bioassessment of Non-perennial Streams affected by oil and gas extraction.
<https://www.youtube.com/watch?v=6uB4s-H8OXc>

- Mazor, R.D., Olson, J., Robison, M., Caudillo, A., and Brown, J. 2019. Assessing the biological condition of dry ephemeral and intermittent streams.
http://ftp.sccwrp.org/pub/download/DOCUMENTS/TechnicalReports/1089_EphemeralStreams.pdf

Electrofishing

- Reynolds, J.B., and Holliman, F.M. Guidelines for Assessment and Reduction of Electrofishing-Induced Injuries in Trout and Salmon.
https://trainingcenter.fws.gov/courses/CSP/CSP2C01/resources/8-Electrofishing-Effects/2_Guidelines%20for%20Assessment%20and%20Reduction%20of%20EF%20Injury.pdf
- Temple, A.J., and Dean, J. Electrofishing Standardization Protocol.
https://trainingcenter.fws.gov/courses/CSP/CSP2C01/resources/1-Introduction/6_Electrofishing%20Standardization%20Protocol.pdf
- USEPA. 2019. National Rivers and Streams Assessment 2018/19. Field Operations Manual Non-wadeable ver 1.2. https://www.epa.gov/sites/default/files/2019-05/documents/nrsa_1819_fom_nonwadeable_version_1.2.pdf

Invertebrates

Document links below are from - CDFW Resource and Document Website:

<https://wildlife.ca.gov/Conservation/Survey-Protocols>

Branchiopods

- DOI. 2017. Survey Guidelines for Listed Large Branchiopods.
<https://www.fws.gov/sites/default/files/documents/survey-guidelines-for-large-branchiopods.pdf>

Mollusks

- USDA. 2008. Survey Protocol for Aquatic Mollusk Species: Preliminary Inventory & Presence/Absence Sampling. https://www.blm.gov/or/plans/surveyandmanage/files/10-mollusks_v3-1.pdf

Valley Elderberry Longhorn Beetle

- USFWS. 2017. Framework for Assessing Impacts to the Valley Elderberry Longhorn Beetle.
<https://www.fws.gov/sites/default/files/documents/survey-guidelines-for-valley-elderberry-longhorn-beetle.pdf>

Mammals

Document links below are from - CDFW Resource and Document Website:

<https://wildlife.ca.gov/Conservation/Survey-Protocols>

San Joaquin Kit Fox

- USFWS. 2011. Standardized Recommendations for Protection of The Endangered San Joaquin Kit Fox Prior To Or During Ground Disturbance.
<https://www.fws.gov/sites/default/files/documents/survey-protocols-for-the-san-joaquin-kit-fox.pdf>

Morro Bay Kangaroo Rat

- DOI. 1996. Survey Protocol for the Morro Bay Kangaroo Rat.
<https://www.fws.gov/sites/default/files/documents/survey-protocols-for-the-morro-bay-kangaroo-rat.pdf>

Reptiles

Document links below are from - CDFW Resource and Document Website:

<https://wildlife.ca.gov/Conservation/Survey-Protocols>

Barefoot banded gecko

- CDFW. 2011. Survey protocol for presence of or negative finding for the barefoot banded gecko (Coleonyx switaki). <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=39305&inline>

Blunt-nosed leopard lizard

- CDFW. 2019. Approved Survey Methodology for the Blunt-nosed leopard lizard.
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=174900&inline>

Desert Tortoise

- Preparing for any Action that may occur within the range of the Mojave Desert Tortoise.
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=174633&inline>

Western Pond Turtle

- USGS. 2006. USGS Western Pond Turtle trapping survey protocol for the Southcoast Ecoregion.
https://sdmmp.com/upload/SDMMP_Repository/0/q4x2pztbkns61wv9hy30rjc78fg5dm.pdf
- USGS. 2006. USGS Western Pond Turtle Visual Survey Protocol for the Southcoast Ecoregion.
https://sdmmp.com/upload/SDMMP_Repository/0/4fnpv18xm0sqtw29j7d3rz56bkychg.pdf

Vegetation

Document links below are from - CDFW Resource and Document Website:

<https://wildlife.ca.gov/Conservation/Survey-Protocols>

- CDFW. 2018. Protocols for Surveying and Evaluating Impacts to Special Status Native Plant Populations and Sensitive Natural Communities.
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=18959&inline>
- USFWS. 2000. Guidelines for Conducting and Reporting Botanical Inventories for Federally Listed, Proposed and Candidate Plants. https://cnps.org/wp-content/uploads/2019/10/Bot-Cert_US-Fish-and-Wildlife-Service-guidelines-botanical-inventories-LR.pdf

West Coast Joint Assessment Team (WCJAT)

This document contains information about the cooperative damage assessment process conducted by Trustees. It also includes sampling methods.

- WCJAT. 2017. Recommendations for Conducting Cooperative Natural Resource Damage Assessment. <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=200743&inline>

APPENDIX 12: Links to Oil Spill Job Aids and Data Sets on Baseline Resource Condition

Job Aids

- Oil Fact Sheets for Spill Responders –
<https://response.restoration.noaa.gov/resources/oil-fact-sheets-spill-responders>
- California Department of Fish and Wildlife, Office of Spill Prevention and Response Natural Resource Damage Assessment. –
<https://wildlife.ca.gov/OSPR/NRDA>
- Uncrewed Aircraft Systems Oil Spill Response Job Aid –
<https://response.restoration.noaa.gov/oil-and-chemical-spills/oil-spills/uncrewed-aircraft-systems-oil-spill-response-job-aid>
- Oil Spills in Marshes: Planning and Response Considerations:
<https://response.restoration.noaa.gov/oil-and-chemical-spills/oil-spills/resources/oil-spills-marshes.html>
- Shoreline Assessment Job Aid:
<https://response.restoration.noaa.gov/oil-and-chemical-spills/oil-spills/resources/shoreline-assessment-job-aid.html>
- Oil Spill Response and Assessment Guidelines for Sea Turtles:
<https://response.restoration.noaa.gov/oil-and-chemical-spills/oil-spills/oil-spill-response-and-assessment-guidelines-sea-turtles>
- Guidelines for Oil Spill Response and Natural Resource Damage Assessment: Sea Turtles:
<https://www.fisheries.noaa.gov/resource/document/guidelines-oil-spill-response-and-natural-resource-damage-assessment-sea-turtles>
- Guidelines for Assessing Exposure and Impacts of Oil Spills on Marine Mammals:
<https://response.restoration.noaa.gov/guidelines-assessing-exposure-and-impacts-oil-spills-marine-mammals>
- Oil Spill Response and Killer Whales:
<https://response.restoration.noaa.gov/oil-and-chemical-spills/oil-spills/resources/oil-spill-response-and-killer-whales.html>

Monitoring and Historical Data Sources

- CAEPA Central Coast Ambient Monitoring Program – http://rdc-omega.mlml.calstate.edu/ca/view_data.php?org_id=rb3&org_id=rb3
- CA State Water Resources Control Board – SWAMP (Surface Water Ambient Monitoring Program) State Mussel Watch Program –
https://www.waterboards.ca.gov/water_issues/programs/swamp/mussel_watch.html#hd
- CalCOFI (CA Cooperative Oceanic Fisheries Investigations) – Oceanographic and Marine based datasets. <https://calcofi.org/data/>
- CDFW's BIOS (Biogeographic Information and Observation System) – Data collected by CDFW and partner agencies. <https://wildlife.ca.gov/Data/BIOS>
- CEDEN (California Environmental Data Exchange Network) – Data collected from various CA state agencies. Data categories include water quality, toxicity, tissue, benthic and habitat. – http://www.ceden.org/find_data_page.shtml
- MARINe (Multi-Agency Rocky Intertidal Network) - Data on more than 200 rocky intertidal monitoring sites in CA. – <https://marine.ucsc.edu/index.html>
- NOAA's ERMA (Environmental Response Management Application – Southwest) – used to support NRDA process. – <https://erma.noaa.gov/southwest>
- SCCWRP – Southern California Coastal Water Research Project

- Bight Program (Marine monitoring) – Includes sediment toxicity, sediment chemistry, benthic infauna, and megabenthic invertebrate and demersal fish –
<https://www.sccwrp.org/about/research-areas/regional-monitoring/southern-california-bight-regional-monitoring-program/bight-program-data-portal/>
- SFEI (San Francisco Estuary Institute) – Regional data center for SF Bay-Delta. Manages water quality, tissue, wetlands, historical and spatial data. Various data tools available. –
<https://www.sfei.org/sfeidata.htm>
- SFEI (San Francisco Estuary Institute) – Contaminant Data Display & Download (CD3) –
<https://cd3.sfei.org/>

Other Useful Resources

- CDFW's California Natural Diversity Database (CNDDDB) - Inventory of status and locations of rare plants and animals. Public access – <https://wildlife.ca.gov/Data/CNDDDB/Maps-and-Data>
- CDFW Marine Species Portal – Basic species information. –
<https://marinespecies.wildlife.ca.gov/>
- CA Department of Water Resources – CA Data Exchange Center – Hydrologic and climate information – <https://cdec.water.ca.gov/>
- CA Department of Water Resources – Water Data Library Station Map – water quality data –
<https://wdl.water.ca.gov/waterdatalibrary/Map.aspx>
- CA State Water Resources Control Board - water quality control plans and state policies for water quality. Will contain acceptable water quality levels for known contaminants and dischargers. May also specify total maximum daily loads (TMDLs).
- EcoAtlas – Aggregates data from various sources (<https://www.ecoatlas.org/data/>). Mapping tool found here: <https://www.ecoatlas.org/regions/ecoregion/statewide>
- NOAA Tide Predictions - https://tidesandcurrents.noaa.gov/tide_predictions.html
- USGS Current Water Data for California – daily streamflow conditions for monitored waterbodies. – <https://waterdata.usgs.gov/ca/nwis/rt>

Regional Board Water Quality Control Plans (Basin Plans)

- Region 1 - North Coast Region –
https://www.waterboards.ca.gov/northcoast/water_issues/programs/basin_plan/
- Region 2 - San Francisco Bay Region –
https://www.waterboards.ca.gov/sanfranciscobay/basin_planning.html
- Region 3 - Central Coast Region –
https://www.waterboards.ca.gov/centralcoast/publications_forms/publications/basin_plan/
- Region 4 - Los Angeles Region –
https://www.waterboards.ca.gov/losangeles/water_issues/programs/basin_plan/basin_plan_documentation.html
- Region 5 - Central Valley Region –
https://www.waterboards.ca.gov/centralvalley/water_issues/basin_plans/
- Region 6 - Lahontan Region –
https://www.waterboards.ca.gov/lahontan/water_issues/programs/basin_plan/index.html
- Region 7 - Colorado River Basin Region –
https://www.waterboards.ca.gov/coloradoriver/water_issues/programs/basin_planning/
- Region 8 - Santa Ana Region –
https://www.waterboards.ca.gov/santaana/water_issues/programs/basin_plan/index.html

- Region 9 - San Diego Region –
https://www.waterboards.ca.gov/sandiego/water_issues/programs/basin_plan/index.html