


# Dune Protected Areas Network 1.0

*Creating a Blueprint for  
Restoration in the Guadalupe  
Nipomo Dunes Complex*

# 2018



## Monitoring Plan

Prepared by: The Land Conservancy of San Luis Obispo County



## Table of Contents

Chapter 1: Introduction .....	3
Conservation Strategy Overview .....	3
Vision and Goals .....	4
Dune Protected Areas Network.....	5
Chapter 2: Monitoring and Evaluation.....	6
Management Activity Monitoring.....	6
Monitoring to Inform Management .....	8
Baseline Monitoring .....	13
Adaptive Management .....	14
Monitoring Data Management and Reporting.....	15
Monitoring and Deliverable Schedule.....	15
Appendices .....	16

# Chapter 1: Introduction


## Conservation Strategy Overview

There is a regional effort underway to manage the entire Guadalupe Nipomo Dunes Complex through a partnership known as the Dunes Collaborative. This partnership is made up of federal, state, private, and non-profit organizations such as US Fish and Wildlife Service (USFWS), the Land Conservancy of San Luis Obispo County (LCSLO), Guadalupe Nipomo Dunes Center, California State Parks - Oceano Dunes State Vehicular Recreation Area, County of Santa Barbara, State of California Coastal Conservancy and California Department of Fish and Wildlife (CDFW).

The primary purpose of this project is to develop a comprehensive Conservation Strategy for the Guadalupe Nipomo Dunes Complex (GNDC). The Conservation Strategy is comprised of three overlapping components, a *Restoration Plan*, *Work Plan*, and *Monitoring Plan*. This document is the *Monitoring Plan* component of this Strategy.

The *Restoration Plan* outlines the concept and design of the Conservation Strategy and answers the question, what will be done? This provides the framework for the work to be accomplished. It is made up of a Vision for future conservation, Goals, and a Strategy to accomplish those Goals. These are fixed for the life of the Conservation Strategy. Also included in the Restoration Plan is a site assessment that identifies existing resources, the threats to those resources and identifies opportunities for conservation and restoration.

The *Work Plan* identifies how the Conservation Strategy is implemented. It answers the essential questions: How much effort will the Conservation Strategy take and what will it cost? The *Work Plan* includes Objectives, Actions and Methods to achieve those Objectives as well as cost estimates. These are time dependent and fluid. Work plans are meant to change over time based on adaptive management.



The *Monitoring Plan* measures progress towards achieving our Conservation Vision and informs subsequent actions. It is essential to knowing if your management actions are working or if you need to do something different. This can also be referred to as “adaptive management”.

## Vision and Goals

The RTF set forth a vision for future conservation of the Guadalupe Nipomo Dunes Complex:


*The Dunes Collaborative promotes connected and continuous coastal dune complexes, which support a diverse and healthy native ecosystem where plants and wildlife thrive and the dynamic nature of the dunes is preserved. These dunes will provide places of wonder for the local community, visitors, and future generations to explore and enjoy.*

In order to promote this vision, the RTF identified the following goals for effective design of a Conservation Strategy:



## Dune Protected Areas Network

The backbone of this Conservation Strategy is a network of high priority conservation areas which promote the conservations goals, called the “Dune Protected Areas Network”, or DPA Network. The DPA Network is based loosely on the “Green Infrastructure Network” concept (Figure 1) used in urban environments to protect natural habitats and pathways. It is an interconnected system of protected natural areas that conserve ecosystem functions while providing benefits for wildlife (Benedict, Edward, & McMahon, 2002). Each DPA consists of *core areas* and *hubs*, which are connected by *linkages*.



**Figure 1: Green Infrastructure Network. The Dune Protected Network is roughly based on the Green Infrastructure Network used to create wildlife pathways through urban areas.**

**Core areas** are the nucleus of the network and are chosen by their biological significance or pristine example of unique habitat. The core areas were first selected using conservation modeling software; a tool being used around the world to efficiently select unbiased areas for conservation. Core selection was further refined through consultation with the RTF, professional recommendations and available occurrence data of rare and listed species. These selected core areas are relatively undisturbed and have low invasive species intrusion.

**Hubs** buffer the core areas to offer additional protection against invasion and disturbance. These extensions of the core areas allow for less fragmentation of habitat types and offer continuous native cover. Hubs may contain multiple core areas, connecting them together as a unit.

**Linkages** are linear features connecting hubs together to facilitate wildlife movement, seed dispersal, and gene flow between core areas freely. Connectivity between hubs is essential for preservation of species in perpetuity. Connectivity was analyzed using Linkage Mapper software specifically designed to support regional wildlife habitat connectivity analyses (McRae & Kavanagh, 2011). The output of the software was modified to meet the needs of each DPA.

## Chapter 2: Monitoring and Evaluation

This monitoring plan is designed to evaluate each DPA's progress in meeting its restoration objectives as outlined in the GNDC Conservation Strategy. There are three types of monitoring applicable to the management of the DPA Network:

- **Management Activity Monitoring** – This is monitoring that tracks what types of Restoration Methods and Activities are happening where. This is meant to track the management itself and not the effects of management. Activity monitoring seeks to answer questions like, “Are projects being implemented as planned? “Are prescriptions being followed?” “Are targets being met?”
- **Monitoring to Inform Management** – This type of monitoring involves defining threshold values or expected responses, then surveying to measure the response or a closely related indicator. Comparing monitoring results with these expected values indicates whether you should initiate, intensify, or alter management actions. An example would be measuring percent cover of an invasive plant to evaluate management actions designed to reduce the cover to a certain threshold value, say 1-5% cover. This monitoring seeks to answer questions like, “Are management activities resulting in desired outcomes?”
- **Baseline Monitoring** – Essential to the DPA Network management philosophy is the need to maintain viable landscapes and reverse declining trends. To evaluate this, we identify a type of monitoring that evaluates baseline conditions and tracks changes through time. This monitoring seeks to answer questions like, “What are the project effects on ecological conditions?”

Each of these monitoring measures will be described in the monitoring reports following year 1 and year 3 of restoration work. Monitoring reports will be completed for each of the DPAs where restoration objectives are currently being addressed.

### Management Activity Monitoring

Management activities will be monitored using a reputable GPS-linked management software. Information will be collected each day work is completed. This data will provide information for reporting, assisting in adaptive management, and measuring success.

Products of activity monitoring include:

- photos from selected photo points (before and after)
- specific amount (acres) and location in which restoration has taken place
- amount of each chemical applied (if applicable)
- notes on restoration activity effectiveness

The preferred activity management software is AgTerra Technologies GIS data management platform. The AgTerra platform integrates mobile mapping, data collection and reporting solutions. Data is collected in the field each day using smartphones or tablets and then uploaded to a cloud-based server. Data is then easily exported into an ESRI ArcGIS format or GoogleEarth. This occurs at the end of each work day and is considered part of daily management activities. A screenshot of the project interface using AgTerra is given in Figure 2.




Figure 2: AgTerra project interface for management activity monitoring.



## Monitoring to Inform Management

To evaluate the success of restoration projects in the GNDC, contractual objectives were established which must be met to be deemed a success. These objectives contain the SMART criteria: Specific (who, what, where, when, and why); Measurable; Achievable; Results-oriented; and Time-fixed.

In most cases, these objectives focus on control or eradication of the invasive species to a certain threshold value (percent cover). To help measure if an invasive species target threshold is being achieved a protocol was developed by Wildlands Conservation Science for survey mapping in the GNDC capturing the most important information for management accurately and efficiently (Ball & Olthof, 2016). The protocol was established to allow for simple and uniform data sharing between land managers and Dunes Collaborative members. Protocol instructions are included in Appendix A.

The Conservation Strategy's work plan outlines the restoration objectives for each Dune Protected Area (DPA). Project contracts were executed for a selection of those objectives. Progress toward each selected objective will be included in the appropriate DPA monitoring report. The following section identifies specific objectives to be accomplished for each funded DPA; the protocol to follow for monitoring; and what actions will be taken if the Objectives are not met.

### **Black Lake Ecological Area (BLEA)**

***Objective 1: Reduce perennial veldtgrass cover to 1-5% cover class by year 3 in the BLEA DPA (Hub & Core).***

*Performance monitoring* will occur in Years 1 and 3 to document progress towards meeting the objective of perennial veldtgrass maintained at a 1-5% cover class value throughout the BLEA DPA by Year 3. This is considered the “knock down” Phase. Performance monitoring will then switch to long-term maintenance and monitoring with a monitoring interval of once every 5 years to ensure 1-5% cover class values are being maintained.

*Protocol:* Monitoring will follow the invasive plant mapping protocol used in the baseline Assessment (Appendix A).

*Actions if Objective is not met:* If monitoring shows that the Objective is not being met, the methods being used will be re-evaluated by a group of technical advisors (The Dune Collaborative Restoration Task Force or RTF) to determine why they are not working and if a change in methods is required. If the methods are found to be sound, then the RTF will determine if the Objective is a realistic target and if not, revise the Objective or terminate activities.

**Objective 2: Reduce invasive plant cover to 1-5% within a 25-foot buffer of Nipomo lupine populations by year 3. Maintain that infestation level during the maintenance phase in Yrs. 4-10.**

*Performance monitoring* will occur in Years 1 and 3 to document progress towards meeting the objective of *invasive plant cover* maintained at a 1-5% cover class value within the Nipomo lupine buffer zones by Year 3. This is considered the “knock down” Phase. Performance monitoring will then switch to long-term maintenance and monitoring with a monitoring interval of once every 5 years to ensure 1-5% cover class values are being maintained.

*Protocol:* Monitoring will follow the invasive plant mapping protocol used in the baseline Assessment (Appendix A).

*Actions if Objective is not met:* If monitoring shows that the Objective is not being met, the methods being used will be re-evaluated by a group of technical advisors (The Dune Collaborative Restoration Task Force or RTF) to determine why they are not working and if a change in methods is required. If the methods are found to be sound, then the RTF will determine if the Objective is a realistic target and if not, revise the Objective or terminate activities.

**Objective 3: Reduce European beachgrass cover to 1-5% cover class by year 3 in the BLEA DPA (Hub & Core).**

*Performance monitoring* will occur in Years 1 and 3 to document progress towards meeting the objective of European beachgrass maintained at a 1-5% cover class value throughout the BLEA DPA by Year 3. This is considered the “knock down” Phase. Performance monitoring will then switch to long term maintenance and monitoring with a monitoring interval of once every 5 years to ensure 1-5% cover class values are being maintained.

*Protocol:* Monitoring will follow the invasive plant mapping protocol used in the baseline Assessment (Appendix A).

*Actions if Objective is not met:* If monitoring shows that the Objective is not being met, the methods being used will be re-evaluated by a group of technical advisors (The Dune Collaborative Restoration Task Force or RTF) to determine why they are not working and if a change in methods is required. If the methods are found to be sound, then the RTF will determine if the Objective is a realistic target and if not, revise the Objective or terminate activities.

## **National Wildlife Refuge**

***Objective 1: Reduce European beachgrass cover to 1-5% cover class by year 3 in the NWR/CSD DPA (Hub & Core).***

*Performance monitoring* will occur in Years 1 and 3 to document progress towards meeting the objective of *European beachgrass* maintained at a 1-5% cover class value throughout the NWR/CSD DPA by Year 3. This is considered the “knock down” Phase. Performance monitoring will then switch to long term maintenance and monitoring with a monitoring interval of once every 5 years to ensure 1-5% cover class values are being maintained.

*Protocol:* Monitoring will follow the invasive plant mapping protocol used in the baseline Assessment (Appendix A)

*Actions if Objective is not met:* If monitoring shows that the Objective is not being met, the methods being used will be re-evaluated by a group of technical advisors (The Dune Collaborative Restoration Task Force or RTF) to determine why they are not working and if a change in methods is required. If the methods are found to be sound, then the RTF will determine if the Objective is a realistic target and if not, revise the Objective or terminate activities.

***Objective 2: Reduce priority invasive plant cover to 1-5% cover class by year three within a fenced buffer around Myrtle Pond.***

*Performance monitoring* will occur in Years 1 and 3 to document progress towards meeting the objective to reduce invasive plant cover at a 1-5% cover class value throughout the Myrtle Pond enclosure area by Year 3. This is considered the “knock down” Phase. Performance monitoring will then switch to long-term maintenance and monitoring with a monitoring interval of once every 5 years to ensure 1-5% cover class values are being maintained.

*Protocol:* Monitoring will follow the invasive plant mapping protocol used in the baseline Assessment (Appendix A). All rare plants documented during this assessment will also be recorded in the online database Calflora ([www.calflora.org](http://www.calflora.org)).

*Actions if Objective is not met:* If monitoring shows that the Objective is not being met, the methods being used will be re-evaluated by a group of technical advisors (The Dune Collaborative Restoration Task Force or RTF) to determine why they are not working and if a change in methods is required. If the methods are found to be sound, then the RTF will determine if the Objective is a realistic target and if not, revise the Objective or terminate activities.

**Objective 3: Reduce iceplant cover to 1-5% cover class by year 3 in selected area to expand on current efforts.**

Performance monitoring will occur in Years 1 and 3 to document progress towards meeting the objective of iceplant (*Carpobrotus* spp.) maintained at a 1-5% cover class value throughout the selected work area by Year 3. This is considered the “knock down” Phase. Performance monitoring will then switch to long-term maintenance and monitoring with a monitoring interval of once every 5 years to ensure 1-5% cover class values are being maintained.

*Protocol:* Monitoring will follow the invasive plant mapping protocol used in the baseline Assessment (Appendix A)

*Actions if Objective is not met:* If monitoring shows that the Objective is not being met, the methods being used will be re-evaluated by a group of technical advisors (The Dune Collaborative Restoration Task Force or RTF) to determine why they are not working and if a change in methods is required. If the methods are found to be sound, then the RTF will determine if the Objective is a realistic target and if not, revise the Objective or terminate activities.

**Objective 4: Reduce feral pig populations by 90% annually on the entire GNDNWR for 3 yrs. Assess removal interval after year five.**

*Performance monitoring* will occur during each removal event. The act of flying the GNDNWR will also provide an assessment of the pig population as a snap-shot in time. In Years 1 and 3 trends will be looked at to document progress towards meeting the objective of feral pig populations reduced by 90% annually

*Protocol:* During aerial removal flights, all pig locations and numbers will be documented using GPS/GIS technology. These numbers will be compared to the total number of pigs removed during the culling event.

*Actions if Objective is not met:* If monitoring shows that the Objective is not being met, the methods being used will be re-evaluated by a group of technical advisors (The Dune Collaborative Restoration Task Force or RTF) to determine why they are not working and if a change in methods is required. If the methods are found to be sound, then the RTF will determine if the Objective is a realistic target and if not, revise the Objective or terminate activities.

## **Point Sal**

### ***Objective 1: Eradicate jubata grass within the Point Sal Reserve DPA by year three.***

*Performance monitoring* will occur in Years 1 and 3 to document progress towards meeting the objective of jubata grass eradicated throughout the Point Sal Reserve DPA by Year 3. Performance monitoring will then switch to long-term early detection and rapid response monitoring once every 5-years to ensure eradication is maintained.

*Protocol:* Monitoring will follow the invasive plant mapping protocol used in the baseline Assessment (Appendix A)

*Actions if Objective is not met:* If monitoring shows that the Objective is not being met, the methods being used will be re-evaluated by a group of technical advisors (The Dune Collaborative Restoration Task Force or RTF) to determine why they are not working and if a change in methods is required. If the methods are found to be sound, then the RTF will determine if the Objective is a realistic target and if not, revise the Objective or terminate activities.

### ***Objective 2: Create a sustainable self-sustaining trail system in the Point Sal Reserve DPA by year five.***

*Performance monitoring* criteria will be determined as part of the initial Trails Assessment.

## Baseline Monitoring

To determine if the Conservation Strategy is achieving its higher-level goals, it is important to set up a monitoring program that will track changes over time. Ecosystems are dynamic, none more so than the coastal dune environment. There will be multiple successional trajectories that are possible, but tracking species composition and functional groups as they change through time will help us evaluate if our management actions are indeed keeping the dune ecosystem viable and sustainable through time.

The most efficient way to achieve this is by setting up and monitoring vegetation relevés. All relevé monitoring will follow California Native Plant Society standardized relevé protocols (Appendix B). This monitoring method allows for quick classification over a large area. It relies on ocular estimates of plant cover rather than counts of the “hits” of particular species along a transect line or precise measurements of cover/biomass by planimetric or weighing techniques. Monitoring will take place in years 1 and 3 to assess how native biodiversity is changing throughout management as well as assist in adaptive management.

Relevé plots should be selected based on both compositional and structural integrity. This means plots will be selected where species composition as well as horizontal and vertical spacing are similar throughout the plot. Vegetation alliances from The Manual of California Vegetation (MCV) is often used to delineate plot types. At least 2 plots per vegetation type (more if deemed necessary) will be selected for monitoring. Plot size and shape vary depending on vegetation cover (recommendations are given in Appendix B).

## Adaptive Management

Adaptive management is a process for evaluating how well the methods of a plan are meeting the stated objectives and using these evaluations to refine future methods and approaches of the plan. In real life, this happens in the field. On a day-to-day basis, land practitioners are evaluating the tools and techniques they are using and determining ways to increase productivity while meeting the desired goals. Decisions are made quickly and typically by those that are present. Adaptive management essentially happens without having to name it or formalize the process. However, there is also merit in having a formalized way of gathering information to have more formal processes to reflect on the success of a program and if changes in method, strategy or direction are warranted. This allows more time to work with experts to ensure management is based on the best available science and critical thought.

For the DPA Network Conservation Strategy, formal program evaluations will occur in year 1 and 3 coinciding with years that monitoring occurs. In those years, monitoring reports will be prepared to evaluate:

- What happened (Management Activity Monitoring),
- are we meeting our stated Objectives (Monitoring to Inform Management),
- and is our Conservation Strategy working (Baseline Monitoring).

Meetings will be held in those years to discuss the monitoring evaluations and refine our management methods and Conservation Strategy based on the findings. Changes will be incorporated into new workplans to guide management in the field. After Year three, the program will move into a long-term maintenance mode and these monitoring and evaluation events will occur on five year intervals in perpetuity.




Image credit: CEDA, dredging.org

## Monitoring Data Management and Reporting

Monitoring Reports will be submitted to the Dunes Collaborative concluding monitoring years and will include pertinent information about each of the monitoring methods, progress towards objectives, and recommendations for adaptive management. Deliverables are delineated in the monitoring reporting outline (Appendix C).

All monitoring data, occurrence updates, and filled data gaps will be housed with The Land Conservancy, on behalf of the Dunes Collaborative. All partners will contribute annually to the database to insure all available data is used in future evaluations and management decisions. The database will also be available for all partners to use and can be accessed by contacting The Land Conservancy.

The Land Conservancy of San Luis Obispo County  
 1137 Pacific Street Suite A  
 San Luis Obispo, CA 93401  
 info@lcslo.org

## Monitoring and Deliverable Schedule

	Year 1				Year 2			
	Fall 2018	Winter 2019	Spring 2019	Summer 2019	Fall 2019	Winter 2020	Spring 2020	Summer 2020
Implement Restoration								
Monitoring Events								
Prepare Report			Year 1 Report					
Adaptive Management Meeting/Review								

	Year 3				Year 4
	Fall 2020	Winter 2021	Spring 2021	Summer 2021	Fall 2021
Implement Restoration					
Monitoring Events					
Prepare Report			Final Report		
Adaptive Management Meeting/Review				Long Term Management Planning	



## Appendix A: Monitoring to Inform Management: Assessment Protocols

To evaluate the success of restoration projects in the GNDC, contractual objectives were established which must be met to be deemed a success. These objectives contain the SMART criteria: Specific (who, what, where, when, and why); Measurable; Achievable; Results-oriented; and Time-fixed.

In most cases, these objectives focus on control or eradication of the invasive species to a certain threshold value (percent cover). To help measure if an invasive species target threshold is being achieved a protocol was developed by Wildlands Conservation Science for survey mapping in the GNDC capturing the most important information for management accurately and efficiently (Ball & Olthof, 2016). The protocol was established to allow for simple and uniform data sharing between land managers and Dunes Collaborative members.

A target invasive plant species list was selected by the Dunes Collaborative and was informed by the *Invasive Plant Inventory and Early Detection Prioritization Tool* (Olsen & Hall, 2015). However, invasive plants aren't the only species causing widespread damage in the GNDC. Feral pig localities, numbers observed, and habitat damage are to be documented. In addition to invasive species, surveys will also target special status native plants such as Nipomo lupine (*Lupinus nipomensis*), La Graciosa thistle (*Cirsium scariosum* var. *loncholepis*), beach spectaclepod (*Dithyrea maritima*), and surf thistle (*Cirsium rhothophilum*) which are known to occur throughout the GNDC. A list of species targeted for inventory surveys is found in Table 1. Additional invasive and rare plant species can be included on the target species list, depending on the needs of the land managers.

When target species are encountered, their location, distribution and ground cover will be recorded using one of three mapping methods herein referred to as point, polygon, or grid. Point and polygon mapping is restricted to plant populations with a discernible boundary extent, these mapping units are herein referred to as populations or stands. An individual population is defined by a single contiguous infestation or a cluster of infestations separated by no more than 30-meters.

**Table 1: Species List for Survey Inventory**

Method	Species	Common Name	Family	Conservation Status	Cal-IPC Ranking
<b>Documented Invasive Plants</b> <i>Map Using a Grid System</i> <i>(5 Species)</i>	<i>Ammophila arenaria</i>	European beachgrass	Poaceae	<i>invasive plant</i>	High
	<i>Carpobrotus chilensis</i>	ice-plant / sea fig	Aizoaceae		Moderate
	<i>Carpobrotus edulis</i>	freeway ice-plant	Aizoaceae		High
	<i>Concosia pugioniformis</i>	slender leaf ice-plant	Aizoaceae		Limited
	<i>Ehrharta calycina</i>	perennial veldt grass	Poaceae		High
<b>Documented Invasive Plants</b> <i>Map Using Points &amp; Polygons</i> <i>(12 Species)</i>	<i>Arundo donax</i>	giant reed	Poaceae	<i>invasive plant</i>	High
	<i>Delairea odorata</i>	cape ivy	Asteraceae		High
	<i>Thinopyrum junceiforme</i>	russian wheatgrass	Poaceae		Red Alert
	<i>Tamarix sp.</i>	tamerisk	Tamaricaceae		High
	<i>Senecio elegans</i>	purple ragwort	Asteraceae		n/a
	<i>Brassica tournefortii</i>	saharan mustard	Brassicaceae		High
	<i>Hedera sp.</i>	algerian/english ivy	Araliaceae		High
	<i>Lepidium draba</i>	hoary cress	Brassicaceae		Moderate
	<i>Vinca major</i>	greater periwinkle	Apocynaceae		Moderate
	<i>Centaurea solstitialis</i>	yellow star thistle	Asteraceae		High
	<i>Cortaderia jubata</i>	pampas grass	Poaceae		High
	<i>Glebionis coronarium</i>	crowndaisy	Asteraceae		Moderate
<b>Dune Protected Areas Only (Grasses)</b> <i>Map Using a Grid System</i> <i>(4 Species)</i>	<i>Bromus madritensis ssp rubens</i>	red brome	Poaceae	<i>invasive plant</i>	High
	<i>Bromus tectorum</i>	downy brome	Poaceae		High
	<i>Cynodon dactylon</i>	bermudagrass	Poaceae		Moderate
	<i>Cenchrus clandestinus</i>	kikuyugrass	Poaceae		Limited
<b>Dune Protected Areas Only (Non-grasses)</b> <i>Map Using a Grid System</i> <i>(4 Species)</i>	<i>Cirsium vulgare</i>	bull thistle	Asteraceae	<i>invasive plant</i>	Moderate
	<i>Conium maculatum</i>	poison hemlock	Apiaceae		Moderate
	<i>Myoporum laetum</i>	ngaio tree	Myoporaceae		Moderate
	<i>Foeniculum vulgare</i>	sweet fennel	Apiaceae		High
<b>Early Detection Invasive Plants (Undocumented)</b> <i>Map Using Grid, Points or Polygons</i> <i>(9 Species)</i>	<i>Eichornia crassipes</i>	common water-hyacinth	Pontederiaceae	<i>invasive plant</i>	High
	<i>Alternanthera philoxeroides</i>	alligator weed	Amaranthaceae		High
	<i>Genista monspessulana</i>	french broom	Fabaceae		High
	<i>Lepidium latifolium</i>	perennial pepperweed	Brassicaceae		High
	<i>Limonium sp.</i>	Algerian sea lavender	Plumbaginaceae		Limited
	<i>Salvinia molesta</i>	giant salvinia	Salviniaceae		High-Alert
	<i>Taeniatherum caput-medusae</i>	medusahead	Poaceae		High
	<i>Hydrilla verticillata</i>	hydrilla	Hydrocharitaceae		High
	<i>Ludwigia sp.</i>	Uruguay waterprimrose	Onagraceae		High
<i>Emex spinosa</i>	Spiny emex	Polygonaceae	Moderate		
<b>Documented Special Status Native Plants</b> <i>Map Using Grid, Points or Polygons</i> <i>(6 Species)</i>	<i>Cirsium rhotophilum</i>	surf thistle	Asteraceae	CT; 1B.2 FE; CT; 1B.1	<i>na</i>
	<i>Cirsium scariosum var. loncholepis</i>	La Graciosa thistle	Asteraceae		
	<i>Dithyrea maritima</i>	beach spectaclepod	Brassicaceae		
	<i>Lupinus nipomoensis</i>	Nipomo Lupine	Fabaceae		
	<i>Nasturtium gambelii</i>	gambel's watercress	Brassicaceae		
	<i>Arenaria paludicola</i>	marsh sandwort	Caryophyllaceae		
<b>Undocumented Special Status Native Plants</b> <i>Map Using Grid, Points or Polygons</i> <i>(1 Species)</i>	<i>Layia carnosa</i>	Beach layia	Asteraceae	FE; CET; 1B.1	<i>na</i>
<b>Non-native Vertebrates</b> <i>Map Using Points &amp; Polygons</i> <i>(1 Species)</i>	<i>Sus scrofa</i>	Feral Pig	Suidae	<i>invasive animal</i>	<i>na</i>

**Total- 36 Invasive Plants, 5 Special Status Native Plants & 1 Invasive Animal**

Descriptions of the three mapping methodologies are provided below:

**Point** - Discrete populations with easily identifiable (circular) boundaries will be mapped using a single data point collected at the population centroid. For each population, diameter and percent ground cover and attribute information listed in Table 2 will be collected. Plant populations mapped as points will be later buffered by their infestation radius and converted to polygons for the final product. All feral pigs and rare plant occurrences will be mapped using discrete point data.

**Polygon** – Populations with a discernible, irregular-shaped boundary are mapped using a polygon drawn atop a high-resolution orthophotograph. Additional population attributes listed in Table 2 must be collected.

**Grid** - European beachgrass (*Ammophila arenaria*), perennial veldt grass (*Ehrharta calycina*), sea-fig iceplant (*Carpobrotus chilensis*), hottentot fig iceplant (*Carpobrotus edulis*) and Narrow-leaved iceplant (*Conicosia pugioniformis*) cannot be mapped using point or polygon methods because there are no discernible population boundaries to be delineated. Therefore, these widespread and/or diffusely occurring species will be mapped by estimating ground cover within the pre-established grid system. Within each grid cell, additional population attribute information is collected (Table 3).

These methods can be implemented either by aerial collection (100 meter<sup>2</sup> grids) or by ground collection (50 meter<sup>2</sup> grids). For aerial surveys, a 100-meter grid size was selected because it is a cost-effective scale for large property surveys while allowing for data resolution that is useful for weed population tracking and treatment planning. For surveys done on foot, a 50-meter grid size is more effective. The entire GNDC has a working 100-meter grid with nested 50-meter grid cells that should be used for mapping to ensure seamless integration. This grid is available from The Land Conservancy of San Luis Obispo County. A survey using both methods was recently used in the GNDC Rancho Guadalupe County Park and Point Sal Reserve (Roddick & Hall, 2018).

**Table 2: Attribute field information associated with polygon data to be recorded during the survey.**

Field Name	Attribute Description
Stand_ID	Individual stand identification code
Date	Date in which the survey was performed
Com_Name	Common name of the documented population stand
Species	Scientific name of documented population stand
Num_Indv	Estimated number of plants within documented population stand
Pop_Dens	The vegetative cover of the documented invasive species within the mapped population based off the CNPS cover class diagrams. The cover-classes are used to visually estimate cover within the polygon. Value ranges: <b>0-1%, 1-5%, 5-25%, 25-50%, 50-75%, 75-95%, 95-100%</b> .
Age_Class	The common age of plants within the population stand. Age is divided into <b>seedling, immature, mature</b> , mixed classes with more young plants ( <b>MixedYoung</b> ) and mixed age classes with more old plants than young ( <b>MixedOld</b> ).
ID_Confid	Confidence level ( <b>High, Med, Low</b> ) that the surveyor was able to identify the documented plant to species.
Photo	A photo taken of the population stand, if necessary
Surveyor	The name of the surveyor recording the data
Comment	Miscellaneous notes regarding the documented population stand
Gross_Acres	Total area (Acres) of the polygons including the interstitial spaces between the documented invasive plants within a populations (post-survey).
Net Acres	Net area (acres) covered by the documented invasive plants within the polygon, not including the interstitial spaces between plants. Calculated by multiplying the midpoint value of the pop_Dens x the Gross_Acres value (post-survey).
Rank	Plant ranking for the documented invasive species or rare plant (post-survey).
Point_X	X coordinate of the polygon centroid in NAD_1983_StatePlane_California_V_FIPF_0405_Feet
Point_Y	Y coordinate of the polygon centroid in NAD_1983_StatePlane_California_V_FIPF_0405_Feet

**Table 3: Attribute field information associated with grid data to be recorded during the survey.**

Field Name	Attribute Description
ID	Individual grid cell identification code
Date	Date in which the survey was performed
AMAR_Cover	The vegetative cover of European beachgrass within the grid cell based on the CNPS class cover diagrams. Value ranges: <b>0-1%, 1-5%, 5-25%, 25-50%, 50-75%, 75-95%, 95-100%</b> .
CACH_Cover	The vegetative cover of Hottentot fig iceplant within the grid cell based on the CNPS class cover diagrams. Value ranges: <b>0-1%, 1-5%, 5-25%, 25-50%, 50-75%, 75-95%, 95-100%</b> .
CAED_Cover	The vegetative cover of sea-fig icelant within the grid cell based on the CNPS class cover diagrams. Value ranges: <b>0-1%, 1-5%, 5-25%, 25-50%, 50-75%, 75-95%, 95-100%</b> .
COPU_Cover	The vegetative cover of slender-leaved iceplant within the grid cell based on the CNPS class cover diagrams. Value ranges: <b>0-1%, 1-5%, 5-25%, 25-50%, 50-75%, 75-95%, 95-100%</b> .
EHCA_Cover	The vegetative cover of perennial veldt grass within the grid cell based on the CNPS class cover diagrams. Value ranges: <b>0-1%, 1-5%, 5-25%, 25-50%, 50-75%, 75-95%, 95-100%</b> .
Gross_Acre	Total area (acres) of each mapped grid cell including the interstitial spaces between documented invasive species within a population (Post-survey).
AMAR_Acres	Net Area (acres) covered by European beachgrass within the grid cell, not including the interstitial spaces between plants. Calculated by multiplying the midpoint value of AMAR_Cover x the Gross_Acres (Post-survey).
CACH_Acres	Net Area (acres) covered by Hottentot fig iceplant within the grid cell, not including the interstitial spaces between plants. Calculated by multiplying the midpoint value of CACH_Cover x the Gross_Acres (Post-survey).
CAED_Acres	Net Area (acres) covered by sea-fig icelant within the grid cell, not including the interstitial spaces between plants. Calculated by multiplying the midpoint value of CAED_Cover x the Gross_Acres (Post-survey).
COPU_Acres	Net Area (acres) covered by slender-leaved iceplant within the grid cell, not including the interstitial spaces between plants. Calculated by multiplying the midpoint value of COPU_Cover x the Gross_Acres (Post-survey).
EHCA_Acres	Net Area (acres) covered by perennial veldt grass within the grid cell, not including the interstitial spaces between plants. Calculated by multiplying the midpoint value of EHCA_Cover x the Gross_Acres (Post-survey).

## Reporting

The findings of the assessment protocol will be included in the overall reporting established by the Monitoring Plan. The Monitoring Plan will include adaptive management decisions based on the findings of this assessment protocol.

Results from this portion of the monitoring plan will include visual and numerical information regarding the following:

- Invasive species
- Special Status Species
- Feral Pigs
- Habitats most affected by invasive species

### *Invasive Species*

Results regarding invasive species will include descriptive information about total acres covered during the survey and how many of the survey species were detected. It will include timing of the survey and any needed information about how the survey was implemented which might affect the outcome (flowering time, major obstacles, etc.). Most importantly, gross and net acres of the detected survey species (Grid: Example Table 4 and polygon/point: example Table 5) will be documented with associate maps indicating location (Figure 3).


**Table 4: Example table for grid survey invasive species reporting**

<b>Species</b>	<b>Gross Acres</b>	<b>Net Acres</b>
<i>Ehrharta calycina</i>	198.3	29.21
<i>Cakile maritima</i>	185.95	3.97
<i>Conicosia pugoniformis</i>	87.1	1.43
<i>Carpobortus edulis</i>	30.27	0.67
<i>Carpobortus chilensis</i>	75.37	1.94
<i>Ammophila arenaria</i>	0	0
<b>Total Acres</b>	<b>393.52*</b>	<b>37.22</b>

\*Multiple species were found within grid cells so this number is not a sum of each species' gross acres. Each grid cell was only counted once in the gross acres calculation.

**Table 5: Example table of polygon/point invasive species reporting**


Species	Approximate # Individuals	Gross Acres	Net Acres
Survey Species List			
Annual Grass	N/A	4.4819	1.2269
<i>Cirsium vulgare</i>	156	0.9083	0.0236
<i>Conium maculatum</i>	1125	1.3293	0.38
<i>Delairea odorata</i>	N/A	0.1338	0.0165
<i>Foeniculum vulgare</i>	13	0.0122	0.0002
<i>Tamarix sp</i>	2	0.001	0.0001
<i>Vinca major</i>	N/A	0.0241	0.0036
Additional Survey Species			
<i>Lathyrus latifolius</i>	N/A	0.2692	0.1009
<i>Tetragonia tetragonioides</i>	N/A	0.1063	0.0163
<i>Zantedeschia aethiopica</i>	1	0.001	0
<b>Total Acres</b>		<b>7.2671</b>	<b>1.7682</b>



**Figure 1: Example figure of Invasive species results.**

### *Special Status Species*


Results regarding documented special status plant species will include descriptive information about survey methods, how many of the survey species were detected, current condition, and population size. It will include timing of the survey and any needed information about how the survey was implemented which might affect the outcome (flowering time, major obstacles, etc.). Most importantly, acres or number of individuals, whichever is most appropriate, of the detected survey species will be documented with associate maps indicating location (Figure 4).



**Figure 2: Example figure for rare plant species survey results**

### *Feral Pigs*

Results regarding documented feral pigs will include descriptive information about survey methods, how many of the survey species were detected and/or evidence of their presence (vegetation damage and rooting), and estimated population size. It will include timing of the survey and any needed information about how the survey was implemented which might affect the outcome (seasonality, major obstacles, etc.). Most importantly, acres damaged, and number of individuals detected will be documented in addition to associated maps indicating location (Figure 5).



**Figure 3: Example figure for feral pig survey results**

*Habitats most affected by invasive species*

This portion of the results will focus on priorities for management. It will outline threatened habitats, especially those with high invasive species cover which are also inhabited by special status plant species. The major invasive species present in each habitat type will be documented in this section.



## References

- Ball, M., & Olthof, K. (2016). *Aerial Invasive Plant Survey and Treatment Prioritization Analysis*. Guadalupe Nipomo Dune Wildlife Refuge.
- Olsen, H., & Hall, J. (2015). *Invasive Plant Species and Area Prioritization Report: Guadalupe Nipomo Dunes Complex*.
- Roddick, L., & Hall, J. (2018). *Filling Major Data Gaps – Task 5 Implementation from Conservation Strategy/Management Plan Development for the Guadalupe-Nipomo Dunes Complex Project*. San Lu.

## Appendix B

CALIFORNIA NATIVE PLANT SOCIETY RELEVÉ PROTOCOL  
CNPS VEGETATION COMMITTEE  
October 20, 2000 (Revised 8/23/2007)

### Introduction

In *A Manual of California Vegetation* (Sawyer and Keeler-Wolf 1995), CNPS published a Vegetation Sampling Protocol that was developed as a simple quantitative sampling technique applicable to many vegetation types in California. Investigators use an ocular estimation technique called a relevé to classify and map large areas in a limited amount of time.

The relevé method of sampling vegetation was developed in Europe and was largely standardized by the Swiss ecologist Josias Braun-Blanquet. He helped classify much of Europe's vegetation, founded and directed a synecology center in France, and was editor of *Vegetatio* for many years. The relevé was, and is, a method used by many European ecologists, and others around the world. These ecologists refer to themselves as phytosociologists. The use of relevé in the United States has not been extensive with the exception of the US Forest Service.

The relevé is particularly useful when observers are trying to quickly classify the range of diversity of plant cover over large units of land. In general, it is faster than the point intercept technique. One would use this method when developing a classification that could be used to map of a large area of vegetation, for example. This method may also be more useful than the line intercept method when one is trying to validate the accuracy of mapping efforts.

The relevé is generally considered a “semiquantitative” method. It relies on ocular estimates of plant cover rather than on counts of the “hits” of a particular species along a transect line or on precise measurements of cover/biomass by planimetric or weighing techniques.

### Selecting a stand to sample:

A stand is the basic physical unit of vegetation in a landscape. It has no set size. Some vegetation stands are very small, such as alpine meadow or tundra types, and some may be several square kilometers in size, such as desert or forest types. A stand is defined by two main unifying characteristics:

- 1) It has compositional integrity. Throughout the site the combination of species is similar. The stand is differentiated from adjacent stands by a discernable boundary that may be abrupt or indistinct, and
- 2) It has structural integrity. It has a similar history or environmental setting that affords relatively similar horizontal and vertical spacing of plant species throughout. For example, a hillside forest originally dominated by the same species that burned on the upper part of the slopes, but not the lower, would be divided into two stands. Likewise, a sparse woodland occupying a slope with very shallow rocky soils would be considered a different stand from an adjacent slope with deeper, moister soil and a denser woodland or forest of the same species.

The structural and compositional features of a stand are often combined into a term called homogeneity. For an area of vegetated ground to meet the requirements of a stand it must be homogeneous.

Stands to be sampled may be selected by assessment prior to a site visit (e.g. delineated from aerial photos or satellite images), or may be selected on site (during reconnaissance to determine extent and boundaries, location of other similar stands, etc.). Depending on the project goals, you may want to select just one or a few representative stands for sampling (e.g., for developing a classification for a vegetation mapping project), or you may want to sample all of them (e.g., to define a rare vegetation type and/or compare site quality between the few remaining stands).

### **Selecting a plot to sample within in a stand:**

Because most stands are large, it is difficult to summarize the species composition, cover, and structure of an entire stand. We are also usually trying to capture the most information with the least amount of effort. Thus, we are typically forced to select a representative portion to sample.

When sampling a vegetation stand, the main point to remember is to select a sample that, in as many ways possible, is representative of that stand. This means that you are not randomly selecting a plot; on the contrary, you are actively using your own best judgement to find a representative example of the stand.

Selecting a plot requires that you see enough of the stand you are sampling to feel comfortable in choosing a representative plot location. Take a brief walk through the stand and look for variations in species composition and in stand structure. In many cases in hilly or mountainous terrain look for a vantage point from which you can get a representative view of the whole stand. Variations in vegetation that are repeated throughout the stand should be included in your plot. Once you assess the variation within the stand, attempt to find an area that captures the stand's common species composition and structural condition to sample.

### Plot Size

All relevés of the same type of vegetation to be analyzed in a study need to be the same size. It wouldn't be fair, for example, to compare a 100 m<sup>2</sup> plot with a 1000 m<sup>2</sup> plot as the difference in number of species may be due to the size of the plot, not a difference in the stands.

A minimal area to sample is defined by species/area relationships; as the sampler identifies species present in an area of homogeneous vegetation, the number will increase quickly as more area is surveyed. Plot shape and size are somewhat dependent on the type of vegetation under study. Therefore general guidelines for plot sizes of tree-, shrub-, and herb-dominated upland, and fine-scale herbaceous communities have been established. Sufficient work has been done in temperate vegetation to be confident the following conventions will capture species richness:

- Alpine meadow and montane wet meadow: 100 sq. m
- Herbaceous communities: 10 sq. m plot, 100 sq. m plot or 400 sq. m plot (Consult with CNPS, and use one consistent size)
- Shrublands: 400 sq. m plot
- Forest and woodland communities: 1000 sq. m plot
- Open desert vegetation: 1000 sq. m plot

## Plot Shape

A relevé has no fixed shape, plot shape should reflect the character of the stand. If the stand is about the same size as a relevé, you need to sample the entire stand. If we are sampling a desert wash, streamside riparian, or other linear community our plot dimensions should not go beyond the community's natural ecological boundaries. Thus, a relatively long, narrow plot capturing the vegetation within the stand, but not outside it would be appropriate. Species present along the edges of the plot that are clearly part of the adjacent stand should be excluded.

If we are sampling broad homogeneous stands, we would most likely choose a shape such as a circle (which has the advantage of the edges being equidistant to the center point) or a square (which can be quickly laid out using perpendicular tapes). If we are trying to capture a minor bit of variety in the understory of a forest, for example a bracken fern patch within a ponderosa pine stand, we would want both bracken and non-bracken understory. Thus, a rectangular shape would be appropriate.

## **GENERAL PLOT INFORMATION**

The following items appear on each data sheet and are to be collected for all plots. Where indicated, refer to attached code sheet.

**Polygon or Relevé number:** Assigned either in the field or in the office prior to sampling.

**Date:** Date of sampling.

**County:** County in which located.

**USGS Quad:** The name of the USGS map the relevé is located on; note series (15' or 7.5').

**CNPS Chapter:** CNPS chapter, or other organization or agency if source is other than CNPS chapter.

**Landowner:** Name of landowner or agency acronym if known. Otherwise, list as private.

**Contact Person:** Name, address, and phone number of individual responsible for data collection.

**Observers:** Names of individuals assisting. Circle name of recorder.

**Plot shape:** indicate the sample shape as: square, rectangle, circle, or the entire stand.

**Plot size:** length of rectangle edges, circle radius, or size of entire stand. NOTE: See page 2 for standard plot sizes.

**Study Plot Revisit:** If the relevé plot is being revisited for repeated sampling, please circle "Yes".

**Photo interpreter community code:** If the sample is in area for which delineation and photo interpretation has already been done, the code which the photointerpreters applied to the polygon. If the sample site has not been photointerpreted, leave blank.

**Other polygons of same type** (yes or no, if applicable), if yes, mark on map: Other areas within view that appear to have similar vegetation composition. Again, this is most relevant to areas that have been delineated as polygons on aerial photographs as part of a vegetation-mapping project. If one is not working from aerial photographs, draw the areas as on a topographic map.

**Is plot representative of whole polygon?** (yes or no, if applicable), if no explain: Detail what other vegetation types occur in the polygon, and what the dominant vegetation type is if there is more than one type.

**Global Positioning System Readings:** Due to the recent availability of very accurate and relatively low cost GPS units, we highly recommend obtaining and using these as a standard piece of sampling equipment. Now that the military intentional imprecision (known as “selective availability”) has been “turned off” (as of July 2000), it is typical for all commercial GPS units these units to be accurate to within 5 m of the actual location. Also note that the GPS units can be set to read in UTM or Latitude and Longitude coordinates and can be easily translated. Thus, the following fields for Latitude, Longitude, and legal description are now optional. In order for all positional data to be comparable within the CNPS vegetation dataset, we request using UTM coordinates set for the NAD 83 projection (see your GPS users manual for instructions for setting coordinates and projections).

**Caveat:** Although GPS units are valuable tools, they may not function properly due to the occasionally poor alignment of satellites or due to the complexity of certain types of terrain, or vegetation. We thus also recommend that you carry topographic maps and are aware of how to note your position on them in the event of a non-responsive or inaccurate GPS.

**UTMN and UTME:** Northing and easting coordinates using the Universal Transverse Mercator (UTM) grid as delineated on the USGS topographic map, or using a Global Positioning System.

**UTM zone:** Universal Transverse Mercator zone. Zone 10S for California west of the 120<sup>th</sup> longitude; zone 11S for California east of 120<sup>th</sup> longitude.

**Legal Description:** Township/Range/Section/Quarter Section/Quarter-Quarter section/Meridian: Legal map location of the site; this is useful for determining ownership of the property. California Meridians are Humboldt, Mt. Diablo, or San Bernardino. (This is optional, see above discussion of GPS units)

**Latitude and Longitude:** Degrees north latitude and east longitude. This is optional (see above)

**Elevation:** Recorded in feet or meters. Please indicate units.

**Slope:** Degrees, read from clinometer or compass, or estimated; averaged over relevé

**Aspect:** Degrees from true north (adjust declination), read from a compass or estimated; averaged over relevé.

**Macrotopography:** Characterize the large-scale topographic position of the relevé. This is the general position of the sample along major topographic features of the area. *See attached code list.*

**Microtopography:** Characterize the local relief of the relevé. Choose the shape that mimics the lay of the ground along minor topographic features of the area actually within the sample. *See attached code list.*

## VEGETATION DESCRIPTION

**Dominant layer:** Indicate whether the community is dominated by the Low layer (L), Mid-layer (M), or Tall (T) layer.

**Preliminary Alliance name:** Name of series, stand, or habitat according to CNPS classification (per Sawyer and Keeler-Wolf 1995); if the type is not defined by the CNPS classification, note this in the space.

**Dominant Vegetation Group:** Use code list to choose group

**Phenology:** Based on the vegetative condition of the principal species, characterize the phenology of each layer as early (E), peak (P), or late (L).

## WETLAND COMMUNITY TYPES

**Community type:** Indicate if the sample is in a wetland or an upland; note that a site need not be officially delineated as a wetland to qualify as such in this context.

**Dominant vegetation form:** This is a four letter code which relates the vegetation of the plot to the higher levels of the NBS/NPS National Vegetation Classification System hierarchy. *See attached code list.*

**Cowardin class:** See “Artificial Keys to Cowardin Systems and Names” (attached). If the plot is located in a wetland, record the proper Cowardin system name. Systems are described in detail in Cowardin et al. 1979. Classification of wetlands and deepwater habitats of the United States. US Dept. of the Interior, Fish and Wildlife Service, Office of Biological Services, Washington, D.C.

**Marine:** habitats exposed to the waves and currents of the open ocean (subtidal and intertidal habitats).

**Estuarine:** includes deepwater tidal habitats and adjacent tidal wetlands that are usually semi-enclosed by land but have open, partly obstructed, or sporadic access to the open ocean, and in which ocean water is at least occasionally diluted by freshwater runoff from the land (i.e. estuaries and lagoons).

**Riverine:** includes all wetlands and deepwater habitats contained within a channel, excluding any wetland dominated by trees, shrubs, persistent emergent plants, emergent mosses, or lichens. Channels that contain oceanic-derived salts greater than 0.5% are also excluded.

**Lacustrine:** Includes wetlands and deepwater habitats with all of the following characteristics: 1) situated in a topographic depression or a dammed river channel; 2) lacking trees or shrubs, persistent emergents, emergent mosses or lichens with greater than 30% aerial coverage; and total area exceeds 8 ha (20 acres). Similar areas less than 8 ha are included in the lacustrine system if an active wave-formed or bedrock shoreline feature makes up all or part of the low tide boundary, or if the water in the deepest part of the basin exceeds 2 m (6.6 feet) at low tide. Oceanic derived salinity is always less than 0.5%.

**Palustrine:** Includes all nontidal wetlands dominated by trees, shrubs, persistent emergents, emergent mosses or lichens, and all such wetlands that occur in tidal areas where salinity derived from oceanic salts is less than 0.5%. Also included are areas lacking vegetation, but with all of the following four characteristics: 1) areas less than 8 ha (20 acres); active wave-formed or bedrock shoreline features lacking; 3) water depth in the deepest part of the basin less than 2 m (6.6 feet) at low water; and 4) salinity due to ocean-derived salts less than 0.5%.

Vertical distance from high water mark of active stream channel: If the plot is in or near a wetland community, record to the nearest meter or foot the estimated vertical distance from the middle of the plot to the average water line of the channel, basin, or other body of water.

Horizontal distance from high water mark of active stream channel: If the plot is in or near a wetland community, record to the nearest meter or foot the estimated horizontal distance from the middle of the plot to the average water line of the channel, basin, or other body of water.

Stream channel form: If the plot is located in or near a community along a stream, river, or dry wash, record the channel form of the waterway. The channel form is considered S (single channeled) if it consists of predominately a single primary channel, M (meandering) if it is a meandering channel, and B (braided) if it consists of multiple channels interwoven or braided.

Adjacent alliance: Adjacent vegetation series, stands or habitats according to CNPS classification; list in order of most extensive to least extensive. Give the name of the alliance, the direction in relation to stand and list up to four species under Description.

**Photographs:** Write the name or initials of the camera owner and the JPEG numbers for photos taken. Write the camera's view direction from compass bearings. Take four or eight photos (depending on the project) from the same point as the GPS reading (center of a circle or NW corner of rectangle). Using a compass, take the first photo from the north, and rotate clockwise, taking the photos in sequence, N, NE, E etc, or N, E, S, W. Keep camera at same orientation, zoom level, and distance from ground for all four (or eight) photos., You may take photos close to the ground, if for instance, you are photographing a low herbaceous stand. Additional photos of the stand may also be helpful. If using a digital camera or scanning in the image into a computer, relevé numbers and compass directions can be recorded digitally. If using a 35mm camera, please note the roll number, frame number, compass direction, and the initials of the person whose camera is being used. (e.g. Roll 5, #1, to the NW, SE)

## STAND AND ENVIRONMENTAL INFORMATION

Vegetation trend: Based on the regenerating species and relationship to surrounding vegetation, characterize the stand as either increasing (expanding), stable, decreasing, fluctuating, or unknown.

Impacts: Enter codes for potential or existing impacts on the stability of the plant community. Characterize each as either 1. Light, 2. Moderate, or 3. Heavy. *See attached code list.*

Site location and plot description: A concise, but careful description that makes locating and/or revisiting the vegetation stand and plots possible; give landmarks and directions. Used in conjunction with the GPS position recorded earlier, this should enable precise re-location of the plot. Indicate where the GPS reading was taken within the plot. In general, the location of the GPS reading should be on the Northwestern corner of the plot, if the plot is rectangular (or square), or in the center if the plot is circular. It is also helpful to briefly describe the topography, aspect, and vegetation structure of the site. If you can't take the GPS reading at the Northwest corner (an obstacle in the way) then note where the GPS point was taken. If you can't get a GPS reading, then spend extra time marking the plot location as precise as possible on a topo map.

Site history: Briefly describe the history of the stand, including type and year of disturbance (e.g. fire, landslides or avalanching, drought, flood, or pest outbreak). Also note the nature and extent of land use such as grazing, timber harvest, or mining.

Unknown plant specimens: List the numbers of any unknown plant specimens, noting any information such as family or genus (if known), important characters, and whether or not there is adequate material for identification. Do not take samples of plants of which there are only a few individuals or which you think may be rare. Document these plants with photographs.

Additional comments: Feel free to note any additional observations of the site, or deviations from the standard sampling protocol. If additional data were recorded, e.g. if tree diameters were measured, please indicate so here.

## SURFACE COVER AND SOIL INFORMATION

Surface cover: Estimate the cover class of each size at or near the ground surface averaged over the plot. Always remember to estimate what you actually see on the surface as opposed to what you think is hiding under, organic litter, big rocks, etc. However, rocks, organic litter, or fine material visible under the canopy of shrubs or trees should be included in the cover estimate.

One way to consider this is to assume that all of the components of surface cover plus the basal cross-section of living plant stems and trunks (at ground level) will add up to 100%. Thus, estimate the cover value of each of the items in the box on the form for surface cover (including the basal area of plant stems) so that they will add up to 100%. Remember that the basal area of plant stems is usually minimal (e.g., if there were 10 trees, each 1 m in diameter at ground level on a 1000 square meter plot, they would cover less than 1% {0.79% } of the plot).



These data are asked for because certain categories of surface cover of rock and other materials have been shown to correlate with certain vegetation types and are thus likely influencing the type of vegetation that is growing in a given area. These estimates should be made quickly with the main point to keep in mind being a rough estimate of the relative proportions of different coarse fragments on the plot.

Fines: Fine mineral fragments including sand, silt, soil, “dirt” < 2 mm in diameter

Gravel: rounded and angular fragments 0.2-7.5 cm (0.08 -3 in.) diameter

Cobble: rounded and angular fragments >7.5-25 cm (3 -10 in.) in diameter

Stone: rounded and angular coarse fragments >25 cm-60 cm (10 -24 in.) in diameter

Boulder: rounded and angular coarse fragments >60 cm (>24 in.) in diameter

Bedrock: continuous, exposed, non-transported rock

Litter: extent of undecomposed litter on surface of plot (this includes all organic matter, e.g. fallen logs, branches, and twigs down to needles and leaves).

Living stems of vascular plants: basal area of living stems of the plants at ground surface

% Bioturbation: Estimate percent cover of ground disturbance by animals (e.g., small mammal burrowing trails, cow hoof marks) across the entire plot surface.

Soil texture: Record the texture of the upper soil horizon, below the organic layer if one is present. *See attached key and code list.*

Parent Material: Geological parent material of site. *See attached code list.*

## **VEGETATION DATA**

### **Assessment of Layers**

Data are recorded for five layers (tree overstory, tree understory, shrub, herb, and non-vascular). The layer a species occupies is determined by life-form. The estimates need not be overly precise and will vary among vegetation types. A young tree, if shrub sized, is considered an understory tree. A caveat: if several relevés are being sampled within the same vegetation type, it is important to be consistent when assigning layers. Some types will have more than five layers (e.g. two tree layers of different maximum height); this should be indicated in the relevé description.

### **Species List**

The collection of vegetation data continues with making a comprehensive species list of all vascular plants within the relevé. This list is achieved by meandering through the plot to see all

microhabitats. During list development, observers document each taxon present in each layer in which it occurs separately, recording it on a different line of the data form and noting which layer is represented. This is important for data entry because each layer of each represented taxon will be entered separately. Each individual plant is recorded in only one layer, the layer in which the tallest portion of the individual is found. One should reach a point at which new taxa are added to the list only very slowly, or sporadically. When one has reached that point, the list is probably done.

The following sections explain how to perform the actual relevé, the Estimation of Cover Values. The sections prefaced by bold-faced titles explain the technique, and the sections with regular font titles refer to the steps needed to complete the accompanying Field Form.

**DBH** – see separate field form (optional)

DBH if >10 cm:

The diameter at breast height (dbh) is important in certain studies. It may be recorded next to each tree species name. First indicate the species name by code and then record the number of sprouts/trunks in clonal trees. You should measure the tree dbh of every tree trunk/sprout that has diameter  $>$  or  $=$  10 cm at breast height in the plot, and each measurement should be in centimeters (cm) using a dbh tape measure. For trunks that may be fused below breast height and branched at breast height, each trunk at breast height gets a separate measurement.

Also indicate if each tree/clone is in the overstory or understory. Trees in the overstory are generally at canopy level. Trees in the understory are entirely below the general level of the canopy.

If snags are encountered in plot, record the dbh and denote it as dead by circling its dbh measurement. If you are unable to identify the snag to species, put the four letter code “SNAG” in the species column.

Depending on the density of trees in each plot, you can record dbh of trees for every tree trunk in the plot, or you can sub-sample the trunks to estimate dbh for every tree species in relatively dense plots. For woodland/forest plots, sub-sampling is appropriate for half the plot if there are at least 50 trees/resprouts present (e.g., 200 m<sup>2</sup> sub-sample in riparian and 500 m<sup>2</sup> sub-sample in upland).

When sub-sampling, make sure to denote this as a sub-sample (note on the data form) and record the sub-sample of dbh's for each tree species in the appropriate row on the Field Form. Once the data are post-processed and entered into a database, then you will need to record each sub-sampled dbh reading three additional times to come up with a full sample of dbh readings. For example, with a sub-sampled tree dbh of 15 cm, this value of 15 should be entered four times (not just once) when it is entered in the database.

Lifeform and size class: If dbh  $<$ 15.2 cm, counts should be made for conifers and hardwoods in two different size classes. Count seedlings ( $\leq$  2.54 cm) and saplings ( $>$  2.54 but  $<$  15.2 cm). First estimate if there are more than 50 seedlings in one half (50% subsample) of the plot. If so, then do counts of seedlings and saplings in five sub-plots of 2x2 m squares. If the plot shape is a

circle, place one square in the center of the plot, and four other squares 10 m to the N, S, E, and W of the plot center. If there are less than 50 seedlings in the 50% subsample plot, then record counts for that subsample instead.

### **Estimating Cover:**

There are many ways to estimate cover. Many people who have been in the cover estimation “business” for a long time can do so quickly and confidently without any props and devices. However, to a novice, it may seem incomprehensible and foolhardy to stand in a meadow of 50 different species of plants and systematically be able to list by cover value each one without actually “measuring” them in some way.

Of course, our minds make thousands of estimates of various types every week. We trust that estimating plant cover can be done by anyone with an open mind and an “eye for nature.” It’s just another technique to learn.

It is very helpful to work initially with other people who know and are learning the technique. In such a group setting, typically a set of justifications for each person’s estimate is made and a “meeting of the minds” is reached. This consensus approach and the concomitant calibration of each person’s internal scales is a very important part of the training for any cover estimate project.

An underlying point to remember is that estimates must provide some level of reliable values that are within acceptable bounds of accuracy. If we require an accuracy level that is beyond the realm of possibility, we will soon reject the method for one more quantitative and repeatable. As with any scientific measurement, the requirement for accuracy in the vegetation data is closely related to the accuracy of the information needed to provide a useful summary of it. Put into more immediate perspective - **to allow useful and repeatable analysis of vegetation data, one does not need to estimate down to the exact percent value the cover of a given plant species in a given stand.**

This point relates to two facts: there is inherent variability of species cover in any environment. For example, you would not expect to always have 23% *Pinus ponderosa*, 14% *Calocedrus decurrens*, and 11% *Pinus lambertiana* over an understory of 40% *Chamaebatia foliosa*, 3% *Clarkia unguiculata*, and 5% *Galium bolanderi* to define the Ponderosa pine-Incense cedar/mountain misery/bolander bedstraw plant community. Anyone who has looked at plant composition with a discerning eye can see that plants don’t space themselves in an environment by such precise rules. Thus, we can safely estimate the representation of species in a stand by relatively broad cover classes (such as <1%, 1-5 %, 5-25%, etc.) rather than precise percentages.

The data analysis we commonly use to classify vegetation into different associations and series (TWINSPAN and various cluster analysis programs, for example) is likewise forgiving. When analyzed by quantitative multivariate statistics information on species cover responds to coarse differences in cover and presence and absence of species, but not to subtle percentage point differences. This has been proven time and again through quantitative analysis of vegetation classification. Many of the world’s plant ecologists estimate cover rather than measure it precisely. Some of the seminal works in vegetation ecology have been based on cover estimates taken by discerning eyes.

With this as a preamble, below we offer some suggestions on estimating cover that have proven helpful. These are simply “tricks” to facilitate estimation, some work better for different situations. You may come up with other methods of estimation that may seem more intuitive, and are equally reliable in certain settings. All values on the relevé protocol that require a cover class estimate, including coarse fragment and vegetation layer information, may rely on these techniques. Just make the appropriate substitutions (using the coarse fragment example substitute, bedrock, stone, cobbles, gravel, and litter for vegetation).

#### Method 1: The invisible point-intercept transect:

This method works well in relatively low, open vegetation types such as grasslands and scrubs where you can see over the major stand components. For those who have worked with the original CNPS line intercept methodology it’s like counting hits along an imaginary line at regular intervals of the 50 m tape. Here’s how it goes:

Envision an imaginary transect line starting from your vantage point and running for 50 m (or however many meters you wish, as long as you are still ending up within the same stand of vegetation you’re sampling - never keep counting outside of your homogeneous stand). Now “walk” your eye along this tape for 50 m and visually “take a point” every 0.5 m. Don’t worry about precision, just try to “walk” your eye along the line and stop every 0.5 m or at any other regular interval until you reach its end and mentally tally what species you hit. Once you come up with a number of hits for each major species in one imaginary transect, take another transect in another direction and estimate the number of hits on that one. Do this several times (usually 3-4 is enough if you are in a homogeneous stand), then average your results.

This can go quickly in simple environments and in environments where the major species are easily discernable (chaparral, bunch-grassland, coastal scrub, desert scrub). Your average number of hits need not be a total of 100 as in the original transect method, but could be 50 along a 25 m imaginary line (in which case you would multiply by two to get your estimated cover), or 25 along a 12.5 m line (multiply average by 4), etc.

#### Method 2: Subdivision of sample plot into quadrants:

Many plots, whether they are square, circular, or rectangular, may be “quartered” and have each quadrant’s plant cover estimated separately. If the plot is a given even number of square meters (such as 100, 400, or 1000 m<sup>2</sup>) then you know that a quarter of that amount is also an easily measurable number. If you can estimate the average size of the plants in each of the quarters (e.g, small pinyon pines may be 5 m<sup>2</sup> (2.2m x 2.2m), creosote bush may be 2m<sup>2</sup> (or 1.41 m x 1.41 m), burrobush may be 0.5m<sup>2</sup>) then you simply count the number of plants in each size class and multiply by their estimated size for the cover in a given quadrant. Then you average the 4 quadrants together for your average cover value.

This method works well in vegetation with open-to-dense cover of low species such as grasses or low shrubs, in open woodlands, and desert scrubs.

#### Method 3; “Squash” all plants into a continuous cover in one corner of the plot :

Another way to estimate how much of the plot is covered by a particular species is to mentally group (or “march”, or “squash”) all members of that species into a corner of the plot and estimate the area they cover. Then calculate that area as a percentage of the total plot area. This technique works well in herb and shrub dominated plots but is not very useful in areas with trees.

Method 4: How to estimate tree cover:

Cover estimates of tall trees is one of the most difficult tasks for a beginning relevé sampler. However it is possible to do this with consistency and reliability using the following guidelines.

1. Have regular sized and shaped plots that you can easily subdivide.
2. Estimate average crown spread of each tree species separately by pacing the crown diameter of representative examples of trees of each species and then roughly calculating the crown area of each representative species.
3. Add together the estimated crown area of each individual of each species of tree on the plot for your total cover.

Method 5: The process of elimination technique:

This method is generally good for estimating cover on sparsely vegetated areas where bare ground, rocks, or cobbles cover more area than vegetation. In such a situation it would be advisable to first estimate how much of the ground is not covered by plants and then subdivide the portion that is covered by plants into rough percentages proportional to the different plant species present. For example, in a desert scrub the total plot not covered by plants may be estimated at 80%. Of the 20% covered by plants, half is desert sunflower (10% cover), a quarter is California buckwheat (5% cover), an eighth brittlebush (2.5% cover), and the rest divided up between 10 species of herbs and small shrubs (all less than 1% cover).

Any of these techniques may be used in combination with one another for a system of checks and balances, or in stands that have characteristics lending themselves for a different technique for each layer of vegetation.

In a relevé, cover estimates, using the techniques described above, are made for each taxon as it is recorded on the species list. Estimates are made for each layer in which the taxon was recorded. For example, if individuals of coast live oak occur in the tree overstory (canopy trees) and tree understory (seedlings and saplings), an estimate is made for both layers should be recorded.

In a traditional relevé, cover is estimated in “cover classes,” not percentages, because of the variability of plant populations over time and from one point to another, even within a small stand. This protocol uses the following 6 cover classes:

Cover Class 1: the taxon in that layer covers < 1 % of the plot area

Cover Class 2: the taxon in that layer covers 1 % - 5 % of the plot area

Cover Class 3a: the taxon in that layer covers >5 - 15 % of the plot area

Cover Class 3b: the taxon in that layer covers >15 - 25 % of the plot area  
Cover Class 4: the taxon in that layer covers >25 - 50 % of the plot area  
Cover Class 5: the taxon in that layer covers >50 - 75 % of the plot area  
Cover Class 6: the taxon in that layer covers > 75% of the plot area

### **Percentages (optional)**

This CNPS protocol also encourages observers to estimate percentages if they feel confident in their estimation abilities. This optional step allows the data to be compared more easily to data collected using different methods, such as a line or point intercept. It also instills confidence in the cover estimate of borderline species that are close calls between two cover classes (e.g., a cover class 2 at 5% as opposed to a cover class 3 at 6%). It is particularly useful for calculating cover by the process of elimination techniques and for estimating total vegetation cover (see below) and coarse fragment cover.

### **Overall Cover of Vegetation**

In addition to cover of individual taxa described above, total cover is also estimated for each vegetation layer. This is done using the same cover classes as described above but combines all taxa of a given category. They can be calculated from the species percent cover estimates, but please make sure to disregard overlap of species within each layer. These estimates should be absolute aerial cover, or the “bird’s eye view” of the vegetation cover, in which each category cannot be over 100%.

To come up with a specific number estimate for percent cover, first use the cover intervals, used in the species cover estimates, as a reference aid to get a generalized cover estimate: While keeping these intervals in mind, you can then refine your estimate to a specific percentage for each category below.

**% Overstory Conifer/Hardwood Tree:** The total aerial cover (canopy closure) of all live tree species that are specifically in the overstory or are emerging, disregarding overlap of individual trees. Estimate conifer and hardwood covers separately. Please note: These cover values should not include the coverage of suppressed understory trees.

**%Low-Medium Tree:** The total aerial cover (canopy closure) of all live understory low to medium height tree species, disregarding overlap of individual trees and shrubs. This category contains recruits of overstory tree species (with seedlings and saplings in the understory) and understory tree species that typically do not make up the overstory canopy (e.g. trees that typically do not attain a height >10m).

**% Shrub:** The total aerial cover (canopy closure) of all live shrub species disregarding overlap of individual shrubs.

**% Herb:** The total aerial cover (canopy closure) of all herbaceous species, disregarding overlap of individual herbs.

**% Total Vascular plants:** The total aerial cover of all vegetation. This is an estimate of the absolute vegetation cover, disregarding overlap of the various tree, shrub, and/or herb layers.

**% Total Non-vascular plants:** The total cover of all lichens and bryophytes (mosses, liverworts, hornworts) on substrate surfaces (not standing or inclined trees).

**Modal height for conifer/hardwood tree, shrub, and herbaceous categories (optional)**

If height values are important in your vegetation survey project, provide an ocular estimate of height for each category listed. Record an average height value per each category by estimating the mean height for each group. Please use the following height intervals to record a height class: 01=<1/2m, 02=1/2-1m, 03=1-2m, 04=2-5m, 05=5-10m, 06=10-15m, 07=15-20m, 08=20-35m, 09=35-50m, 10=>50m.

**Caveats**

Please consult with the members of the vegetation committee for advice and feedback on proposed vegetation surveys prior on initiating projects.

Notes on the Order and Division of Labor for Data Collection: As with every procedure, there are always more and less efficient ways to collect the information requested. Although we respect each field crews' option to choose in what order they collect the data, we suggest the following general rules:

- Work with teams of two for each plot collected.
- Both team members can determine the plot shape and size and lay out the tapes and mark the edges for the plot boundary (see below).
- The two person teams can also divide up tasks of data collection with one member collecting location, environmental (slope, aspect, geology, soil texture, etc.) and plot description information while the other begins the species list. Thus, two clipboards are useful and data sheets that are at first separated (not stapled).
- Following the making of the initial species list and collection of location and environmental data both team members convene to do the estimation of plant cover by species followed by the estimation of total vegetation cover and cover by layer.
- Following that process, the estimation of cover by the up to 10 height strata classes and the listing of the diagnostic species for each is done collaboratively.
- This is followed by the estimation of the coarse fragment information, again done collaboratively.

For egalitarian and familiarization purposes we suggest that the roles be switched regularly between the team members and that if multiple teams are being used in a larger project, that each team member switches frequently between teams, building all-important calibration, and camaraderie among the whole group.

Suggestions for Laying out Plots: If you are laying out a circular plot, work with two or more people. One person stands at the center of the plot and holds the tape case while the other walks the end of the tape out to the appointed distance (radius 5.6 for 100 m<sup>2</sup> circle, radius 11.3 m for a 400 m<sup>2</sup> circle, and radius 17.6 m for a 1000m<sup>2</sup> circle). The walker then fixes the tape end with a pin flag and walks back to the center where he/she instructs the center person to walk in the opposite direction of the already laid out tape radius, stretching the rest of the tape to an equal

length (another 11.3 or 17.6 m) to the opposite edge of the plot, where he/she affixes it with another pin flag. This process is again repeated with another tape laid out perpendicular to the first so that an “+” shape is created. The margins of the circle can be further delineated by measuring to the center of the circle with an optical tape measure (rangefinder) and marking mid points between the four ends of the crossed tapes.

When laying out square or rectangular plots work with two or more people per team. If doing a rectangle, determine the long axis of the plot first and have one person be stationed at the zero m end of the tape while the other person walks the unrolling tape case out to the appropriate length. The stationary end person can guide the walker, keeping them moving in a straight line. Once that tape is laid out and the far end staked, the team lays out another tape perpendicular to the first, either at one end, using the same type of process. This establishes the width of the rectangle (or square). Using an optical rangefinder and pin-flags, or colored flagging the team can further mark additional points along the other parallel long axis and short axis of the plot (every 5 m for shorter plots or every 10 m for longer plots is suggested) so that the entire plot boundary can be easily visualized.

#### References:

Barbour M.G., J.H. Burk, and W.D. Pitts 1987. Terrestrial Plant Ecology, Second Edition. Benjamin/Cummings Publishing Co. Menlo Park, CA. 634 pages.

Sawyer and Keeler-Wolf. 1995. Manual of California Vegetation. California Native Plant Society, Sacramento, CA. 471 pages

The Nature Conservancy and Environmental Systems Research Institute. 1994. Final Draft, Standardized National Vegetation Classification System. Prepared for United States Department of the Interior, National Biological Survey, and National Park Service. Arlington, VA.

Complete document available at the following website:

<http://biology.usgs.gov/npsveg/fieldmethods.html>

#### Suggested Equipment:

Equipment List: Prices as of May 2000, toll free orders from Forestry Suppliers (1-800-647-5368) (item numbers in parentheses)

Chaining pins, surveyor steel (#39167)	\$21.50
Fiberglass tapes 2 - 165'/50 m (#39972)	\$42.90
Logbook cover 8 ½ “ x 12” (#53200)	\$23.95
Perforated flagging (#57960)	\$1.95
UTM Coordinate Grid (#45019)	\$16.95
Rangefinder, 10-75m (#38973)	\$51.60
Silva Compass w/ clinometer (#37036)	\$43.90
Garmin GPS 12XL (#39095, #39111)	\$244.90



**CALIFORNIA NATIVE PLANT SOCIETY RELEVÉ FIELD FORM CODE LIST** (revised 3/0107)

**MACRO TOPOGRAPHY**

- 00 Bench
- 01 Ridge top (interfluvial)
- 02 Upper 1/3 of slope
- 03 Middle 1/3 of slope
- 04 Lower 1/3 of slope (lowslope)
- 05 Toeslope (alluvial fan/bajada)
- 06 Bottom/plain
- 07 Basin/wetland
- 08 Draw
- 09 Other
- 10 Terrace (former shoreline or floodplain)
- 11 Entire slope
- 12 Wash (channel bed)
- 13 Badland (complex of draws & interfluvial)
- 14 Mesa/plateau
- 15 Dune/sandfield
- 16 Pediment
- 17 Backslope (cliff)

**MICRO TOPOGRAPHY**

- 01 Convex or rounded
- 02 Linear or even
- 03 Concave or depression
- 04 Undulating pattern
- 05 Hummock or Swale pattern
- 06 Mounded
- 07 Other

**SITE IMPACTS**

- 01 Development
- 02 ORV activity
- 03 Agriculture
- 04 Grazing
- 05 Competition from exotics
- 06 Logging
- 07 Insufficient population/stand size
- 08 Altered flood/tidal regime
- 09 Mining
- 10 Hybridization
- 11 Groundwater pumping
- 12 Dam/inundation
- 13 Other
- 14 Surface water diversion
- 15 Road/trail construction/maint.
- 16 Biocides
- 17 Pollution
- 18 Unknown
- 19 Vandalism/dumping/litter
- 20 Foot traffic/trampling
- 21 Improper burning regime
- 22 Over collecting/poaching
- 23 Erosion/runoff
- 24 Altered thermal regime
- 25 Landfill
- 26 Degrading water quality
- 27 Wood cutting
- 28 Military operations
- 29 Recreational use (non ORV)
- 30 Nest parasitism
- 31 Non-native predators
- 32 Rip-rap, bank protection
- 33 Channelization (human caused)
- 34 Feral pigs
- 35 Burros
- 36 Rills
- 37 Phytogenic mounding
- 38 Sudden oak death syndrome (SODS)

**PARENT MATERIAL**

- IGTU Igneous (type unknown)
- VOLC General volcanic extrusives
- RHYO Rhyolite
- ANDE Andesite
- BASA Basalt
- ASHT Ash (of any origin)
- OBSI Obsidian
- PUMI Pumice
- PYFL Pyroclastic flow
- VOFL Volcanic flow
- VOMU Volcanic mud
- INTR General igneous intrusives
- GRAN Granitic (generic)
- MONZ Monzonite
- QUDI Quartz diorite
- DIOR Diorite
- GABB Gabbro
- DIAB Diabase
- PERI Peridotite
- METU Metamorphic (type unknown)
- GNBG Gneiss/biotite gneiss
- SERP Serpentine
- SCHI Schist
- SESC Semi-schist
- PHYL Phyllite
- SLAT Slate
- HORN Hornfels
- BLUE Blue schist
- MARB Marble
- SETU Sedimentary (type unknown)
- BREC Breccia (non-volcanic)
- CONG Conglomerate
- FANG Fanglomerate
- SAND Sandstone
- SHAL Shale
- SILT Siltstone
- CACO Calcareous conglomerate
- CASA Calcareous sandstone
- CASH Calcareous shale
- CASI Calcareous siltstone
- DOLO Dolomite
- LIME Limestone
- CALU Calcareous (origin unknown)
- CHER Chert
- FRME Franciscan melange
- GREE Greenstone
- ULTU Ultramafic (type unknown)
- MIIG Mixed igneous
- MIME Mixed metamorphic
- MISE Mixed sedimentary
- MIRT Mix of two or more rock types
- GLTI Glacial till, mixed origin, moraine
- LALA Large landslide (unconsolidated)
- DUNE Sand dunes
- LOSS Loess
- CLAL Clayey alluvium
- GRAL Gravelly alluvium
- MIAL Mixed alluvium
- SAAL Sandy alluvium (most alluvial fans and washes)
- SIAL Silty alluvium
- OTHE Other than on list

**SOIL TEXTURE**

- COSA Coarse sand
- MESN Medium sand
- FISN Fine sand
- COLS Coarse, loamy sand
- MELS Medium to very fine, loamy sand
- MCSL Moderately coarse, sandy loam
- MESA Medium to very fine, sandy loam
- MELO Medium loam
- MESL Medium silt loam
- MESI Medium silt
- MFCL Moderately fine clay loam
- MFSA Moderately fine sandy clay loam
- MFSL Moderately fine silty clay loam
- FISA Fine sandy clay
- FISC Fine silty clay
- FICL Fine clay
- SAND Sand (class unknown)
- LOAM Loam (class unknown)
- CLAY Clay (class unknown)
- UNKN Unknown
- PEAT Peat
- MUCK Muck

**DOMINANT VEGETATION GROUP**

***Trees:***

- TBSE Temperate broad-leaved seasonal evergreen forest
- TNLE Temperate or subpolar needle-leaved evergreen forest
- CDF Cold-deciduous forest
- MNDF Mixed needle-leaved evergreen-cold deciduous forest
- TBEW Temperate broad-leaved evergreen woodland
- TNEW Temperate or subpolar needle-leaved evergreen woodland
- EXEW Extremely xeromorphic evergreen woodland
- CDW Cold-deciduous woodland
- EXDW Extremely xeromorphic deciduous woodland
- MBED Mixed broad-leaved evergreen-cold deciduous woodland
- MNDW Mixed needle-leaved evergreen-cold deciduous woodland

***Shrubs:***

- TBES Temperate broad-leaved evergreen shrubland
- NLES Needle-leaved evergreen shrubland
- MIES Microphyllous evergreen shrubland
- EXDS Extremely xeromorphic deciduous shrubland
- CDS Cold-deciduous shrubland
- MEDS Mixed evergreen-deciduous shrubland
- XMED Extremely xeromorphic mixed evergreen-deciduous shrubland

***Dwarf Shrubland:***

- NMED Needle-leaved or microphyllous evergreen dwarf shrubland
- XEDS Extremely xeromorphic evergreen dwarf shrubland
- DDDS Drought-deciduous dwarf shrubland
- MEDD Mixed evergreen cold-deciduous dwarf shrubland

***Herbaceous:***

- TSPG Temperate or subpolar grassland
- TGST Temperate or subpolar grassland with sparse tree
- TGSS Temperate or subpolar grassland with sparse shrublayer
- TGSD Temperate or subpolar grassland with sparse dwarf shrub layer
- TFV Temperate or subpolar forb vegetation
- THRV Temperate or subpolar hydromorphic rooted vegetation
- TAGF Temperate or subpolar annual grassland or forb vegetation

***Sparse Vegetation:***

- SVSD Sparsely vegetated sand dunes
- SVCS Sparsely vegetated consolidated substrates

**Simplified Key to Soil Texture**  
(Adapted from Brewer and McCann 1982)

Place about three teaspoons of soil in the palm of your hand. Take out any particles  $\geq 3$  mm in size.

**A. Does soil remain in ball when squeezed in your hand palm?**

Yes, soil does remain in a ball when squeezed..... **B**

No, soil does not remain in a ball when squeezed..... **sand**

Very coarse texture..... SAND Sand (class unknown)  
Moderately coarse texture..... COSA Coarse sand  
Moderately fine texture..... MESN Medium sand  
FISN Fine sand

**B. Add a small amount of water until the soil feels like putty. Squeeze the ball between your thumb and forefinger, attempting to make a ribbon that you push up over your finger. Does soil make a ribbon?**

Yes, soil makes a ribbon; though it may be very short..... **C**

No, soil does not make a ribbon..... **loamy sand**

Very gritty with coarse particles..... COLS Coarse, loamy sand  
Moderately to slightly gritty with medium to fine particles.....MELS Medium to very fine, loamy sand

**C. Does ribbon extends more than one inch?**

Yes, soil extends > 1 inch. .... **D**

No, soil does not extend > 1 inch. .... Add excess water

Soil feels gritty..... **loam or sandy loam**

Very gritty with coarse particles..... LOAM Loam (class unknown)  
Moderately gritty with medium to fine particles..... MCSL Moderately coarse, sandy loam  
Slightly gritty..... MESA Medium to very fine, sandy loam  
MELO Medium loam

Soil feels smooth..... **silt loam**

MESIL medium silt loam

**D. Does soil extend more than 2 inches?**

Yes, ribbon extends more than 2 inches, and does not crack if bent into a ring..... **E**

No, soil breaks when 1–2 inches long; cracks if bent into a ring..... Add excess water

Soil feels gritty..... **sandy clay loam or clay loam**

Very gritty..... MFSA Moderately fine sandy clay loam  
Slightly gritty..... MFCL Moderately fine clay loam

Soil feels smooth..... **silty clay loam or silt**

Moderately fine texture..... MFSL Moderately fine silty clay loam  
Very fine texture..... MESI Medium silt

**E. Soil makes a ribbon 2+ inches long; does not crack when bent into a ring..... Add excess water**

Soil feels gritty..... **sandy clay or clay**

Very gritty..... CLAY Clay (class unknown)  
Slightly gritty..... FISA Fine sandy clay  
FICL Fine clay

Soil feels smooth..... **silty clay**

FISC Fine silty clay

## Artificial Key to the Systems and Classes

### Key to the Systems

1. Water regime influenced by oceanic tides, and salinity due to ocean-derived salts 0.5% or greater.
  2. Semi-enclosed by land, but with open, partly obstructed or sporadic access to the ocean. Halinity wide-ranging because of evaporation or mixing of seawater with runoff from land..... ESTUARINE
  - 2'. Little or no obstruction to open ocean present. Halinity usually euhaline; little mixing of water with runoff from land..... 3
    3. Emergents, trees, or shrubs present ..... ESTUARINE
    - 3'. Emergents, trees, or shrubs absent..... MARINE
- 1'. Water regime not influenced by oceanic tides, or if influenced by oceanic tides, salinity less than 0.5%
  4. Persistent emergents, trees, shrubs, or emergent mosses cover 30% or more of the area..... PALUSTRINE
  - 4'. Persistent emergents, trees, shrubs, or emergent mosses cover less than 30% of substrate but nonpersistent emergents may be widespread during some seasons of year .....5
    5. Situated in a channel; water, when present, usually flowing.....RIVERINE
    - 5'. Situated in a basin, catchment, or on level or sloping ground; water usually not flowing ..... 6
      6. Area 8 ha (20 acres) or greater ..... LACUSTRINE
      - 6'. Area less than 8 ha.....7
        7. Wave-formed or bedrock shoreline feature present or water depth 2 m (6.6 feet) or more . . . . . LACUSTRINE
        - 7'. No wave-formed or bedrock shoreline feature present and water > 2 m deep ..... PALUSTRINE

### Key to the Classes

1. During the growing season of most years, aerial cover by vegetation is less than 30%.
  2. Substrate a ridge or mound formed by colonization of sedentary invertebrates (corals, oysters, tube worms) . . . REEF
  - 2'. Substrate of rock or various-sized sediments often occupied by invertebrates but not formed by colonization of sedentary invertebrates ..... 3
    3. Water regime subtidal, permanently flooded, intermittently exposed, or semipermanently flooded. Substrate usually not soil..... 4
      4. Substrate of bedrock, boulders, or stones occurring singly or in combination covers 75% or more of the area ..... ROCK BOTTOM
      - 4'. Substrate of organic material, mud, sand, gravel, or cobbles with less than 75% areal cover of stones, boulders, or bedrock..... UNCONSOLIDATED BOTTOM
    - 3'. Water regime irregularly exposed, regularly flooded, irregularly flooded, seasonally flooded, temporarily flooded, intermittently flooded, saturated, or artificially flooded. Substrate often a soil ..... 5
      5. Contained within a channel that does not have permanent flowing water (i.e., Intermittent Subsystem of Riverine System or Intertidal Subsystem of Estuarine System) ..... STREAMBED
      - 5'. Contained in a channel with perennial water or not contained in a channel..... 6
        6. Substrate of bedrock, boulders, or stones occurring singly or in combination covers 75% or more of the area ..... ROCKY SHORE
        - 6'. Substrate of organic material, mud, sand, gravel, or cobbles; with less than 75% of the cover consisting of stones, boulders, or bedrock. .... UNCONSOLIDATED SHORE
  - 1'. During the growing season of most years, percentage of area covered by vegetation 30% or greater.
    7. Vegetation composed of pioneering annuals or seedling perennials, often not hydrophytes, occurring only at time of substrate exposure.....8
      8. Contained within a channel that does not have permanent flowing water. .... STREAMBED (VEGETATED)
      - 8'. Contained within a channel with permanent water, or not contained in a channel  
..... UNCONSOLIDATED SHORE (VEGETATED)
    - 7'. Vegetation composed of algae, bryophytes, lichens, or vascular plants that are usually hydrophytic perennials . . . . .9
      9. Vegetation composed predominantly of nonvascular species ..... 10
        10. Vegetation macrophytic algae, mosses, or lichens growing in water or the splash zone of shores . . . . . AQUATIC BED
        - 10'. Vegetation mosses or lichens usually growing on organic soils and always outside the splash zone of shores.....MOSS-LICHEN WETLAND
      - 9'. Vegetation composed predominantly of vascular species ..... 11
        11. Vegetation herbaceous..... 12
          12. Vegetation emergents ..... EMERGENT WETLAND
          - 12'. Vegetation submergent, floating-leaved, or floating ..... AQUATIC BED
        - 11'. Vegetation trees or shrubs..... 13
          13. Dominants less than 6 m (20 feet) tall ..... SCRUB-SHRUB WETLAND
          - 13'. Dominants 6 m tall or taller ..... FORESTED WETLAND

**CALIFORNIA NATIVE PLANT SOCIETY RELEVÉ FIELD FORM**

(Revised 8/23/07)

Page \_\_\_\_\_ of Relevé # \_\_\_\_\_

**FOR OFFICE USE ONLY**

Polygon # _____ or Relevé # _____	Permanent Number: _____
Date _____ Airphoto # _____ MMDD / YYYY	Community Name: _____ Community Number: _____ Occurrence Number: _____
County _____	Source Code: _____
USGS Quad. _____ 7.5' or 15' (Circle one)	Quad Code: _____ Quad Name: _____ Map Index Number: _____
CNPS Chapter _____	Update: Yes No (Circle one)

Landowner \_\_\_\_\_

Contact Person \_\_\_\_\_

Address \_\_\_\_\_

City _____	Zip _____	Phone number _____
------------	-----------	--------------------

Observers \_\_\_\_\_

Relevé plot shape (square, rectangle, triangle, circle, entire stand) \_\_\_\_\_ **NOTE:** Forest/woodland plots should be 1000m<sup>2</sup> if upland or 400m<sup>2</sup> if riparian.  
 Relevé plot size (length and width of rectangle, or circle-diameter) \_\_\_\_\_ (m.) All shrub plots should be 400m<sup>2</sup>. Herb plots should be 100 or 10m<sup>2</sup>\*.  
 (1000m<sup>2</sup>) \*Please consult with CNPS Vegetation Ecologist on herb plots.  
 Study Plot Revisit? Yes or No (Circle one) For circle radiuses: 5.64m (100m<sup>2</sup>), 11.28m (400m<sup>2</sup>), 17.84m  
 Photo Interpreter Community Code for Polygon \_\_\_\_\_  
 Other polygons of same type? Yes or No Is plot representative of whole polygon? Yes or No (Circle one) If not, why not? \_\_\_\_\_

GPS File # \_\_\_\_\_ GPS name (or points in file) \_\_\_\_\_ Start Time \_\_\_\_:\_\_\_\_ (am or pm) GPS Datum (from GPS setup) (e.g. WGS 84, NAD 27) \_\_\_\_\_  
 File type: Point or Polygon (circle one) Releve: UTME \_\_\_\_\_ UTMN \_\_\_\_\_ Error ± \_\_\_\_\_ ft/m UTM Zone \_\_\_\_\_  
 Transect: Start UTME \_\_\_\_\_ UTMN \_\_\_\_\_ End: UTME \_\_\_\_\_ UTMN \_\_\_\_\_

Elevation (ft.) \_\_\_\_\_ Slope (°) \_\_\_\_\_ Aspect (°) \_\_\_\_\_ Topography: Macro \_\_\_\_\_ Micro \_\_\_\_\_

**VEGETATION DESCRIPTION** *See code list for italicized fields*

Dominant Layer \_\_\_ 0-0.5 m, \_\_\_ 0.5-5 m, \_\_\_ >5 m Preliminary Alliance Name \_\_\_\_\_  
 Stand Size \_\_\_ <1 acre, \_\_\_ 1-5 acres, \_\_\_ >5 acres Dominant Vegetation Group \_\_\_\_\_ (use codes from code list)

Phenology: Ground \_\_\_\_\_ Shrub \_\_\_\_\_ Tree \_\_\_\_\_  
 (Early, Peak, Late)

Wetland Community Type \_\_\_\_\_ (Wetland or Upland)  
 If Community Type = Wetland (see Artificial Keys to Cowardin Systems and Names)  
 Cowardin System \_\_\_\_\_ Subsystem \_\_\_\_\_ Class \_\_\_\_\_  
 Distance to water (m): Vertical \_\_\_\_\_ Horizontal \_\_\_\_\_ Channel form (if riverine) \_\_\_\_\_  
 (Straight, Meandering, Braided)

Adjacent Alliance	Location (e.g., North, South, East, or West of stand)	Description (up to 4 species by layer)

**Photographs** – Note which camera, photo JPEG/frame numbers, and photo direction relative to plot.

--	--	--

**CALIFORNIA NATIVE PLANT SOCIETY RELEVÉ FIELD FORM**

Page \_\_\_\_\_ of Relevé # \_\_\_\_\_

**STAND AND ENVIRONMENTAL DESCRIPTION**

Trend code \_\_\_\_\_ *Site Impact codes* \_\_\_\_\_

1. Increasing 2. Stable 3. Decreasing (List codes in order, with most significant first)  
4. Fluctuating 5. Unknown

Site Intensity \_\_\_\_\_

1. Light 2. Moderate 3. Heavy (List beneath each impact code)

**Site Location and Plot Description** – Describing where the plot is located and what the main vegetation and environmental features are


**Site History** – Including observations of fire scars, insect/disease damage, grazing/browsing, human disturbance


**Sensitive Species** – List species observed and GPS UTM's; Estimate size and extent of local populations


**Unknown Specimens** – List code, identification notes (e.g. Genus, condition of specimen) of unknowns


**Additional Comments** – Including animal observations, anthropological observations, abiotic features


**Surface Coarse Fragments and Soils Information** (see cover class intervals-below ↓)

Type:	Fines	Gravel	Cobble	Stone	Boulders	Bedrock	Litter	Water	Living stems
Descriptor:	Including sand, mud	2mm-7.5 cm diameter	7.5-25 cm diam	25-60cm diam.	>60cm diam.	Including outcrops	Organic matter covering ground	Standing or running water	Vascular plants at ground surface
Cover class (see below):									
% Cover*:									

\*note all surface fragments, non-vegetation, living stems, etc., should add up to 100%

Cover Class Intervals: 1 (<1%), 2 (1-5%), 3a (>5-15%), 3b (>15-25%), 4 (>25-50%), 5 (>50-75%), 6 (>75%)

**% Bioturbation** \_\_\_\_\_ **Soil Texture** \_\_\_\_\_ (Code) **Parent Material** \_\_\_\_\_ (Code)





## **APPENDIX C. MONITORING REPORT OUTLINE**

### **I. Monitoring Report Content (one report per DPA)**

#### **A. Project Information**

1. Project name
2. Grantee name, address, and phone number
3. DPA Network Map (highlighting current DPA)
4. Brief historical management summary
5. Important habitat qualities present/ special status species
6. Major threats impacting this DPA
7. Objectives to mitigate major threats

#### **B. Management Activity Monitoring**

1. Location of the restoration actions (DPA map necessary)
2. Methodology of monitoring (including before and after photos)
3. Dates summary of work completed during monitoring period
4. Specific amount (acres) in which restoration actions were taken
5. Amount of chemical applied (if applicable)
6. Summary of effectiveness of restoration actions

#### **C. Monitoring to Inform Management**

1. Location of the monitoring (e.i. grid map)
2. Methodology of monitoring (following protocol given in Appendix A)
3. Dates summary of work completed during monitoring period
4. Results with appropriate tables and maps as outlined in Appendix A (Invasive Species, Special status Species, Feral Pigs and Habitats most affected by invasive species)
6. Summary of findings (change in percent cover, species found etc.)



D. Baseline Monitoring

1. Location of the monitoring (e.i. releve plot map)
2. Methodology of monitoring (following protocol given in Appendix B)
3. Dates summary of work completed during monitoring period
4. Results with appropriate tables and maps as outlined in Appendix B
5. Summary of findings (dominant vegetation alliances and species, changes since last monitoring event)

E. Adaptive Management

1. How restoration actions are meeting stated objectives
2. How restoration actions are not meeting stated objectives
3. Recommendations to better meet stated objectives (if necessary)

Monitoring Report Visuals

Report Section	Visuals
A. Project Information	DPA Network Map (highlighting current DPA)
B. Management Activity Monitoring	Location Map
	Restoration Area Map (highlighting treatment)
	Before and after photos (current monitoring period and overall)
	Treatment area (acres)
C. Monitoring to Inform Management	Chemical applied (if applicable)
	Grid Location Map highlighting Grids visited in current monitoring period
D. Baseline Monitoring	All tables and maps outlined in Monitoring Protocol (Appendix A)
	Releve Location Map
E. Adaptive Management	All tables and maps outlined in Monitoring Protocol (Appendix B)
	Any necessary visuals for potential changes in management strategy

II. Appendices

- A. Original Data Sheets and Technical Appendices
- B. Photographic Record of the Site during most recent monitoring visit at record stations