## State of California

Department of Fish and Wildlife

## Memorandum

Date: September 3, 2025

To: Erin Chappell

Regional Manager

Bay Delta Region

From: Margaret Johnson

**Environmental Scientist** 

Bay Delta Region

Subject: 2025 Summer Townet Survey Delta Smelt Abundance Index

The annual California Department of Fish and Wildlife (CDFW) Summer Townet Survey abundance index for Delta Smelt (*Hypomesus transpacificus*) in 2025 is 0.0. This is the tenth year in which the abundance index was equal to zero. The first zero index was reported in 2015 and has reported a 0.0 index every year since 2018 (Figure 1).

The Summer Townet Survey sampling period consists of six biweekly surveys, beginning in June and ending in late August. During each survey, young pelagic fishes are sampled from 40 fixed stations which consist of 31 index stations and 9 supplemental non-index stations (Figure 2). These sampling stations are located in the upper San Francisco Estuary from eastern San Pablo Bay and Napa River and upstream through to the Sacramento Deep Water Ship Channel (SDWSC) and lower San Joaquin River.

The Delta Smelt annual index is generated from an average of the first two survey indices. Survey indices for Delta Smelt are calculated by multiplying the catch at each index station by a station-specific weighting factor, summing the weighted catches of all index stations, and dividing the total by 1000. More information on index calculation can be found at on the department website for the <a href="Summer Townet Survey">Summer Townet Survey</a> (ca.gov).

The first and second Summer Townet surveys of 2025 were conducted during June 2-6 and June 16-18. During these sampling periods, no Delta Smelt were collected at index or non-index stations, indicating that Delta Smelt continue to occur at very low levels and detection is limited. Additionally, no Delta Smelt were observed at index or non-index stations during subsequent surveys 3-6 in 2025.

However, other monitoring studies may have observed small numbers of Delta Smelt during the summer season. Preliminary data from the U.S. Fish and Wildlife Service Enhanced Delta Smelt Monitoring Program (EDSM) indicate that two potential Delta Smelt have been detected through Kodiak trawl sampling on June 20 and July 15. These fish were caught in Suisun Marsh and retained for genetic confirmation. More information on EDSM catch can be obtained from the EDSM Daily Report.

Cc: Nicole Athearn

Steven Slater

Kathy Hieb

Dr. Tim Malinich

Colin Brennan

Stephanie Fong




Figure 1. Summer Townet Survey Delta Smelt annual abundance indices 1959-2025.




Figure 2. Distribution of Summer Townet Survey index and non-index stations in the upper San Francisco Estuary.