

**California Wildlife Habitat Relationships System**  
**California Department of Fish and Wildlife**  
**California Interagency Wildlife Task Group**

---

DUSKY-FOOTED WOODRAT

*Neotoma fuscipes*

Family: MURIDAE  
M127

Order: RODENTIA

Class: MAMMALIA

Written by: P. Brylski

Reviewed by: H. Shellhammer

Edited by: R. Duke

Updated by: CWHR Program Staff, August 2005 and January 2008

#### DISTRIBUTION, ABUNDANCE, AND SEASONALITY

The dusky-footed woodrat is common in California. It is found in both the Coast Ranges and interior. Also widespread along the western slope of the Sierra Nevada, mostly below 2150 m (7000 ft). What was the southern part of the range of *N. fuscipes* is now considered the range of *N. macrotis*. Once a subspecies of *N. fuscipes*, *N. macrotis* has been elevated to species status (Matocq 2002a, 2002b). (Note: Because *N. fuscipes* and *N. macrotis* are still undergoing taxonomic revision, the range of the federally-endangered subspecies *N. fuscipes riparia* has been mapped in CWHR as part of the range of both species.) Generally absent from cultivated land and open grasslands of Central Valley. Common to abundant in forest habitats of moderate canopy and moderate to dense understory. Can be abundant in chaparral habitats.

#### SPECIFIC HABITAT REQUIREMENTS

**Feeding:** Feeds mainly on woody plants, especially live oak, maple, coffeeberry, alder, and elderberry when available (Linsdale and Tevis 1951). Other woody plants eaten elsewhere; English (1923) listed 37 species of plants eaten by the dusky-footed woodrat. Also eats fungi, flowers, grasses, acorns. Forages on ground, in bushes, and in trees.

**Cover:** Prefers moderate canopy in a variety of habitats. Houses are built of sticks and leaves at the base of, or in a tree, around a shrub, or at the base of a hill. Houses may measure 2.4 m (8 ft) in height and 2.4 m (8 ft) in diameter (English 1923).

**Reproduction:** Nests are located in the stick house, and are constructed of shredded grass, leaves, and other miscellaneous materials (e.g., bird feathers). Vestal (1938) reported the following average dimensions: 112 mm (4.4 in) x 101 mm (4 in) with a depth of 85 mm (3.4 in). Abundance probably limited by availability of nest-building materials. Competition for houses is constant and intense (Linsdale and Tevis 1951).

**Water:** Drinks water, but may be sustained by leafy vegetation and fungi.

**Pattern:** Prefers forest habitats with moderate canopy, year-round greenery, a brushy understory, and suitable nestbuilding materials. Well-developed understory at base of a single evergreen may be suitable for a single individual.

#### SPECIES LIFE HISTORY

**Activity Patterns:** Mostly nocturnal. May reduce activity on moonlit or rainy nights. Active year-round.

**Seasonal Movements/Migration:** None.

**Home Range:** In Sonoma Co., home ranges (for what is now considered *N. macrotis*) averaged 0.23 ha (0.58 ac) for males, 0.19 ha (0.48 ac) for females, and 0.17 ha (0.43 ac) for juveniles; densities reached a peak of 20 individuals/ha (8/ac) in late summer (Cranford 1977). In Monterey Co., an individual may confine its lifetime activity around a single tree, or range over 18.7 ha (46.2 ac) (Linsdale and Tevis 1951). In chaparral habitat, density was reported to reach 18.8/ ha (7.5/ac) (Bleich 1973).

**Territory:** The nest is defended against competitors.

**Reproduction:** Breeds from December to September, with a peak in mid-spring. Litter size averages 2-3 young (range 1-4) (Linsdale and Tevis 1951, Verner and Boss 1980). One to 5 litters per yr. Females probably are promiscuous.

**Niche:** The dusky-footed woodrat is heavily preyed upon by owls, coyotes, bobcats, hawks, and perhaps snakes. Other small mammals and amphibians and reptiles are known to use woodrat houses. Cattle grazing probably reduces carrying capacity for woodrats by removing cover. Wildfires and prescribed burning are likely to be detrimental by destroying houses.

## REFERENCES

- Bleich, V. C. 1973. Ecology of rodents at the United States Naval Weapon Station, Seal Beach, Fallbrook Annex, San Diego County, California. M.A. Thesis, California State Univ., Long Beach. 102pp.
- Cranford, J. A. 1977. Home range and habitat utilization by *Neotoma fuscipes* as determined by radiotelemetry. *J. Mammal.* 58:165-172.
- English, P. F. 1923. The dusky-footed woodrat (*Neotoma fuscipes*). *J. Mammal.* 4:1-9.
- Hooven, E. F. 1959. Dusky-footed woodrat in young Douglas-fir. Oregon For. Lands Res. Cent., Corvallis. Res. Note No. 41. 24pp.
- Linsdale, J. M., and L. P. Tevis, Jr. 1951. The dusky-footed woodrat. Univ. California Press, Berkeley. 664pp.
- Matocq, M. D. 2002a. Phylogeographical structure and regional history of the dusky-footed woodrat, *Neotoma fuscipes*. *Molecular Ecology* 11:229-242.
- Matocq, M. D. 2002b. Morphological and molecular analysis of a contact zone in the *Neotoma fuscipes* species complex. *Journal of Mammalogy* 83(3):866-883.
- Verner, J., and A. S. Boss. 1980. California wildlife and their habitats: Western Sierra Nevada. USDA, For. Serv. Gen. Tech. Rep. PSW-37. 439pp.
- Vestal, E. H. 1938. Biotic relations of the woodrat (*Neotoma fuscipes*) in the Berkeley Hills. *J. Mammal.* 19:1-36.
- Williams, D. F. 1986. Mammalian species of special concern in California. Calif. Dept. Fish and Game, Sacramento. Admin. Rep. 86-1. 112pp.