

**California Wildlife Habitat Relationships System**  
**California Department of Fish and Wildlife**  
**California Interagency Wildlife Task Group**

---

GREAT BASIN COLLARED LIZARD

*Crotaphytus bicinctores*

Family: CROTAPHYTIDAE

Order: SQUAMATA

Class: REPTILIA

R017

Written by: T. Kucera, 1998

Edited by: CWHR Program Staff, March 2000 and February 2008

#### DISTRIBUTION, ABUNDANCE, AND SEASONALITY

The Great Basin collared lizard is widely distributed throughout the arid and semiarid regions of the Mojave, Sonoran, and Southeastern Great Basin deserts. It is generally restricted to areas with rocky substrates, slopes, gullies, washes, canyons, and sometimes rock piles, although occasionally can be found up to a mile from extensive rocky habitat (McGuire 1996). It is most common in desert succulent shrub, desert scrub, and desert wash habitats. The Great Basin collared lizard is active in the spring and summer and to a lesser extent in the early fall. Southern populations are active earlier than northern populations, and remain active later in the year. It probably exhibits considerable variation in habits over its range (Johnson et al. 1948, Stebbins 1985, Fitch 1956, Legler and Fitch 1957, Yedlin and Ferguson 1973, Sanborn and Loomis 1979).

#### SPECIFIC HABITAT REQUIREMENTS

**Feeding:** The diet of this species consists largely of arthropods and small vertebrates (Stebbins 1985, McGuire 1996). Arthropods include orthopterans, coleopterans, hemipterans, hymenopterans, lepidopterans, and arachnids. Vertebrates include *Uta*, *Sceloporus*, *Cnemidophorus*, *Phrynosoma*, and *Xantusia*. Plant material is also occasionally consumed.

**Cover:** This species prefers rocky areas and seeks cover under rocks and in cracks and crevices and rodent holes (Stebbins 1985), occasionally bounding bipedally from stone to stone when disturbed (McGuire 1996).

**Reproduction:** The species lays eggs and presumably constructs its own nest but there are no reports. The closely related *C. collaris* nests in rocky areas (Yedlin and Ferguson 1973) and sometimes lays eggs in tunnels under rocks (Fitch 1956). Friable, well drained soil is probably required for nesting.

**Water:** No information on water requirements. Probably does not require water, as it occurs in arid habitats.

**Pattern:** The Mojave black-collared lizard occupies slopes, rock outcrops, gullies, washes and other areas with small vertical perches. It often sits on rock perches and watches for prey, predators, or perhaps conspecifics.

#### SPECIES LIFE HISTORY

**Activity Patterns:** This species is diurnal, active all day when temperatures are moderate or warm, and showing a bimodal or morning activity when temperatures are high (Stebbins 1985). It hibernates in winter, and may aestivate. It may become active as early as 17 April in the north (McGuire 1996).

Seasonal Movements/Migration: This lizard is not known to migrate..

Home Range: Nothing is known about the size of home ranges of the Mojave black-collared lizard. There are no reports of abundance but it is often possible to see as many as 5 individuals in 1 ha (2.5 ac).

Territory: This lizard is aggressive; adult males behave aggressively toward each other but will tolerate subordinate males (Stebbins 1985).

Reproduction: Courtship occurs in early spring. Eggs are laid in June and July; neonates have been observed in eastern Oregon in August (Brooking 1934). In Utah, females reached reproductive maturity at 85 mm SVL (Andre and MacMahon 1980). Mean clutch size was 5.4, and ranged from 3 to seven. Larger females produced larger clutches. Occasionally 2 clutches are laid (Johnson et al. 1948, Stebbins 1985).

Niche: Great Basin collared lizards are probably eaten by Masticophis, avian predators, and perhaps mammalian predators. The young are taken by loggerhead shrikes.

General Comments: This species has been subject to taxonomic confusion for several decades. McGuire (1996) revised the taxonomy of the Crotaphytidae based on morphological and genetic data, elevating *C. bicinctores* to specific status, separate from *C. vestigium* of the Peninsular range of southern California and *C. collaris* that occurs further east.

## REFERENCES

- Andre, J. B. and J. A. MacMahon. 1980. Reproduction in three sympatric lizard species from west-central Utah. *Great Basin Naturalist* 40:68-72.
- Brooking, W. J. 1934. Some reptiles and amphibians from Malheur County, inn eastern Oregon. *Copeia* 1934:93-95.
- Fitch, H. S. 1956. An ecological study of the collared lizard (*Crotaphytus collaris*). *Univ. Kans. Publ. Mus. Nat. Hist.* 8:213-274.
- Johnson, D. H., M. D. Bryant, and A. H. Miller. 1948. Vertebrate animals of the Providence Mountains area of California. *Univ. Calif. Publ. Zool.* 48:221-376.
- Legler, J. and H. S. Fitch. 1957. Observations on hibernation and nests of the collared lizard, *Crotaphytus collaris*. *Copeia* 1957:305.
- McGuire, J. A. 1996. Phylogenetic systematics of crotaphytid lizards (Reptilia: Iguania: Crotaphytidae). *Bulletin of Carnegie Museum of Natural History* 32:1-120.
- Sanborn, S. R. and R. B. Loomis. 1979. Systematics and behavior of collared lizards (*Crotaphytus collaris*) in southern California. *Herpetologica* 35:101-106.
- Stebbins, R. C. 1985. A field guide to western reptiles and amphibians. 2nd ed., revised. Houghton Mifflin, Boston. 336pp.
- Yedlin, I. and G. Ferguson. 1973. Variations in aggressiveness of free-living male and female collared lizards, *Crotaphytus collaris*. *Herpetologica* 29:268-275.

R017

---

Life history accounts for species in the California Wildlife Habitat Relationships (CWHR) System were originally published in: Zeiner, D.C., W.F.Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. *California's Wildlife*. Vol. I-III. California Department of Fish and Game, Sacramento, California. Updates are noted in accounts that have been added or edited since original publication.