

California Wildlife Habitat Relationships System
California Department of Fish and Wildlife
California Interagency Wildlife Task Group

SIDEWINDER

Crotalus cerastes

Family: VIPERIDAE
R075

Order: SQUAMATA

Class: REPTILIA

Written by: R. Marlow
Reviewed by: T. Papenfuss
Edited by: S. Granholm

DISTRIBUTION, ABUNDANCE, AND SEASONALITY

The sidewinder is widely distributed and locally abundant in the Colorado and Mojave deserts and north just into the southern Great Basin. It is found from sea level to 1800 m (6000 ft) in a wide variety of desert habitats, including woodlands, shrub types, barren areas, and dunes. It is most abundant in desert washes and flats with shrub cover and wind-blown sand, especially at the bases of bushes where the snakes often wait, partially buried. This snake is active from mid-spring to early fall, but activity may be restricted during the hottest part of the summer (Cowles 1945, Klauber 1944, 1972, Lowe and Norris 1950, Stebbins 1954).

SPECIFIC HABITAT REQUIREMENTS

Feeding: This snake eats small mammals (*Dipodomys*, *Perognathus*, etc.), lizards (*Cnemidophorus*, *Uta*, *Uma*, *Phrynosoma*, etc.) and occasionally birds (Stebbins 1954, Klauber 1972). It is an active forager, but it also waits under bushes for prey, partially buried in sand.

Cover: This snake has a set of behavioral and morphological specializations for living in sand. It normally buries itself in a coil in fine sand at the base of a bush, if available, or in the open. It may occasionally use mammal burrows or surface cover objects.

Reproduction: Young are live-born and probably require a safe and secure place for birth. Rodent burrows are probably utilized.

Water: Probably not required.

Pattern: This snake occurs in a variety of habitats that are characterized by the presence of wind-blown sand. Rarely, this species occurs in rocky or non-sandy habitats.

SPECIES LIFE HISTORY

Activity Patterns: This snake is primarily nocturnal, but in the early spring it is active at dusk and even occasionally during the day. It is active from early to mid-spring until late summer or early fall. Populations of southerly or warmer areas become active earlier. It sometimes ceases activity in mid-summer, when temperatures are highest (Stebbins 1954, Klauber 1972).

Seasonal Movements/Migration: Not known to migrate.

Home Range: No data.

Territory: Males may be territorial but evidence is not clear (Lowe and Norris 1950).

Reproduction: Copulation occurs shortly after emergence in April and May and the young are born in mid-summer. Litters average 11 young and range from 7-13 (Stebbins 1954, Klauber 1972).

Niche: This snake is eaten by kingsnakes and roadrunners, and probably by most avian and mammalian desert predators.

REFERENCES

- Cowles, R. B. 1945. Some of the activities of the sidewinder. *Copeia* 1945:220-222.
- Klauber L. M. 1944. The sidewinder, *Crotalus cerastes*, with description of a new subspecies. *Trans. San Diego Soc. Nat. Hist.* 10:91-126.
- Klauber, L. M. 1972. Rattlesnakes: their habits, life histories, and influence on mankind. 2nd ed. Univ. California Press, Berkeley. 1533pp.
- Lowe, C. H., and K. S. Norris. 1950. Aggressive behavior in male sidewinders, *Crotalus cerastes*, with a Discussion of aggressive behavior and territoriality in snakes. *Chicago Acad. Sci. Nat. Hist. Misc. Publ.* 66:1-13.
- Stebbins, R. C. 1954. *Amphibians and reptiles of western North America.* McGraw-Hill, New York. 536pp.
- Stebbins, R. C. 1954. *Amphibians and reptiles of western North America.* McGraw-Hill, New York. 536pp.

R075

Life history accounts for species in the California Wildlife Habitat Relationships (CWHHR) System were originally published in: Zeiner, D.C., W.F.Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. *California's Wildlife. Vol. I-III.* California Department of Fish and Game, Sacramento, California. Updates are noted in accounts that have been added or edited since original publication.