

Literature Cited

- Alstatt, J.M., Ambrose, R.F., Engle, J.M., Haaker, P.L., Lafferty, K.D., and Raimondi, P.T. 1996. Recent declines of black abalone *Haliotis cracherodii* on the mainland coast of central California. Mar. Ecol. Prog. Ser. 142: 185-192.
- Ault JS, DeMartini JD. 1987. Movement and dispersion of red abalone, *Haliotis rufescens*, in Northern California. Calif. Fish and Game 73(4):196-213.
- Bonnot, P. 1948. The abalones of California. Calif. Fish and Game 34:140-169.
- Burton, R. S., and M. J. Tegner. 2000. Enhancement of Red Abalone, *Haliotis rufescens* stocks at San Miguel Island: reassessing a success story. Marine Ecology Progress Series. 202: 303-308.
- California Department of Fish and Game. 1991. Final supplemental environmental document abalone sport fishing. Sacramento, California: the Resources Agency.
- California Department of Fish and Game. 1993a. Final environmental document black abalone closure. Sacramento, California: the Resources Agency California.
- California Department of Fish and Game. 1993b. Final supplemental environmental document abalone ocean sport fishing. Sacramento, California: the Resources Agency.
- California Department of Fish and Game. 1995a. The red abalone (*Haliotis rufescens*) sport fishery in central and northern California from creel 1975-94, aerial 1975-1985, and telephone based survey. Sacramento, California: the Resources Agency.
- California Department of Fish and Game. 1995b. Draft environmental document pink, green and white abalone fishery closure. Sacramento, California: the Resources Agency.
- California Department of Fish and Game. 2001b. Draft supplement to the environmental document - ocean sport fishing regulations concerning abalone sport fishing. Sacramento, California: the Resources Agency.
- California Department of Parks and Recreation. 1988. Five views: an ethnic historic site survey for California. Office of Historic Preservation. Sacramento, California - The Resources Agency.
- Carr M, Syme C, and Caselle J. 2001. MLMA nearshore reef monitoring network (NRMN) proposal. Draft 10-20-01.
- Cicin-Sain, B., Moore, J.E., and Wyner, A.J. . 1977. Management approaches for marine fisheries: The case of the California abalone. Univ. Calif. Sea Grant College Program, Sea Grant Pub.54, 223 pp.
- Clavier, J. & Richard, O. 1986. Growth of juvenile *Haliotis tuberculata* (Mollusca: Gastropoda) in the natural environment. J. Mar. Biol. Assoc. UK 66:497-503.
- Conte, F.S. and S. McBride. 1996. California abalone aquaculture: California Aquaculture. ASAQ-A10-6/96. 5 pp.
- Cooper, J., M. Wieland, and A. Hines. 1977. Subtidal abalone populations in an area inhabited by sea otters. Veliger 20:163-167.
- Cox KW. 1960. Review of the abalone in California. Calif. Fish and Game 46:381-406.

- Cox KW . 1962. California abalones, family Haliotidae. Calif. Department of Fish and Game. Fish Bulletin 118. 133 p.
- Crisp, DS. 1974. Factors affecting settlement of marine invertebrate larvae. In Chemoreception in marine Organisms (ed) by P. Grant A.M. Mackie pp. 177-265. Academic press New York and London.
- Culver CS, Kuris, AM. 2000. The apparent eradication of a locally established introduced marine pest. Biological Invasions 2: 245-253
- Daniels R, Floren R. 1998. Poaching pressures on Northern California's Abalone Fishery. Journal of Shellfish Research. 17(3): 859-862.
- Daume S., Brokerand-Gardner S. & Woerkerling Wm.J. 1999. Preferential settlement of abalone larvae: diatom films vs. non geniculate coralline red algae. Aquaculture 174, 243-254.
- Davis GE. 1988. Kelp forest monitoring handbook. National Park Service.
- Davis GE. 1989a. Design of a long-term ecological monitoring program for the Channel Islands National Park, California. Natural Areas Journal. 9(2):80-89.
- Davis GE. 1993. Mysterious Demise of the southern California black abalone, *Haliotis cracherodii* Leach 1814. Journal of Shellfish Research 12(2): 183-184.
- Davis GE, Richards DV, Haaker PL, Parker DO. 1992. Abalone populations declines and fishery management in Southern California. In: Shepherd, Tegner, and Guzman del Proo, eds. Abalone of the World: Biology, Fisheries and Culture. Blackwell Scientific, Oxford, U.K. p. 237-249.
- Davis GE, Haaker PL, Richards DV. 1998. The perilous condition of white abalone, *Haliotis sorenseni*, Bartsch, 1940. Journal of Shellfish Research 17(3): 871-875.
- Day EG. 1998. Ecological interactions between abalone (*Haliotis midae*) juveniles and sea urchins (*Parechinus angulosus*) off the south-west coast of South Africa. [PhD dissertation]. University of Cape Town, South Africa. 174 p.
- Day, E.G. and G.M. Branch. 2002. Influences of the sea urchin *Parechinus angulosus* (Leske) on the feeding behavior and activity rhythms of juveniles of the South African abalone *Haliotis midae* Linn. J. Exp. Marine Biology and Ecology, 276(1-2):1-17.
- Dayton, P.K. and M.J. Tegner. 1984. Catastrophic storms, El Niño, and patch stability in a Southern California kelp community. Science 224:283
- Dayton P.K., and M.J. Tegner. 1989. Bottoms beneath troubled waters: Benthic impacts of the 1982-1984 El Niño in the temperate zone. Pages 433-472 in: P.W. Glynn (ed). Global ecological consequences of the 1982-1983 El Niño Southern Oscillation. Elsevier Oceanography Series, No. 52. Amsterdam, The Netherlands.
- Deacon, J. 1977. Habitat selection and competition among abalone and sea urchins at The Sea Ranch, California. MS Thesis. University of California, Davis. 83 pp.
- Dean, T.A., S.C. Schroeter, and J.D. Dixon. 1984. Effects of grazing by two species of sea urchins (*Strongylocentrotus franciscanus* and *Lytechinus anamesus*) on recruitment and survival of two species of kelp.
- Donovan, D.A. and T.H. Carefoot. 1997. Locomotion in the abalone *Haliotis kamtschatkana*: pedal morphology and cost of transport. The J. of Exper. Biol. 200: 1145-1153.

- Dugan, J.E. and G.E. Davis. 1993. Applications of marine refugia to coastal fisheries management. Can. J. Fish. Aquat. Sci. 50:2029-2042.
- Duggins, D.O. 1980. Kelp beds and sea otters: An experimental approach. Ecology 61:447-453.
- Ebeling, A.W., Laur, D.R., and Rowley, R.J. 1985. Severe storm disturbances and reversal of community structure in a southern California kelp forest. Marine biology, 84(3):287-294.
- Ebert, E.E., 1968. A food habits study of the southern sea otter, *Enhydra lutris nereis*. Calif. Fish and Game, 54(1):33-42.
- Ebert TB, EE Ebert 1988. An innovative technique for seeding abalone and preliminary result of laboratory and field trials. California Fish and Game, 74(2):68-81.
- Ebert, T.A., Dixon, J.D., Schroeter, S.C., Kalvass, P.E., Richmond, N.T., Bradbury, W.A. and D.A. Woodby. 1999. Growth and mortality of red sea urchin *Strongylocentrotus franciscanus* across a latitudinal gradient. Mar. Ecol. Prog. Ser. 190:189-209.
- Ebert, TA, MP Russell.1988. Latitudinal variation in size structure of the west coast purple sea urchin: A correlation with headlands. Limnol. Oceanogr. 33(2):286-294.
- Edwards, C.L. 1913. The abalone industry in California. Calif. Dept. of Fish and Game, Fish Bull. 1:5-15.
- Estes, J.A., and J.F. Palmisano. 1974. Sea otters: Their role in structuring nearshore communities. Science 185:1058-1060.
- Forward, R.G. JR., Wellins, C.A. & Buswell C.U. 1989. Behavioural responses of larvae of the crab *Neopanope sayi* to hydrostatic pressure. Mar. Ecol. Prog. Ser. 57:267-277.
- Foster, M.S., and D.R. Schiel. 1985. The ecology of giant kelp forests in California: A community profile. U.S. Fish Wildl. Serv. Biol. Rep. 85(7.2). 152 pp.
- Friedman CS, Thomson M, Chun C, Haaker PL, Hedrick RP. 1997. Withering Syndrome of the black abalone, *Haliotis cracherodii* (Leach): water, temperature, food availability, and parasites as possible causes. Journal of Shellfish Research 16(2):403-411.
- Friedman CS, Biggs W, Shields JD, Hedrick RP. 2002. Transmission of withering syndrome in black abalone, *Haliotis cracherodii* (Leach). Journal of Shellfish Research. Forthcoming.
- Friedman CS, Andree KB, Beauchamp K, Moore JD, Robbins TT, Shields JD, Hedrick RP. 2000. "Candidatus Xenohaliotis californiensis", a newly described pathogen of abalone, *Haliotis* spp., along the west coast of North America. International Journal of Systematic and Evolutionary Microbiology 50:847-855.
- Gaffney, P.M., V.P. Rubin, D. Hedgecock, D. Rowers, G. Morris, and L. Hereford. 1996. Genetic effects of artificial propagation: signals from wild hatchery populations of red abalone in California. Aquaculture. 143:257-266.
- Garland, C.D. Cooke, S.L. Grant, J.F. & McMeekin T.A. 1985. Ingestion of bacteria on the cuticle of crustose (non-articulated) coralline algae by post-larval juvenile abalone (*Haliotis rubra* Leach) from Tasmanian waters J. Exp. Mar. Biol. Ecol. 91:137-149.
- Geibel, J.J., and D.J. Miller. 1984. Estimation of sea otter , *Enhydra lutris*, population, with confidence bounds, from air and ground counts. Calif. Fish and Game 70:225-233.

- Geiger, D. 2000. Distribution and biogeography of the recent Haliotidae (Gastropoda: Vetigastropoda) world-wide. International Journal of Malacology 35(5-12):57-120.
- Gerard, V.A. 1976. Some aspects of material dynamics and energy flow in a kelp forest in Monterey Bay, California. PhD. Dissertation, University of California, Santa Cruz. 176pp.
- Gotshall,D.W., R.N. Lea, L.L. Laurent, T.L. Hoban and G.D. Farrens. 1974. Mendocino Power Plant site ecological study Final Report. PG and E Coop. Res. Agree S-1902. Marine Resources CDFG Admin. Report. 74-7.
- Giorgi, A.E. and DeMartini, J.D. 1977. A study of the reproductive biology of the red abalone (*Haliotis rufescens*) near Mendocino, California. Calif. Fish and Game 63:80-94.
- Gulland, J.A. 1983. Fish Stock Assessment: a Manual of Basic Methods. John Wiley and Sons, Chichester, U.K. 223 pp.
- Guzman-Del Proo, S. A. 1992. A review of the biology of abalone and its fishery in Mexico. In: S. A. Shepherd, M. J. Tegner, and S.A. Guzman del Proo (eds.). Abalone of the World: Biology, Fisheries and Culture. Blackwell Scientific, Oxford, U.K. pp. 341-360.
- Haaker, P.L., D.V. Richards, C.S. Friedman, G.E. Davis, D. Parker, and H.A. Togstad. 1992. Mass mortality and withering syndrome in Black Abalone, *Haliotis cracherodii*. In: Abalone of the World: Biology, Fisheries and Culture. S.A. Shepherd, M.J. Tegner, and S. Guzman del Proo. (eds.) Blackwell Scientific.
- Haaker PL, Parker DO, Chun CS. 1995. Growth of black abalone, *Haliotis cracherodii* (Leach) at San Miguel Island and Point Arguello, California. Journal of Shellfish Research 14(2):519-525.
- Haaker, P.L., Parker, K, Barsky, C., and Chun, C.S. 1998. Growth of red abalone, *Haliotis rufescens* (Swainson) at Johnsons Lee, Santa Rosa Island, California. Journal of Shellfish Research. 17(3):747-753.
- Hahn, K.O. 1989. Biotic and abiotic factors affecting the culture of abalone. Pages 113-134 in: K.O. Hahn (ed). The culture of abalone and other marine gastropods. CRC Press, Boca Raton, FL.
- Hamm, D.E. and R.S. Burton. 2000. Population genetics of black abalone, *Haliotis cracherodii*, along the central California coast. J. of Exper. Mar. Biol. and Ecol. 254:235-247.
- Hannan, C.A. 1984. Planktonic larvae may act like passive particles in turbulent near-bottom flows. Limnol. Oceanogr. 29:1108-1116.
- Hansen, J.D. 1970. Commensal activity as a function of age in two species of California abalones (Mollusca: Gastropoda). Veliger 13(1):90-94.
- Harrold, C. and D.C. Reed. 1985. Food availability, sea urchin grazing and kelp forest community structure.
- Harrold, C. and J.S. Pearse. 1987. The ecological role of echinoderms in kelp forests. Pp. 137-233. In: M. Jangoux and J.M. Lawrence (eds.) Echinoderm Studies 2. AA. Balkema, Rotterdam, The Netherlands.
- Henderson, K.C., D.O. Parker, and P.L. Haaker. 1988. The survival and growth of transplanted adult pink abalone, *Haliotis corrugata*, at Santa Catalina Island. Calif. Fish and Game 74:82-86.
- Hilborn R, Walters CJ. 1992. Quantitative fisheries stock assessment: choice, dynamics and uncertainty. Chapman and Hall. 570 p.

- Hines AH, Pearse JS. 1982. Abalone, shells, and sea otters: dynamics of prey populations in Central California. *Ecology* 63(5):1547-1560.
- Hobday AJ, Tegner MJ. 2000. Status review of white abalone (*Haliotis sorenseni*) throughout its range in California and Mexico. NOAA Technical Memorandum NOAA-TM-NMFS-SWR-035. U.S. Department of Commerce, National Oceanic and Atmospheric Administration. National Marine Fisheries Service - Southwest Region Office.
- Hutchings, J.A. and Myers, R.A. 1994. What can be learned from the collapse of a renewable resource? Atlantic cod *Gadus morhua*, of Newfoundland and Labrador. *Can J. Fish. Aquat. Sci.* 51: 2126-2146.
- Karpov KA. 1991. A combined telephone and creel survey of the red abalone, *Haliotis rufescens*, (Swainson), sport fishery in California from Monterey to the Oregon border. April through November 1989. California Department of Fish and Game, Marine Resources Division, Admin. Rep. No. 91-2. 72 p.
- Karpov KA, and Tegner MJ. 1992. Abalones. In: California's Living Marine Resources and Their Utilization. Leet WS, Dewees CM, Haugen CW (eds). Sea Grant Extension Publication UCSGEP-92-12. p33-36.
- Karpov KA, Geibel JJ, Law PM. 1997. Relative abundance and size composition of subtidal abalone, *Haliotis* spp., sea urchin, *Strongylocentrotus* spp., and abundance of sea stars off Fitzgerald Marine Reserve, September 1993. CDFG Mar. Res. Div. Admin. Rpt. 97-1. 28 p.
- Karpov KA, Haaker P, Albin D, Taniguchi IK, and Kushner D. 1998. The red abalone, *Haliotis rufescens*, in California: Importance of depth refuge to abalone management. *J. Shel. Res.* 17(3):863-870.
- Karpov KA, Haaker PL, Taniguchi IK, and Rogers-Bennett L. 2000. Serial depletion and the collapse of the California abalone (*Haliotis* spp.) fishery. In: Campbell A. (ed.). Workshop on rebuilding abalone stocks in British Columbia. *Can. Spec. Publ. Fish. Aquat. Sci.* 130. p 11-24.
- Karpov KA, Tegner MJ, Rogers-Bennett L , Kalvass PE, and Taniguchi IK. 2001. Interactions among red abalones and sea urchins in fished and reserve sites of northern California: implications of competition to management. *J. Shellfish Res.* 20(2):743-753.
- Kenyon, K.W. 1969. The sea otter in the eastern Pacific Ocean. North American Fauna. No. 68. Bureau of Sport Fisheries and Wildlife. U.S. Govt. Printing Office. Washington, D.C. 352 pp.
- Keough, M.J., and G.P. Quinn. 1998. Effects of periodic disturbance from trampling on rocky intertidal algal beds. *Ecolog. Appl.* 8: 141-161.
- Kirby, V. L., R. Villa, and D. A. Powers. 1998. Identification of microsatellites in the California Red Abalone, *Haliotis rufescens*. *J. of Shell. Res.* 17(3): 801-804.
- Kitting, K.L., and D.E. Morse. 1997. Feeding effects of postlarval Red Abalone, *Haliotis rufescens* (Mollusca:Gastropoda) on encrusting coralline algae. *Moll. Res.* 18:183-196.
- Kojima, H. 1981. Mortality of young Japanese black abalone *Haliotis discus* after translocation. *Bull. Of Jap. Soc. Of Sci. Fish.* 47(2):151-159.
- Lapota, D., G. Rosen, J. Chock, C.H. Liu. 2000. Red and green abalone seed growout for reseeding activities off Point Loma, California. *J. Shellfish Res.* 19(1):431-438.
- Lafferty KD, and Kuris AM. 1993. Mass mortality of abalone, *Haliotis cracherodii*, on the California Channel Islands: Tests of epidemiological hypotheses. *Mar. Ecol. Prog. Ser.* 96:239-248.

Leet, W.S., DeWees, C.M., Klingbeil, R. and E.J. Larson. 2001. California's Living Marine Resources: A Status Report. University of California. 592 pp.

Leighton, D.L. 1959. Diet and its relation to growth in the black abalone, *Haliotis cracherodii* (Leach). Masters Thesis, Univ. California, Los Angeles, 61 pp.

Leighton, D.L. 1966. Studies of food preference in algivorous invertebrates of southern California kelp beds. Pac. Sci. 20:104-113.

Leighton, D.L., 1968. A comparative study of food selection and nutrition in the abalone, *Haliotis rufescens* (Swainson), and the sea urchin, *Strongylocentrotus purpuratus* (Stimpson). Ph.D. Thesis, Univ. California, San Diego, 197pp.

Leighton, D.L. 1972. Laboratory observations on the early growth on the abalone, *Haliotis sorenseni*, and the effect of temperature on larval development and settling success. Fish. Bull., U.S. 70:373-381.

Leighton DL. 1974. The influence of temperature on larval and juvenile growth in three species of southern California abalones. Fish. Bull. 72(4):1137-1145.

Leighton, D. L. and R. A. Boolootian. 1963. Diet and growth in the black abalone, *Haliotis cracherodii*. Ecology 44:227-238.

Levitin, D.R., M.A. Sewell, and F.S. Chia. 1992. How distribution and abundance influence fertilization success in the sea urchin, *Strongylocentrotus franciscanus*. Ecology 73:248-254.

Lowry, L.F. and J.S. Pearse. 1973. Abalones and Sea Urchins in a Sea Otter Habitat. University Of California, Santa Cruz. 15 pp.

MacCall, A.D. 1989. Against marine fish hatcheries: Ironies of fishery politics in the technological era. Calif. Coop. Ocean Fish. Inv. Rep. 30: 46-48.

MacGinitie, G.E., and N. MacGinitie. 1949. Natural history of marine animals. McGraw Hill Book Co., New York, NY. 473 pp.

McMillen, R. and J. Phillips. 1974. In search of the elusive *Haliotis walallensis*. Conchologists Amer. Bull. 1:2.

McMullen, J., and T. Thompson. 1989. Abalone culture in an urban environment. Pages 227-238 In: Hahn, K.O. (ed). The culture of abalone and other marine gastropods. CRC Press, Boca Raton, FL.

McShane, P.E. 1992. Early life history of abalone: a review. Pages 120-138. In: Abalone of the World: Biology, Fisheries and Culture. S.A. Shepherd, M.J. Tegner, and S. Guzmán del Próo (eds). Abalone of the world: Biology, fisheries and culture. Blackwell Scientific, Oxford, U.K.

Miller DJ, Geibel JJ, Houk JL. 1974. Results of the 1972 skindiving assessment survey; Pismo Beach to Oregon. Calif. Dept. Fish and Game. Technical Report 23:61.

Moore JD, Robbins TT, Friedman CS. 2000. Withering syndrome in farmed red abalone, *Haliotis rufescens*: Thermal induction and association with a gastrointestinal Rickettsiales-like prokaryote. Journal of Aquatic Animal Health. 12:26-34.

Moore JD, Robbins TT, Hedrick R.P., and Friedman CS. 2001. Transmission of the Rickettsiales-like prokaryote "Candidatus Xenohaliotis californiensis" and its role in withering syndrome of California abalone, *Haliotis* spp. J. Shellfish Res. 20:867-874.

- Morse, D.E., N. Hooker, L. Jensen, and H. Duncan. 1979. Induction of larval abalone settling and metamorphosis by gamma-aminobutyric acid and its congeners from crustose red algae. II. Applications to cultivation, seed production and bioassays; principal causes of mortality and interference. *J. World Maricul. Soc.* 10:81-91.
- Murray SN, and TG Denis. 1997. Vulnerability of the rockweed *Pelvetia compressa* to anthropogenic disturbance on southern California rocky shores. *Phycologia* 36:75–76.
- North, W.J. and J.S. Pearse. 1970. Sea urchin population explosion in southern California coastal waters. *Science* 167:209.
- North, W.J. 1971. The biology of giant kelp beds (*Macrocystis*) in California.. *Nova Hedwigia* 32:1-660.
- O'Connell, C. 1955. The life history of the cabezon *Scorpaenichthys marmoratus* (Ayres). Calif. Department of Fish and Game. Fish Bulletin 93. 76 pp.
- Orensanz, J.M.L., Armstrong, J., Armstrong, D., Hilborn, R. 1998 Crustacean resources are vulnerable to serial depletion - the multifaceted decline of crab and shrimp fisheries in the Greater Gulf of Alaska. *Reviews in Fish Bio. and Fish.* 8(2):117-176.
- Parker DO, Haaker PL, Henderson KC. 1988. Densities and size composition of red abalone, *Haliotis rufescens*, at five locations on the Mendocino and Sonoma county coasts, September 1986. Calif. Dept. Fish and Game, Mar. Res. Div., Admin. Rep. No. 88-5, 65 p.
- Parker, D.O., P.L. Haaker and H.A. Togstad. 1992. Case histories for three species of California abalone, *Haliotis corrugata*, *H. fulgens* and *H. cracherodii*. Pages 384-394. In: S.A. Shepherd, M.J. Tegner, and S. Guzmán del Próo (eds). *Abalone of the world: Biology, fisheries and culture*. Blackwell Scientific, Oxford, U.K. *Abalones of the World*.
- Pennington, J.T. 1985. The ecology of fertilization of echinoid eggs: the consequences of sperm dilution, adult aggregation, and synchronous spawning. *Biol. Bull.* 169:417-430.
- Pilson, M.F.O., and M.B. Taylor. 1961. Hole drilling by octopus. *Science*. 134: no. 3487, 1366-1368.
- Post JR, Sullivan M, Cox S, Lester NP, Walters CJ, Parkinson EA, Paul AJ, Jackson L, Shuter BJ. 2002. Canada's recreational fisheries: the invisible collapse? *Fisheries* 27:6-17.
- Prince, J.D., T.L. Sellers, W.B. Ford and S.R. Talbot. 1987. Experimental evidence for limited dispersal of haliotid larvae (genus *Haliotis*; Mollusca: Gastropoda) *J. Exp. Mar. Ecol.*, 106:243-263.
- Raimondi PT, Wilson CM, Ambrose RF, Engle JM, Minchinton TE. 2002. Continued declines of black abalone along the coast of California: are mass mortalities related to El Nino events? *Marine Ecology Progress Series*, 242: 143-152
- Ricker, W. E. 1975. Computation and interpretation of biological statistics of fish populations. *Bull. Fish. Res. Board Can.* 191:1-382.
- Rogers-Bennett L, and Pearse JS. 1998. Experimental seeding of hatchery-reared juvenile red abalone in northern California. *J. Shellfish Res.* 17(3):877-880.
- Rogers-Bennett, L. and J. Pearse. 2001. Indirect benefits of marine protected areas for juvenile abalone. *Conserv. Biol.* 15:1-6.
- Rogers-Bennett, L., P.K. Haaker, T.O. Huff, and P.K. Dayton. 2002. Estimating Historic Abundances of Abalone in California for Restoration. *CalCOFI* 43:63-74.

- Saito, K. 1981. The appearance of 0-year old Ezo abalone. Bull. Jap. Soc. Fish., 47:1393-1401.
- Saito, K. 1984. Ocean ranching of abalones and scallops in northern Japan. Aquaculture, vol. 39, no. 1-4, pp. 361-373
- Shepherd, S.A. 1973. Competition between sea urchins and abalone. Australian Fisheries 6(32).
- Shepherd SA, and Brown LD. 1993. What is an abalone stock: Implications for the role of refugia in conservation. Can. Jour. Fish. and Aqua. Sci. 50:2001-2009.
- Shepherd S.A. and J.A. Turner 1985. Studies on southern Australian abalone (genus *Haliotis*) VI. Habitat preference and abundance and predators of juveniles. J. Exp. Mar. Biol. Ecol. 93: 285-298.
- Siqueiros-Beltrones, D.A. and D. Voltolina 2000. Grazing Selectivity of Red Abalone, *Haliotis rufescens*, Postlarvae on Benthic Diatom Films under Culture Conditions. J. of Wld. Aquacul. Soc. 31(2):239-246.
- Sluzanowski, P.R., 1984. A management oriented model of an abalone fishery whose substocks are subject to pulse fishing. Can.J. Fish. Aquat. Sci., 1:1008-1014.
- Smith, J.R. 2002. The effects of bait collection and trampling on *Mytilus californianus* communities in the southern California intertidal zone. MS Thesis, Department of Biological Sciences, California State University, Fullerton
- Tarr, R.J.Q., Williams, P.V.G., and A.J. MacKenzie. 1996. Abalone, sea urchins and rock lobster: a possible ecological shift that may affect traditional fisheries. S. African J. of Marine Science. 17:319-323.
- Tegner MJ, 1980. Multispecies considerations of resource management in southern California kelp beds. Can. Tech. Rep. Fish. Aquat. Sci. 954:125-143.
- Tegner MJ. 1989a. The feasibility of enhancing red sea urchin *Strongylocentrotus franciscanus* stocks in California; an analysis of the options. Mar. Fish. Rev. 51(2):1-22.
- Tegner MJ. 1989b. The California abalone fishery: production, ecological interactions, and prospects for the future. In: Caddy JF. (ed). Marine Invertebrate Fisheries: Their Assessment and Management. John Wiley and Sons, New York. p 401-420.
- Tegner MJ. 1992. Brood-stock transplants as an approach to stock enhancement. In: Shepherd SA, Tegner MJ, Guzman del Proo S. (eds). Abalone of the world: Biology, fisheries and culture. Blackwell Scientific, Oxford, U.K. p 461-473.
- Tegner, M.J. 1993. Southern California abalones: Can stocks be rebuilt using marine harvest refugia? Can. J. Fish. Aquat. Sci. 50:2010-2018.
- Tegner, M.J. and P.K. Dayton. 1977. Sea urchin recruitment patterns and implications of commercial fishing. Science 196:324-326.
- Tegner MJ, and Butler RA. 1985a. The survival and mortality of seeded and native red abalones, *Haliotis rufescens*, on the Palos Verdes Peninsula. Calif. Fish and Game 71(3):150-163.
- Tegner MJ, and Butler RA. 1989. Abalone seeding. In: Hahn K (ed). Handbook of culture of abalones and other gastropods. CRC Press, Boca Raton, Fl. p 157-182.
- Tegner MJ, Dayton PK. 2000. Ecosystem effects of fishing in kelp forest communities. ICES. J. Mar. Sci. 57:579-589.

- Tegner, M.J. and L.L. Levin. 1982. Do sea urchins and abalones compete in the California kelp forest communities? Pages 265-271 In: J.W. Lawrence (Ed.), International Echinoderms Conference, Tampa Bay. A.A. Bakema, Rotterdam.
- Tegner MJ, Breen PA, and Lennert CE. 1989. Population biology of red abalone, *Haliotis rufescens*, in Southern California and management of the red and pink, *H. corrugata*, abalone fisheries. California Department of Fish and Game. Fish Bulletin 87:313-339.
- Tegner MJ, DeMartini JD, and Karpov KA. 1992. The California red abalone fishery; a case study in complexity. In: Shepherd SA, Tegner MJ, Guzman del Proo S. (eds). Abalone of the world: Biology, fisheries and culture. Blackwell Scientific, Oxford, U.K. p 370-383.
- Tegner, M.J., Basch, L.V., and P.K. Dayton. 1996. Near extinction of an exploited marine invertebrate. Trends Ecol. Evol. 11:278-280.
- Tegner, M.J. P.L. Haaker, K.L. Riser, and L. Ignacio Vilchis. 2001. Climate variability, kelp forests, and the southern California red abalone fishery. J. Shellfish Res. 20: 755-764.
- Tutschulte, T.C. 1976. The comparative ecology of three sympatric abalones. Ph.D. dissertation. University of California, San Diego, CA.
- Tutschulte, T.C. and J.H. Connell. 1988b. Growth of three species of abalones (*Haliotis*) in southern California. The Veliger 31 (3/4): 204- 213.
- U.S. Fish and Wildlife Service and Institute of Marine Sciences, University of California, Santa Cruz. 1987. Final environmental impact statement for translocation of sea otters. Volume I. May 1987. U.S. Fish and Wildlife Service, Sacramento, CA. 515 pp.
- U.S. Fish and Wildlife Service. 2000. Reinitiation of formal consultation on the containment program for the southern sea otter (1-8-99-FW-81). California/Nevada Operations Office. 19 July.
- U.S. Fish and Wildlife Service. 2003. Final Revised Recovery Plan for the Southern Sea Otter (*Enhydra lutris nereis*). Portland, Oregon. xi + 165 pp.
- Valiela, I. 1984. Marine ecological processes. Springer-Verlag, Berlin. 546 pp.
- VanBlaricom, G.R., and J.A. Estes (eds). 1988. The community ecology of sea otters. Springer Verlag, Berlin, Germany. 247 pp.
- Watson J. 2000. The effects of sea otters (*Enhydra lutris*) on abalone (*Haliotis* spp.) populations. In: Campbell A (ed). Workshop on rebuilding abalone stocks in British Columbia. Can. Spec. Publ. Fish. Aquat. Sci. 130:123-132.
- Watson, J. C. and T.G. Smith. 1996. The effects of sea otters on invertebrate fisheries in British Columbia: a review. In Invertebrate working papers. Reviewed by the Pacific Stock Assessment Review Committee (PSARC) in 1993 and 1994. Edited by C.M. Hand and B.J. Waddell. Can . Tech. Rep. Fish Aquat. Sci. 2089: 262-303.
- Wendell F. 1994. Relationship between sea otter range expansion and red abalone abundance and size distribution in Central California. California Fish and Game (80)2: 45-56.
- Wilson, K.C. and R. McPeak. 1983. Kelp restoration. Pages 199-216 In: W. Bascom (ed). The effects of waste disposal on kelp communities. Institute of Marine Research, University of California, San Diego, CA.

Woodby, D., Larson, R. and J. Rumble. 2000. Decline of the Alaska abalone (*Haliotis spp.*) Fishery and prospects for rebuilding the stock. In A. Campbell (ed.) Workshop on Rebuilding Abalone Stocks in British Columbia. Can. Spec. Publ. Fish. Aquat. Sci. 130:25-31.