

2010 SCOTT RIVER SALMON STUDIES

FINAL REPORT



Prepared By:

Morgan Knechtle and Diana Chesney

**California Department of Fish and Game
Northern Region
Klamath River Project
1625 South Main Street
Yreka, California 96097
(530) 842-3109**

SCOTT RIVER SALMON STUDIES, 2010

California Department of Fish and Game
Northern Region
Klamath River Project

ABSTRACT

The California Department of Fish and Game's (Department), Klamath River Project (KRP) operated a video fish counting facility and conducted cooperative spawning ground surveys (carcass surveys) on the Scott River during the 2010 fall-run Chinook salmon (*Oncorhynchus tshawytscha*) and coho salmon (*Oncorhynchus kisutch*) spawning season. The purpose of these surveys is to describe the run characteristics of adult fall-run Chinook salmon and coho salmon into the Scott River. Video fish counting operations began on September 29, 2010 and ended on January 7, 2011. The total number of Chinook salmon that entered the Scott River during the 2010 season is estimated to be **2,508** fish. Based on the proportion of male and female Chinook salmon that were sampled during the spawning ground surveys the run was comprised of approximately 923 (36.8%) males and 1,585 (63.2%) females. Based on a scales age analysis, adults comprised approximately 84.0% (2,107 fish) and grilse comprised 16.0% (401 fish) of the run. Males ranged in fork length (FL) from 42cm to 105cm and averaged 71.9cm. Females ranged in FL from 45cm to 96cm and averaged 80.2cm. KRP staff estimated that none of the Chinook salmon were of hatchery origin.

The first adult coho salmon was observed at the Scott River Fish Counting Facility on October 25, 2010 and the last coho salmon was observed on December 13, 2011. A total of 911 coho salmon were observed moving upstream through the Scott River Fish Counting Facility (SRFCF) during the season. Eight coho redds were estimated in areas below the counting facility representing an estimated 16 coho. The total number of coho salmon that entered the Scott River during the 2010 season is estimated to be **927**. Based on the proportion of male and female coho salmon that were sampled during the season the run was comprised of approximately 291 (31.4%) males and 636 (68.6%) females. Based on video data, adults comprised approximately 98.5% (913 fish) and grilse comprised 1.5% (14 fish) of the run. Males ranged in fork length (FL) from 59cm to 77cm and averaged 70.1cm. Females ranged in FL from 60cm to 75cm and averaged 69.1cm. None of the coho salmon were of estimated to be of hatchery origin.

INTRODUCTION

STUDY LOCATION AND RUN TIMING

The Scott River is a major tributary of the Klamath River located in Siskiyou County, and enters the Klamath River at river mile 143 (Figure 1). The Scott River fish counting facility is located at river mile 18.2 near the downstream edge of Scott Valley between the Indian Scotty Campground and Jones Beach picnic area (041° 38' 10.93" N; 123° 04' 3.08" W). Fall-run Chinook salmon typically return to the Scott River to spawn from mid September to late December. The coho salmon spawning run occurs from mid October to early January and steelhead run from November to March.




Figure 1. Location of the Scott River, tributary to the Klamath River, Siskiyou County.

KLAMATH RIVER PROJECT AND THE SCOTT RIVER STUDY

The Scott River study is one component of the KRP (initiated in 1978). The goals of the KRP include obtaining information on species composition, spawning distribution, FL frequency and sex ratios for salmonids, primarily fall-run Chinook salmon (Chinook), in various tributaries to the Klamath River including the Salmon, Scott, and Shasta rivers, as well as Bogus Creek and a dozen other smaller tributaries. The Scott River is particularly important because it is a major salmon spawning tributary. For example, during the 1996-98 spawning seasons, an average of 30.6% (8,914) of the total number of natural area Klamath River adult Chinook salmon spawners above the Trinity River confluence were estimated to have entered the Scott River to spawn. Therefore, a significant portion of natural escapement to the Klamath Basin would be unaccounted for if the Scott River studies were not conducted. In addition to providing valuable escapement estimates to the Pacific Fisheries Management Council for the effective management and allocation of fall Chinook salmon originating from the Klamath River Basin, the Scott River studies provide an opportunity to monitor an independent population within the state and federally listed Southern Oregon/Northern California Coast coho salmon (SONCC) range.

In the early years of the KRP, spawning ground surveys were conducted in the major spawning areas of the river which included about 5.5 miles of the Scott River near Etna and 4.75 miles of river downstream of the State Highway 3 Bridge crossing near Fort Jones. From 1989 through 1991 spawning ground surveys were limited to the lower river. In 1985 a temporary fish marking weir was installed on the lower river at river mile 1.6 and was operated during each spawning season until 1991. Operation of the

weir was often hampered by high flows, and beginning in 1992 operation of the weir was dropped in favor in conducting more intensive mark recapture spawning ground surveys in cooperation with USDA Forest Service (USFS) fisheries staff.

In 1994 the California State Legislature passed the Leslie Amendment (SB 779). The passage of SB 779 required Departmental staff to obtain landowner permission prior to accessing private lands to conduct biological investigations. The entire length of the Scott River within the Scott Valley (above river mile 24) passes through privately owned agricultural lands. As a result, since 1994, spawning ground surveys have been limited to those areas of the river on private land where landowner permission has been granted each year. The level of cooperation from local landowners has varied over the years. However, since the 2001 and 2002 spawning seasons, the number of landowners that have denied permission for access has increased dramatically. Controversies associated with the listing of SONCC coho salmon under the California Endangered Species Act (CESA) and other regulatory actions have substantially reduced the amount of cooperation provided by local landowners to the extent that the Department has been denied permission to survey nearly all of the Chinook spawning reaches present in the Scott Valley. As a result of the limited landowner access to the valley reaches the Department proposed installation of a fish counting facility to be located at the upper end of the canyon reach. The location of the fish counting station allows for monitoring fish abundance into the valley. The counting facility is also located upstream of several tributaries that can produce significant fall and winter stream flows, thereby reducing the probability of having the counting facility inoperable due to high flow events.

SCOTT RIVER STUDY OBJECTIVES SUMMARIZED:

- A) Determine the in-river run size (escapement) of Chinook and coho salmon returning to the Scott River.
- B) Determine run timing, spawning distribution, length frequency (FL) distribution, and sex ratio for Chinook and coho salmon in the Scott River.
- C) Collect scale samples and recover heads (containing coded wire tags) from ad-clipped Chinook in order to determine age composition and hatchery contribution rates of the run.
- D) Collect biological data for all steelhead observed during the Chinook and coho salmon spawning seasons.

METHODS

OPERATION OF THE SCOTT RIVER FISH COUNTING FACILITY

The video fish counting system was installed at the Scott River Fish Counting Facility on September 29, 2010 at 1153 hours Pacific Standard Time (PST). A temporary resistance board weir was installed to direct migrating fish into a flume where they pass in front of a video camera (Figure 2). The underwater video system consisted of a digital color video camera, water proof camera housing, viewing window, and counting flume which allowed for recording unimpeded fish passage through the facility. The facility was operated 24 hours a day, seven days a week throughout the Chinook and coho salmon migration period. A Splash Cam digital color video camera equipped with a 3.6mm wide angle lens with an auto iris was used to collect the photo image and an Ever Focus Digital Video Recorder (Model EDSR100) was used to record the image to external hard drives. The time lapse DVR was set to record continuously and drive changes were made at least twice a week.



Figure 2. Scott River Fish Counting Facility located in Siskiyou County, California 2010.

All hard drives were immediately returned to the office where each was subsequently downloaded and reviewed by project staff in the video lab. During each review staff recorded the date, time (hour:min:sec), and species of each fish observed on each video image. If the species could not be determined due to poor visibility or picture quality, staff recorded that observation as fish unknown. Staff also noted any ad-clipped fish observed, recorded the presence of lamprey scars and any other distinguishable marks that were visible on the tape. All data was then entered into computer files and each data file was subjected to one independent edit prior to commencement of data analysis.

SPAWNING GROUND SURVEYS

Spawning ground surveys were conducted twice a week on Mondays and Thursdays throughout the Chinook salmon spawning season starting October 11, 2010 and ending November 29, 2010. A total of twelve surveys were performed during the spawning season. On the morning of each survey, crews of at least two people were given daily instructions, data sheets, field equipment, vehicle assignments, and were assigned a survey reach. Crews walked their assigned reach in a downstream direction looking for salmon carcasses and spawning redds. All new redds were mapped once a week, recorded on USGS topographic maps, and the information was provided to the Klamath National Forest. All carcasses recovered were identified to species and gender, checked for marks or tags, measured (FL), a scale

sample was collected for age composition analysis, and females were examined for spawning success.

For purposes of the mark recapture estimate, each carcass was categorized into one of four pathways (Paths). Fresh carcasses, those with clear eyes and/or firm flesh were designated as Path 1. Individually numbered jaw tags were attached to the lower right jaw of all Path 1 carcasses and returned to the river for potential recapture during later surveys. Older carcasses, those with cloudy eyes and/or mushy flesh, were categorized as Path 2. All Path 2 carcasses were cut in half and returned to the river after all biological data was collected. Path 3 carcasses included all of the Path 1 carcasses (with jaw tag) that were recaptured during subsequent surveys. Any carcasses that could be observed by a survey crew but could not be retrieved for data collection, because they were located in inaccessible or unsafe locations, were designated as Path 4. Path 4 designations were rarely encountered during the survey.

The final Chinook salmon run-size estimate for reaches below the counting facility is normally calculated using the Schafer Method as follows (Ricker 1975):

$$\text{Escapement} = \sum((R_{ij}) (M_i/R_i)(C_j/R_j))$$

Where:

- M = The total number of fish marked
- M_i = The number of fish marked in period i
- R_i = The total number of marked fish recaptured in period i
- R_j = The total number of marked fish recaptured in period j
- C = The total number of fish recaptured during the season
- C_j = The total number of fish recaptured in period j

Unfortunately during 2010 extreme low numbers of carcasses were recaptured in the spawning ground survey below the counting facility which prevented the use of the Schafer mark recapture estimate. As a result the adult Chinook abundance below the weir was estimated by multiplying the total number of redds observed during the Chinook period by two. The grilse component from the redd survey was then added back into the total (total run= adults/(1-%jacks estimated((derived from aged scale samples)).

SURVEY REACHES

Survey reaches have remained fairly consistent since the beginning of the cooperative spawning ground survey in 1992. During the Chinook salmon spawning season, decisions regarding which reaches should be surveyed were based on the known distribution of the Chinook salmon run each week, the available labor force present during each survey, and on private lands was limited to those areas where permission has been granted by private landowners.

A total of 16 survey reaches, covering approximately 53.6 river miles, have been identified on the Scott River (Table 1, Figure 3). Access to private lands along the Scott River is critically important to the survey in those spawning areas that are present in Scott Valley. The highest observed densities of Chinook salmon spawning areas within Scott Valley are located downstream of the State Highway 3 Bridge crossing (rm 34.6) to the USGS gauging station located at river mile 21 (Reaches 8, 9, and 10), and in that reach of the river located downstream of Young's Dam, river mile 46, to about river mile 42 located upstream of the Eller Lane Bridge crossing (Reaches 12, 13, and 14).

Table 1. Description of cooperative spawning ground survey reach locations along the Scott River during the 2010 season.

Reach Number	Downstream Limit	RM	Upstream limit	RM	Length (miles)
1	Mouth	0.00	Mid Point	2.60	2.60
2	Mid Point	2.60	Pat Ford Ck	4.90	2.30
3	Pat Ford Ck	4.90	George Allen Gulch	7.80	2.90
4	George Allen Gulch	7.80	Townsend Gulch	10.50	2.70
5	Townsend Gulch	10.50	Bridge Flat	14.20	3.70
6	Bridge Flat	14.20	Counting Weir	18.20	4.00
7	Counting Weir	18.20	USGS Stream Gage	21.00	2.80
8	USGS Stream Gage	21.00	Meamber Bridge	24.40	3.40
9	Meamber Bridge	24.40	Dunlop	29.50	5.10
10	Dunlop	29.50	Highway 3 Bridge	35.60	6.10
11	Highway 3 Bridge	35.60	Eller Lane	41.10	5.50
12	Eller Lane	41.10	Sweezy Bridge	42.10	1.00
13	Sweezy Bridge	42.10	Horn Lane	43.90	1.80
14	Horn Lane	43.90	Young's Dam	46.00	2.10
15	Young's Dam	46.00	Fay Lane	49.60	3.60
16	Fay Lane	49.60	East Fork Confluence	53.60	4.00

To assist in developing stock identification baseline information the KRP collected both genetic tissue and otolith samples during the season. DNA samples were collected from 129 Chinook salmon (Tissue collected from the first five fish for each reach and each survey date). All samples were collected following protocols provided by the National Oceanic Atmospheric Administration's (NOAA) Southwest Fisheries Science Center. Samples were sent to the Salmonid Genetic Tissue Repository located at the NOAA Santa Cruz Laboratory for archiving and analysis. Otoliths were collected from 90 Chinook salmon (first fish from each reach from each survey date). All otoliths collected were archived for future microchemistry analysis. All otolith samples were collected following standard protocols described by Stevenson (1992).

POPULATION ESTIMATE

The Chinook salmon spawner escapement for the area of Scott River upstream of the counting facility was derived from a direct count of all Chinook salmon observed at the video counting facility. To estimate total escapement in the Scott River, the number of Chinook salmon redds multiplied by 2 (utilizing data from reach 1 through reach 6 only) for adults were added to the count of all Chinook salmon observed passing through the video counting facility. The grilse component from below the counting facility was also added to the total (total run= adults/(1-%jacks estimated)).




Figure 3. Location of the fish counting facility and spawning ground survey reaches on the Scott River used during the 2010 field season.

The coho salmon spawner escapement for the area of the Scott River upstream of the counting facility was also derived from a direct count of all coho salmon observed at the video counting facility. Spawning ground surveys were conducted on the mainstem Scott River below the weir through the end of November. Spawning ground surveys were conducted on coho spawning tributaries below the weir (Tompkins, Kelsey Creek and Canyon Creek) through the first week of January. To estimate total escapement in the Scott River, the number of coho salmon redds multiplied by 2 (utilizing data from tributary reaches and mainstem reaches below the weir only) for adults were added to the count of all coho salmon observed passing through the video counting facility. The grilse component from below the counting facility was then added back into the total (total run= adults/(1-%jacks estimated)).

RESULTS

OPERATION OF THE SCOTT RIVER FISH COUNTING FACILITY

The SRFCF began recording fish movements on September 29, 2010. The first Chinook salmon was observed at the SRFCF on September 30, 2010 and the last Chinook salmon was observed on November 30, 2010. The run peaked between October 24, 2009 and October 30, 2009 when 54.1% of the Chinook migration was observed (Figure 4). The majority of Chinook salmon passed through the SRFCF during daylight hours and peaked in the afternoon between 1200 and 1600 hours (Figure 5).




Figure 4. Run timing of Chinook salmon through the Scott River Fish Counting Facility during the 2010 season. Both upstream and downstream movements through the counting flume are shown (N=2,283).

A total of 2,149 Chinook salmon were estimated to have passed through the Scott River Fish Counting Facility during the 2010 season. Fourteen unidentified fish were assigned to be Chinook during the season. Any unidentified fish were assigned species composition based on the proportion of known daily observations. Sixty-five Chinook were added to the total as an expansion for periods of time when the camera was not functioning. During the Chinook period the only time the camera was not functioning was from 2030 hours on October 24, 2010 to 0900 on October 25, 2010 for a total of 12.5 hours.

SPAWNING GROUND SURVEYS

A total of 475 Chinook carcasses were sampled during the spawning ground survey as Path 1 or Path 2 carcasses. Of these 175 (36.8%) were male and 300 (63.2%) were female. Males ranged in FL from 42cm to 105cm and averaged 71.9cm (Figure 6). Females ranged in FL from 45cm to 96cm and averaged 80.2cm (Figure 7). One ad-clipped Chinook was observed during the spawning ground survey on November 15, 2010. The head collected from this ad-clipped Chinook did not contain a coded wire tag indicating that it most likely misclassified as an ad clip during the spawning ground survey. After examination of the length frequency distribution and scales age analysis of Path 1 and Path 2 carcasses, a grilse cut-off of < 63 cm was established for Scott River.




Figure 5. Summary of daily run timing of Chinook salmon observed at the Scott River Fish Counting Facility during 2010.

A total of 300 Chinook salmon female carcasses were observed during the spawning ground survey. Each female carcass was examined to determine if it had successfully spawned prior to death. Females with approximately 50% or more of their eggs still present in the body cavity when examined were identified as a pre-spawn mortality. Of the 300 female Chinook salmon carcasses examined, 272 females (90.7%) were found to have spawned, and 28 females (5.6%) still contained more than 50% of their spawn and were identified as pre-spawn mortalities. A total of 118 redds were observed in Reaches 1-6 yielding an estimated 236 adults (redds *2) and 44 grilse (total run= adults/(1-%jacks estimated)).

In 2010 the Schaffer mark recapture data generated from the spawning ground survey was segregated by reaches and analyzed independently for three areas: 1) all reaches, 2) reaches 7-8 and 3) reach 8 alone. In previous years the basin estimate was derived by adding the Schaffer estimate generated from reaches 1-6 to the total number of Chinook observed passing through the counting facility, but during 2010 the Schaffer estimate generated from reaches 1-6 was unusable due the lack of recaptures (Marked=43; Captured=80; Recaptured=3). The Schaffer estimate for all reaches, reaches 7-8, and reach 8 were 1,031; 549 and 378 respectively. For all reaches 493 Chinook were recovered, 343 were marked and 120 were subsequently recaptured yielding a recapture rate of 35.0%. For reach 7 and 8 together 253 Chinook were recovered, 186 were marked and 52 were recaptured yielding a recapture rate of 27.9%, for reach 8 alone, 209 Chinook were recovered, 159 were marked and 44 were recaptured yielding a recapture rate of 27.7%. The total Chinook salmon run size estimate (based on summing the video estimate and the redd estimate from below the weir) was estimated to be 2,508 fish. Based on scale age analysis, adults comprised approximately 84.0% (2,107 fish) and grilse comprised 16.0% (401 fish) of the run (KRTAT 2011).




Figure 6. Length Frequency distribution of Path 1 and Path 2 male Chinook salmon observed during spawning ground surveys in the Scott River, 2010 (n = 175).




Figure 7. Length frequency distribution of Path 1 and Path 2 female Chinook salmon observed during spawning ground surveys in the Scott River, 2010 (n = 300).

COHO SALMON

The first adult coho salmon was observed at the counting facility on October 25, 2010 and the last coho salmon was observed on December 13, 2010. A total of 878 coho salmon were observed moving

upstream through the SRFCF during the season (Figure 8). Coho salmon migration peaked during the 15 day period from October 25, 2010 to November 8, 2010 when 653 or 71.7% of the coho were observed. Four coho salmon were observed swimming downstream during the season. During the coho migration the weir was inoperable on a few occasions. Specific periods in which the counting facility was inoperable and the number of coho added are listed in Table 2. A total of 16 coho were added to the estimate during periods when the counting facility was not operational. In addition 17 unidentified fish were assigned to be coho. Any unidentified fish were assigned species composition based on the proportion of known daily observations. None of the 878 coho salmon observed were coho salmon that previously entered IGH and were subsequently released (released coho from IGH were externally marked with “T-Bar” anchor Floy tags for identification). During 2010 KRP staff attempted to estimate the number of grilse in the Scott River by enumerating the number of coho observed in the video flume that were shorter or longer than 56 cm. Utilizing this method KRP staff identified 865 (98.5%) adults and 13 (1.5%) grilse.




Figure 8. Run timing of coho salmon observed passing through the Scott River Fish Counting Facility during the 2010 season (N=918). No downstream movements were observed of coho salmon.

Diel movements of coho salmon through the SRFCF were higher in the afternoon hours and peaked between 1400 hours and 1500 hours (Figure 9). This movement pattern is not consistent with observations from previous seasons. Migrations were fairly consistent throughout the day with reduced movement during the late morning hours. The hours between 0700 and 0900 was generally the time during the day when the crew was at the weir conducting daily maintenance.




Figure 9. Diel migration patterns of coho salmon observed moving through the Scott River Fish Counting Facility in 2010 (N=877).

Table 2. Specific dates and times when the filming stopped and restarted, the number of hours without data and the number of coho estimated during that time.

	Date	Time	Number of hours without data	Number of Coho estimated
Filming Stopped	10/24/2010	2030	3.5	
Filming Started	10/25/2010	930	9.5	5
Filming Stopped	12/6/2010	1600	8	4
Filming Started	12/7/2010	930	9.5	4
Filming Stopped	12/10/2010	1000	14	
No Data available	12/11/2010		24	1
No Data available	12/12/2010		24	1
No Data available	12/13/2010		24	1
No Data available	12/14/2010		24	
No Data available	12/15/2010		24	
No Data available	12/16/2010		24	
No Data available	12/17/2010		24	
Filming Started	12/18/2010	615	6.15	
Totals			218.65	16

SPAWNING GROUND SURVEYS

No coho carcasses were observed during the cooperative spawning ground survey on the mainstem Scott River below the weir or in the tributary surveys below the counting facility, although redds were observed in Tompkins Creek (2), Canyon Creek (1) and Kelsey Creek (5). The eight redds observed below the weir were estimated to represent 16 adult coho. No additional jacks were estimated to accompany the 8 observed redds (estimated 0.24 jacks was rounded to zero). Twenty nine coho carcasses were observed in the Scott River during surveys coordinated by the Siskiyou Resource Conservation District (SRCD) above the counting facility (Yokel, 2011). Six additional coho carcasses (one unmeasured) were collected as washbacks at the counting facility (Figure 10). As reported in Quigley 2011 the sex ratio of observed coho salmon carcasses in the Scott River during 2010 was 68.6% (24) female and 31.4% (11) male. None of the 35 carcasses examined had a maxillary clip indicating that the return of adults was 100% of natural origin. Based on the fork length frequency distribution of the limited number of carcasses collected during the season all of the fish would have been estimated to be age three (See video section for further information on age structure). All of the carcasses examined were sampled for tissue and collected samples were supplied to the NOAA Southwest Fisheries Science Center located in Santa Cruz, California for stock identification purposes.




Figure 10. Length frequency distribution of male and female coho salmon observed during RCD spawning ground surveys ((29); Yokel 2011) and as washbacks (5) on the Scott River Fish Counting Facility, during the 2010 spawning season (n = 34).

A total of 911 coho salmon were observed moving upstream through the Scott River Fish Counting Facility (SRFCF) during the season. Eight coho redds were estimated in areas below the counting facility representing an estimated 16 coho. The total number of coho salmon that entered the Scott River during the 2010 season is estimated to be **927**. Utilizing the observed age proportions, derived from video data, the resulting number of age two and three fish are 14 and 913 respectively.

STEELHEAD

In 2010, a net total of 419 adult (>16") steelhead (121 upstream, 2 downstream; Figure 11) and 132 sub-adult (<16") steelhead (135 upstream, 3 downstream; Figure 12) were estimated to have entered and remained in the Scott River during the video recording season from September 29, 2010 to January 7, 2011. Peak migration of adult steelhead was observed during the 13 day period between October 26, 2010 and November 7, 2010 when 81.6% of the adult steelhead were observed. Lines on the back of the video flume were set at 16 inches (40.64 cm) to delineate sub-adults versus adults. The 2010 season was the first year that lines delineating adult steelhead and sub-adults was used. As additional years of data are collected annual comparisons will be presented.




Figure 11. Run timing of steelhead trout (>16'') observed passing through the Scott River Fish Counting Facility during the 2010 season (N=423).

DISCUSSION

CHINOOK SALMON RUNS

Since 1978 the Chinook salmon run in the Scott River has ranged from 14,477 fish (1995) to 467 fish (2004) and has averaged 5,200 fish (Figure 13). The 2010 Chinook salmon run in the Scott River ranks twenty-seventh (2,508 fish) out of 33 years of population estimates. The 2010 run was 52% lower than the 33 year average. A total of 2,228 Chinook salmon were estimated to have passed through the Scott River Fish Counting Facility during the 2010 season. A total of 280 Chinook salmon carcasses were estimated in reach 1 through reach 6, yielding a total run size estimate of 2,508 Chinook salmon. A total of 549 Chinook salmon were estimated in reaches 7 and 8. If the total number of Chinook estimated in reach 7 and 8 are subtracted from the counting facility estimate the proportion of the run that utilized areas in and upstream of the valley can be estimated. During 2010, 75.4% (1,679) of the Chinook run utilized areas of the watershed above Member Bridge. Reach 8, a 3.4 mile section of the river, between Member Bridge and the USGS gauging station accounted for 378 Chinook salmon. These 378 fish that utilized reach 8 accounted for 17.0% and 15.0% of the total spawning above the weir and total spawning throughout the entire watershed respectively. The Schaffer mark recapture carcass estimate for the

entire watershed that was surveyed was 1,031 fish. These 1,031 fish underestimate the total estimate by 58.9%. The addition of the fish counting facility has yielded a more accurate estimate of the




Figure 12. Run timing of steelhead trout (<16'') observed passing through the Scott River Fish Counting Facility during the 2010 season (N=138).

total number of Chinook in the Scott River and has allowed for accurate estimation of Chinook utilization in the valley reaches without having to survey these reaches.




Figure 13. Estimated escapement of adult Chinook salmon returning to the Scott River from 1978 to 2010.

The Scott River is an important component of the Klamath Basin Chinook runs. Table 3 shows that the Scott River has contributed an average of 10 percent of the basin-wide natural spawning escapement during the period from 1978 to 2010. The production of emigrating 0+ Chinook has been estimated in the Scott River since Brood Year 1999 (Chesney et al. 2011). The number of 0+ Chinook produced per adults is a direct measure of in-river productivity and as habitat conditions improve or diminish this measure will reflect those conditions. It is encouraging that the number of 0+ Chinook produced per adult is increasing and has increased each season starting with brood year 2005 (Figure 14).

Table 3. Klamath Basin and Scott River Chinook natural spawner escapements (age 2-5), 1978-2010.

Year	Chinook Natural Spawner Escapement		% Scott
	Klamath Basin	Scott River	
1978	74,906	5,332	7%
1979	37,398	3,824	10%
1980	48,465	4,277	9%
1981	50,364	6,556	13%
1982	50,597	10,176	20%
1983	33,310	3,568	11%
1984	21,349	1,801	8%
1985	61,628	4,408	7%
1986	142,302	8,041	6%
1987	110,489	8,566	8%
1988	91,930	5,200	6%
1989	49,377	4,188	8%
1990	16,946	1,615	10%
1991	12,367	2,165	18%
1992	17,171	2,838	17%
1993	25,683	5,300	21%
1994	38,578	2,863	7%
1995	179,118	14,477	8%
1996	87,500	12,097	14%
1997	50,369	8,561	17%
1998	45,343	3,327	7%
1999	28,904	3,584	12%
2000	89,122	6,253	7%
2001	85,581	6,142	7%
2002	69,502	4,308	6%
2003	89,744	12,053	13%
2004	28,516	467	2%
2005	27,931	756	3%
2006	45,002	4,960	11%
2007	61,741	4,505	7%
2008	48,073	4,673	10%
2009	52,702	2,211	4%
2010	49,027	2,508	5%
Average	58,213	5,200	10%




Figure 14. Number of 0+ Chinook produced per adult spawner in the Scott River by brood year, for Brood Years 1999-2009.

COHO SALMON

Since video operations began in 2007 the estimated escapement of coho salmon in the Scott River has been 1,622, 63, 81 and 927 for 2007, 2008, 2009 and 2010 respectively (Figure 15). The adult run size of coho salmon in the recent past prior to 2007 was unknown and with the addition of the counting facility the Department’s ability to monitor this listed run has greatly improved. Although recent adult run size data is sparse on the Scott River, monitoring of the yearling juvenile emigration has taken place since 2003. The emigration data generated from 2003 through 2010 indicates significant variation in brood year strength (Chesney, 2011). Results of the first four years of adult monitoring at the SRFCF support this observation. The cohort that returned in 2010 is the strongest of the three year classes in the Scott River.




Figure 15. Estimated escapement of adult coho salmon returning to the Scott River from 1978 to 2010.

Utilizing the number of yearling coho produced in the Scott River (Chesney, 2011) and the results of the adult abundance estimates allows for analysis of freshwater production and out of basin survival by brood year. For brood years 2004, 2005 and 2007 the number of yearling coho that were required to produce a single adult coho averaged 58.94 and ranged from a low of 46.30 to a high of 67.11. The corresponding out of basin survival has averaged 1.74 percent and ranged from a low of 1.49 percent to a high of 2.16 percent (Table 4). Utilizing the estimated number of yearling outmigrants and the average percent yearling survival value of 1.74 percent the number of returning adults in 2011 can be forecasted at 37. Due to the difficulty in estimating abundance of outmigrants at low abundance levels, data from brood year 2006 has been omitted from this analysis. Although the proportion of yearlings that survive outside the Scott River watershed is largely driven by uncontrollable factors it is important to track this survival metric to accurately evaluate ongoing restoration efforts taking place within the watershed.

Table 4. Yearling coho outmigrant abundance point estimates, adult coho abundance estimates, ratio of outmigrant yearlings to adult returns and proportion of outmigrant yearlings that returned as adults by brood year for the Scott River, Brood Years 2004-2008.

Brood Year	Yearling Year	Yearling point Estimate	Adult Year	Adult Estimate	Yearlings to adult	Percent yearling survival
2004	2006	75097	2007	1622	46.30	2.16
2005	2007	3931	2008	62	63.40	1.58
2006	2008	941	2009	81	11.62	8.61
2007	2009	62207	2010	927	67.11	1.49
2008	2010	2174	2011	37 ^{/2}	58.94 ^{/2}	1.74 ^{/1}

/1 Average percent yearling survival from brood years 2004, 2005 and 2007

/2 Projected adult estimate and yearling to adult ratio based on yearling point estimate of 62,207 and average percent yearling survival from brood years 2004, 2005 and 2007.

Analyzing the comparisons of yearling coho production estimates to estimated adult coho returns

produces freshwater survival estimates in the form of yearling coho produced per adult return. Brood Years 2007 and 2008, the only years in which the number of yearling coho produced per returning adult can be calculated, and are 38.35 and 34.51 respectively (Table 5). As additional years of data become available the freshwater production of coho salmon in the Scott River can be further evaluated. To give some context to the yearlings produced per adult in the Scott River, this value in the Shasta River has averaged 18.6 and ranged from a low of 4.4 to a high of 38.0 for brood years 2001-2007. The number of yearlings produced per returning adult by brood year is a direct measure of freshwater survival. For levels below carrying capacity, it can be stated that as the number of yearlings produced per returning adult increases it can be inferred that in-river conditions for coho salmon are improving. Conversely as the number of yearlings produced per returning adult decreases it can be inferred that in-river conditions for coho salmon are getting worse.

Table 5. Adult coho estimate, yearling coho production point estimate and ratio of yearling coho produced per adult return for the Scott River, Brood Years 2007 and 2008.

Adult Year Brood Year	Adult Estimate	Yearling Year	Yearling point Estimate	Yearlings produced per adult
2007	1622	2009	62207	38.35
2008	63	2010	2174	34.51

ACKNOWLEDGEMENTS

The California Department of Fish and Game would like to thank Sport Fish Restoration Fund for their financial assistance in the operation of the SRFCF during 2010. The cooperative spawning ground surveys would not be possible without the assistance of the U.S. Fish and Wildlife Service Yreka office and the USFS. Additionally this work would not be possible without the cooperation and participation of the Siskiyou Resource Conservation District, Karuk Tribe, Quartz Valley Tribe, USFS, Northern California Resource Center, Americorps and Etna High School who assisted KRP staff in completing spawning ground surveys on the Scott River. We would also like to express our appreciation to the various landowners who have graciously provided permission to access the Scott River on their lands.

Literature Cited

- Chesney, W. Daniels, S. Debrick, A. Diviney, C. Underwood, K. Stenhouse, S. 2011. Final Report Shasta and Scott River Juvenile Salmonid Outmigrant Study, 2010. P0710307. California Department of Fish and Game Available from California Department of Fish and Game, 1625 South Main Street, Yreka, CA. 96097
- Klamath River Technical Team (KRTAT) 2011. Klamath River Fall Chinook Salmon Age-Specific Escapement, River Harvest, and Run Size Estimates, 2010 Run.
- Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. Canada. Dep. of Environ., Fish. and Mar. Serv. Bull. 191. 382 p.
- Stevenson, David R. and Steven E. Campana. 1992. Otolith Microstructure Examination and Analysis. Department of Marine Resources, Marine Resources Laboratory, McKown Point, West Boothbay Harbor, Maine 04575.
- Yokel, D. 2011. Scott River Adult Coho Spawning Ground Surveys 2010-2011 Season. Siskiyou Resource Conservation District report Agreement# 813339G030