

California Fish and Game 96(4): 223-231; 2010

## **PATHOGEN SURVEY, GILL Na-K-ATPase ACTIVITY, AND LEUKOCYTE PROFILE OF ADULT DELTA SMELT**

J. SCOTT FOOTT and JOHN BIGELOW  
U.S. Fish and Wildlife Service  
California-Nevada Fish Health Center  
24411 Coleman Hatchery Road  
Anderson, CA 96007  
Scott\_Foott@fws.gov

**Delta smelt abundance has declined drastically and the species was listed as threatened in 1993. The role of disease in this decline is largely unknown. One hundred and five adult smelt, collected from the lower Sacramento River between January and May 2010, were surveyed for infectious agents, blood leukocyte profile, and gill Na-K-ATPase activity. Few tissue changes or significant parasitic infections were observed in histological specimens and there was a low incidence of bacterial isolations. Asymptomatic *Mycobacterium sp.* infection was detected in 54% of the samples by PCR; however, this bacterial group was not isolated in culture. Gill Na-K-ATPase activity was lower in the January sample than subsequent month collections. The number of blood granulocytes increased between March and May. Disease did not appear to be an overt influence on the surveyed population in the spring of 2010.**

**Keywords:** California, delta smelt, *Hypomesus transpacificus*, parasites, pathology, potassium, sodium, PCR, Sacramento River Delta

### **INTRODUCTION**

The delta smelt (*Hypomesus transpacificus*) was listed as threatened under the Endangered Species Act in 1993 and is endemic to the upper San Francisco Bay Estuary (Moyle et al. 1992). In the last decade, a clade of pelagic organisms in the delta has declined in abundance (Feyrer et al. 2007). Potential factors associated with these declines in delta smelt include reduction in freshwater flows, entrainment losses at diversions and power plants, inadequate food base and competition for food from exotic species, environmental contaminants, and predation by exotic fishes. Only limited work has been reported on smelt disease (Antonio et al. 2000, Teh 2007). Our objective was to determine the presence of infectious pathogens (viruses, bacteria, or parasites), tissue abnormalities (gill, liver, intestinal tract), peripheral blood cell profile, and gill Na-K-ATPase activity in adult smelt captured in the lower Sacramento River between January and May 2010.

METHODS

One hundred and five adult smelt were sampled from the California Department of Fish and Game (CDFG) Bay-Delta Kodiak trawl collections in the Sacramento River during January–May 2010 (Table 1). After initial evaluation of maturity by a CDFG biologist, the following actions were taken: (1) fish were euthanized in 100 mg/L benzocaine; (2) measured by caliper (standard length; mm), weighed (Pesola micro-line 20010 hanging balance, 0.1 g), and condition factor calculated ( $(\text{g}/\text{mm}^3) \times 10^5$ ); (3) caudal peduncle severed and blood collected in microhematocrit tube; (4) blood was transferred into cold L-15 separation solution or smeared directly onto glass slides; (5) gill tissue placed into 100 microliter Sucrose-EDTA-Imidazole solution and frozen on dry ice; (6) liver aseptically removed with half placed into Brain Heart Infusion (BHI) broth transport media and the other half placed into a tube and frozen on dry ice; (7) anterior kidney removed and placed into cold antibiotic-

**Table 1.** Delta smelt collection dates in 2010, site numbers, (Montezuma slough [606]; Sacramento River from Brannon Island [704] upstream to the deep water ship channel [719]) and associated sample of delta smelt per site, January - April 2010. If no sample was collected it is designated as “NT”. One 30 mm larva was included in the collection from site 719 on 5 May.

| Location | Date   |        |        |        |       |       |
|----------|--------|--------|--------|--------|-------|-------|
|          | 13 Jan | 10 Feb | 10 Mar | 11 Mar | 7 APR | 5 MAY |
| 606      | NT     | NT     | NT     | 2      | NT    | NT    |
| 704      | 8      | NT     | 5      | NT     | 12    | 2     |
| 706      | 6      | 1      | 4      | NT     | NT    | NT    |
| 707      | NT     | NT     | 2      | NT     | NT    | NT    |
| 713      | 1      | NT     | 4      | NT     | NT    | NT    |
| 715      | 3      | 7      | 7      | NT     | NT    | 1     |
| 716      | NT     | 2      | 6      | NT     | NT    | 1     |
| 719      | 5      | 2      | 12     | NT     | 25    | 2     |

antimycotic solution; and, (8) gill and viscera placed into Davidson’s fixative for 24 h and processed for 6  $\mu\text{m}$  histological sections. The remaining carcass (head into ethanol, body and gonad into 10% formalin) was given to CDFG for use by other researchers.

Anterior kidney was pooled (1 to 5 fish) in 1 ml antibiotic /mycotic solution and held on ice for 18 - 24 h until processed (American Fisheries Society-Fish Health Section 2007). The samples were sonicated (Heat system Microson XL2005, Farmingdale NY) for 10 s at #3 setting to disrupt the tissue, centrifuged at  $10,000 \times g \times 4^\circ\text{C}$ , supernatant diluted 2x and 4x in Hanks Buffered Salt Solution, and 100 microliter samples inoculated onto duplicate wells of EPC, CHSE214, BF2, and SSN1 cell cultures maintained in 48 well plates. Cultures were incubated for 18 d at either 15  $^\circ\text{C}$  (EPC and CHSE214) or 25  $^\circ\text{C}$  (BF2 and SSN1) and examined every 3 or 4 days.

Liver in BHI broth was inoculated onto BHI agar and incubated at 20 - 25 $^\circ\text{C}$  for 48 - 72 h as well as Middlebrook 7H10 media. For the 5 May collection, Lowenstein-Jenson 5% NaCl media was also inoculated. Both Middlebrook and Lowenstein-Jenson media were incubated at 25 $^\circ\text{C}$  for 18 – 20 d. Any colony obtained from either Middlebrook or Lowenstein-Jenson media was acid-fast stained to presumptively identify *Mycobacterium* (Antonio et al. 2000, Stine et al. 2010). An ATCC culture of *Mycobacterium marinum* (ATCC 11565) was

grown on Middlebrook 7H10 media to verify its suitability for the survey. Isolates from BHIA were identified to genus or clade (American Fisheries Society-Fish Health Section 2007). DNA from frozen liver was extracted with a Qiagen Cube using the manufacturer's procedure and sent to the Real-time PCR Research and Diagnostics Core Facility, School of Veterinary Medicine, Department of Medicine and Epidemiology, University of California, Davis to be analyzed in their proprietary MTC assay for multiple *Mycobacterium* species.

Fixed tissue was processed for 6  $\mu$ m paraffin sections and stained with hematoxylin and eosin (H&E). Sagittal sections, of a 30 mm larvae collected on 5 May, were stained by both H&E and Nissel stain for neurons. Tissues examined microscopically included gill, liver, intestine, adipose tissue, pancreatic acinar cells, spleen, and heart. Not all tissues were sectioned for each fish.

In March, blood smears were immediately prepared using a cover slip, air-dried, fixed in cold methanol for 5 minutes, and protected from light until further processing. During April and May, blood was dispensed from the hematocrit tubes into Eppendorf tubes containing L-15 separation solution (1,000  $\mu$ l of Lebowitz L-15 media, 75  $\mu$ l of 7.5% fetal bovine serum, and 2  $\mu$ l of 5 KU/ml heparin). Blood smears were then prepared 4-8 h later by brief vortexing, placing 200  $\mu$ l of the suspension into a single well cyto-spin chamber, centrifuged at 1,600 rpm for 4 min in a StatSpin Cytofuge (StatSpin, Inc., Norwood, MA), air-dried and fixed in cold methanol for 5 minutes. For panoptical staining, blood smears were treated according to a modified Leishman-Giemsa staining protocol of Yasutake and Wales (1983). For each peripheral blood smear, 81-115 leukocytes were classified as lymphocytes, granulocytes, or thrombocytes on the basis of panoptical staining characteristics (Blaxhall and Daisley 1973) by examination at 100x magnification and expressed as a proportion of the total number of leukocytes classified. A lymphocyte to granulocyte ratio was calculated for each sample (Modra et al. 1998). Gill Sodium-Potassium-Adenosine Triphosphatase activity (ATPase =  $\mu$ moles ADP/mg protein/hr) was assayed by the method of McCormick and Bern (1989). Statistical analysis was performed with SigmaStat 3.1 software on raw data. Normality was tested with the Kolmogorov-Smirnov method at the  $P=0.05$  level. One-way ANOVA or T-test (data with normal distribution, reported with  $F$  or  $t$  value) or Kruskal-Wallis ANOVA or Mann-Whitney U test with subsequent multiple comparison procedures (Holm-Sidak or Dunns method respectively,  $\alpha \leq 0.05$ ) was used to compare groups.

## RESULTS

Male smelt averaged 62 mm in standard length (SL) while the mean female SL was 66 mm (Table 2). Similarly, female wet weight and condition factor were higher than males due to their gravid condition. No abnormalities (cytopathic effects) were observed in the 4 cell lines over the 18 d incubation periods (Table 3). The eighteen anterior kidney pools (1-5 fish / pool) tested contained a total of 79 smelt. It was necessary to add fresh media to SSN1 and BF2 cultures to maintain them for the full 18 days. Liver samples inoculated onto the general purpose media, Brain Heart Infusion Agar, isolated a low incidence of aquatic bacteria (*Aeromonas* / *Pseudomonas* sp. and *Staphylococcus* / *Micrococcus* sp.). Middlebrook cultures, collected on 13 January, were contaminated with fungi and environmental bacteria related to the outside inoculation conditions and discarded. On subsequent trips, fish dissection and tissue collection were done within the boat's cabin. No acid-fast bacteria were isolated from either the Middlebrook 7H10 or Lowenstein-Jensen 5% NaCl media (Table 3). In contrast, 54% of the livers tested positive for *Mycobacterium* sp. DNA by PCR

**Table 2.** Mean (SE) standard length, weight, and condition factor ((g/mm<sup>3</sup>) x 10<sup>5</sup>) of delta smelt collected from the lower Sacramento River, California, January - May, 2010. Sample number (N) for fish identified to sex is less than total number of fish collected in 2010.

| Metric               | Male                  | Female                |
|----------------------|-----------------------|-----------------------|
| Standard length (mm) | 62 (4), N = 30        | 66 (4), N = 60        |
| Weight (g)           | 2.1 (0.3), N = 26     | 2.7 (0.6), N = 59     |
| Condition factor     | 0.895 (0.119), N = 26 | 0.936 (0.115), N = 59 |

assay. These *Mycobacterium* data indicate a high incidence of asymptomatic, carrier-state infections but not a significant health problem for the sample population.

**Table 3.** Viral assay, bacterial culture (Brain heart infusion agar [BHIA] and mycobacterium media Middlebrook 7H10 [MB]), and *Mycobacterium sp.* (Mycob) PCR results for delta smelt collected from the lower Sacramento River, California, January - May 2010. Data are reported as number positive/monthly sample number (or viral pools) collected in Kodiak trawls and incidence (total positive/total samples [%]). If sample was not performed or collected it is designated as “ND”.

| Assay         | 13 Jan              | 10 Feb      | 10 Mar | 11 Mar      | 7 Apr                                      | 5 May              | Incidence                                          |
|---------------|---------------------|-------------|--------|-------------|--------------------------------------------|--------------------|----------------------------------------------------|
| Viral (pools) | 0 / 5               | 0 / 3       | ND     | 0 / 1       | 0 / 8                                      | 0 / 1              | 0 / 18                                             |
| BHIA          | 5 / 23 <sup>a</sup> | 0 / 12      | ND     | 0 / 2       | 2 / 33 <sup>a</sup><br>2 / 33 <sup>b</sup> | 0 / 5              | 7 / 75 (9) <sup>a</sup><br>2 / 75 (3) <sup>b</sup> |
| MB            | ND                  | 0 / 12      | ND     | 0 / 2       | 0 / 33                                     | 0 / 5 <sup>c</sup> | 0 / 19                                             |
| Mycob PCR     | 18 / 23 (78)        | 6 / 11 (55) | ND     | 2 / 2 (100) | 9 / 27 (33)                                | 1 / 4 (25)         | 36 / 67 (54)                                       |

<sup>a</sup>*Aeromonas* / *Pseudomonas* sp.  
<sup>b</sup>*Staphylococcus* / *Micrococcus* sp.  
<sup>c</sup>Lowenstein – Jenson 5%NaCl media also used for *Mycobacterium* isolation

Few tissue abnormalities and no overt infectious disease was observed in 72 histological specimens (Table 4). Helminth parasites (nematode and trematode) were seen in < 10% of intestine, visceral adipose, or liver sections. No lesions were associated with the parasites. No parasitic infection was seen in gill sections. Focal liquefactive necrosis was observed in 7% of the liver sections and occurred primarily in the 10 February collection. The cause of the lesion is unclear, as there was no association with bacterial or parasitic infection, but post-mortem artifact cannot be ruled out. No acid-fast bacteria (e.g, *Mycobacterium*) were observed in a pancreatic acinar tissue granuloma from a 13 January

sample. Spleen contained multifocal regions of endogenous pigments / melanomacrophage centers. Small foci of lipofuscin were seen in the visceral adipose tissue of 2 fish. No necrotic neurons were seen in the brain and retina (Nissel stain) of the 30 mm larvae collected on 5 May.

**Table 4.** Histological observation of parasites or tissue abnormalities such as nematode (nem), trematode (trm), granulomatous foci (grn), inflammatory cell foci (icf), liquefactive necrosis foci (lqn), and endogenous brown pigment foci (bpf) among delta smelt collected from the lower Sacramento River, January - May 2010. Data are reported as the number of specific tissues positive for either parasites or abnormality/total sections by date of collection and incidence (total positive/total samples [%]). If no specific tissue in sections of a given collection it is designated as "NT".

| Tissue              | Date          |           |               |       |                     |              | Incidence                                                   |
|---------------------|---------------|-----------|---------------|-------|---------------------|--------------|-------------------------------------------------------------|
|                     | 13 Jan        | 10 Feb    | 10Mar         | 11Mar | 7 Apr               | 5 May        |                                                             |
| Gill                | 0 / 12        | 0 / 12    | 0 / 21        | 0 / 2 | 0 / 14 <sup>a</sup> | 0 / 4        | 0 / 65 (0)                                                  |
| Heart               | 0 / 6         | 0 / 4     | 0 / 6         | NT    | 0 / 5               | NT           | 0 / 21 (0)                                                  |
| Intestine           | 2/15<br>nem   | 0 / 12    | 0 / 23        | 0 / 2 | 0 / 16              | 0 / 4        | 2 / 72 (3)<br>nem                                           |
| Stomach             | 0 / 16        | 0 / 12    | 0 / 19        | 0 / 2 | 0 / 12              | 0 / 3        | 0 / 64 (0)                                                  |
| Acinar<br>cells     | 1 / 15<br>grn | NT        | 0 / 21        | NT    | 0 / 12              | 0 / 3        | 1 / 51 (2)<br>grn <sup>b</sup>                              |
| Visceral<br>Adipose | 0 / 15        | 0 / 11    | 1 / 21<br>nem | 0 / 1 | 1 / 11<br>nem       | 1 / 3<br>trm | 2 / 62 (3)<br>nem<br>1 / 62 (2)<br>trm                      |
| Liver               | 1 / 15<br>icf | 4 / 7 lqn | 1 / 21<br>trm | 0 / 1 | 0 / 2               | 0 / 3        | 1 / 59 (2)<br>trm<br>4 / 59 (7)<br>lqn<br>1 / 59 (2)<br>icr |
| Spleen              | 3/3 bpf       | 2 / 2 bpf | NT            | NT    | 2 / 2 bpf           | NT           | 7 / 7 (100)<br>bpf                                          |

<sup>a</sup> Late fixation resulted in variable levels of post-mortem changes in tissue morphology.

<sup>b</sup> No acid fast bacteria observed in granuloma

Gill Na-K-ATPase activity was significantly lower in the 13 January samples compared to subsequent collection groups, but low sample numbers reduced our confidence in its biological significance (Table 5, ANOVA  $F=6.984$ ,  $P=0.001$ ). Twenty-five percent of the samples did not produce enzymatic activity. Protein concentrations of these “no activity” samples were similar to the other samples. We suspect the quantity of muscle and bone was in excess of gill filament in these samples, but were not able to rule out other explanations. Eight of the ten “no activity” samples occurred in the January and February collection groups. Mean activities of smelt ranged from 10.7 to 20.9; in comparison, mean gill Na-K-ATPase activity ranges of juvenile Chinook smolts assayed by the same methods ranged from 7.0 - 11.2  $\mu\text{mole ADP/mg protein/hr}$  (Foott et al. 2007).

**Table 5.** Mean (SE) Gill Na-K-ATPase activity (micromole ADP/mg protein/hr), total gill sample set, and number of samples showing no activity (not included in analysis) from adult delta smelt collected from the lower Sacramento River by sample date, January - May 2010. Letters in common indicate differences were not significant. The single sample from 11 March was not compared to other groups.


|             | Date         |              |        |              |              |
|-------------|--------------|--------------|--------|--------------|--------------|
|             | 13 Jan       | 10 Feb       | 11 Mar | 7 Apr        | 5 May        |
| Mean (SE)   | 10.7 (1.2) A | 20.9 (2.8) B | 8.6    | 17.6 (0.9) B | 19.0 (2.4) B |
| Total       | 13           | 12           | 2      | 8            | 5            |
| No activity | 3            | 5            | 1      | 1            | 0            |

Thrombocytes, lymphocytes, and granulocytes were present in proportions of 0.876, 0.119, and 0.006 in males, of which 8 of 9 males were captured during March (Table 6). Females displayed the same order of prevalence (thrombocyte>lymphocyte>granulocyte)

**Table 6.** Proportion of leukocytes (mean [SE], range, and coefficient of variation [CV]) in peripheral blood of delta smelt collected from the lower Sacramento River, California, January - May 2010.

| Type of leukocyte | Males (n = 9)  | Females (n = 24) | Both Sexes (n = 33) |
|-------------------|----------------|------------------|---------------------|
| Lymphocytes       |                |                  |                     |
| Mean (SE)         | 0.119 (0.0285) | 0.167 (0.0336)   | 0.154 (0.0257)      |
| Range             | 0.030 - 0.300  | 0.000 - 0.610    | 0.000 - 0.610       |
| CV                | 0.719          | 0.989            | 0.962               |
| Granulocytes      |                |                  |                     |
| Mean (SE)         | 0.006 (0.0024) | 0.045 (0.0139)   | 0.034 (0.0105)      |
| Range             | 0.000 - 0.020  | 0.000 - 0.291    | 0.000 - 0.291       |
| CV                | 1.308          | 1.526            | 1.780               |
| Thrombocytes      |                |                  |                     |
| Mean (SE)         | 0.876 (0.0300) | 0.789 (0.0416)   | 0.813 (0.0318)      |
| Range             | 0.690 - 0.970  | 0.301 - 1.000    | 0.301 - 1.000       |
| CV                | 0.103          | 0.258            | 0.225               |

in peripheral blood leukocyte types as males during March, April, and May. Granulocyte numbers increased over time in the females, resulting in a lower lymphocyte:granulocyte ratio in May compared to March or April (Figure 1).



**Figure 1.** Mean number of lymphocytes and granulocytes counted (bar = SE) in peripheral blood of female delta smelt collected from the Sacramento River, California, January - May 2010. Insufficient male samples excluded them from temporal analysis.

## DISCUSSION

Adult delta smelt, collected from the lower Sacramento River in the spring of 2010, were apparently healthy as demonstrated by lack of clinical signs or morbidity, few tissue changes or significant parasitic infections observed in histological specimens, or microbiological isolations. While a high incidence of asymptomatic *Mycobacterium* sp. infection was detected by PCR, this bacterial group was not isolated in culture. Antonio et al. (2000) reported the occurrence of *Mycobacterium* disease in delta smelt broodstock held at  $\geq 16^{\circ}\text{C}$  and stressed by handling. Similar mycobacteria disease has occurred in captive post-spawning smelt held at Livingston Stone National Fish Hatchery. In contrast to a stressful culture situation, Antonio et al. (2000) were not able to culture *Mycobacterium* in recently caught smelt. Similar to the histological data, Teh (2007) reported no significant disease or parasitic infection observed in 385 adult smelt collected from all regions of the Delta in 2005. Teh (2007) noted that liver abnormalities (glycogen depletion, cytoplasm eosinophilic inclusion, fat vacuoles, single cell necrosis, and inflammation) were common observations.

The occurrence of melanomacrophage foci in the spleen has been reported in other fish. Melanomacrophage centers are common in fish hemopoietic tissues and act as depositories of oxidized materials, scavenged iron, and some micro-parasites (Aguis and

Roberts 2003). The macrophages contain a mixture of endogenous pigments such as lipofuscin/ceroid, melanin, and hemosiderin. The extent of such foci has been associated with contaminate exposure or hypoxic conditions (Fournie et al. 2001).

We do not have an adequate explanation for the “no activity” gill Na-K-ATPase samples collected in January and February. While collection, storage, and processing error could be the cause, metal exposure has been reported to impair Na-K-ATPase activity (Lauren and McDonald 1985, Roger et al. 2003). Delta smelt are exposed to a variety of contaminants that could affect their health and physiology (Kuivila and Moon 2004, Foe 1995).

Increased granulocyte numbers in the blood between March and May are counter to that observed by other workers where lymphocyte numbers and overall specific immune function tend to increase with increasing seasonal temperature. Alcorn et al. (2002) reported temperature influence on lymphocyte to granulocyte ratios, with greater numbers of peripheral blood lymphocytes in sockeye salmon reared at 12 C than 8 C. Similarly, Luskova (1998) noted the white blood cell count (leukocyte) increased with higher seasonal water temperatures in brown trout. Higher granulocyte counts result in low L:G ratio values, and can indicate infection, tissue damage, or seasonal blood cell changes (Modra et al 1998). It is unclear what influenced the higher granulocyte numbers in the May collection, given that water temperature increased throughout the spring and we did not detect clinical infections or tissue lesions in the gill or viscera.

The extreme stress of capture and handling associated with the Kodiak trawl would likely influence blood measurements. Delta smelt are extremely sensitive to handling stress (Swanson et al. 1996), and the effect on peripheral blood cells could induce an artifact. It is unlikely that leukocyte evaluation will be a valid biomarker for captured smelt. Physiological measurements that utilize tissue or blood are limited by the small size of smelt. The use of sentinel fish is one option for examining the effects of water quality on the health and physiology of these fish.

#### ACKNOWLEDGMENTS

We thank the California Department of Fish and Game for access to delta smelt collected in the 2010 Kodiak trawl, and especially appreciate the patience and good humor of Julio Adib-Samii. Ann Bolick assisted with DNA extraction and Ron Stone helped with histological processing.

#### LITERATURE CITED

- American Fisheries Society – Fish Health Section. 2007. Fish health section bluebook: suggested procedures for the detection and identification of certain finfish and shellfish pathogens, 2007 edition. AFS-FHS, Bethesda, Maryland, USA.
- Agius, C. and R. J. Roberts. 2003. Melano-macrophage centres and their role in fish pathology. *Journal of Fish Pathology* 26:499-509.
- Alcorn, S. W., A. L. Murray, and R. J. Pascho. 2002. Effects of rearing temperature on immune functions in sockeye salmon (*Oncorhynchus nerka*). *Fish and Shellfish Immunology* 12:303-334.
- Antonio, D. B., C. Swanson, J. J. Cech, R. C. Mager, S. Doroshov, and R. P. Hedrick. 2000. Prevalence of *Mycobacterium* in wild and captive delta smelt. *California Fish and Game* 86:233-243.

- Blaxhall, P. C., and K. W. Daisley. 1973. Routine haematological methods for use with fish blood. *Journal of Fish Biology* 5:771-781.
- Feyrer, F., M. L. Nobriga, and T. R. Sommer. 2007. Multidecadal trends for three declining fish species: habitat patterns and mechanisms in the San Francisco Estuary, California, USA. *Canadian Journal of Fisheries and Aquatic Sciences* 64:723-734.
- Foe, C. 1995. Evaluation of the potential impact of contaminants on aquatic resources in the central valley and Sacramento-San Joaquin Delta estuary. Report D-037262, Central Valley Regional Water Quality Control Board, Sacramento, California, USA.
- Foott, J. S., R. Stone, and K. Nichols. 2007. Proliferative kidney disease (*Tetracapsuloides bryosalmonae*) in Merced River Hatchery juvenile chinook salmon: mortality and performance impairment in 2005 smolts. *California Fish and Game* 93:57-76.
- Fournie, J. W., J. K. Summers, L. A. Courtney, V. D. Engle and V. S. Blazer. 2001. Utility of splenic macrophage aggregates as an indicator of fish exposure to degraded environments. *Journal of Aquatic Animal Health* 13:105-116.
- Kuivila, K. M., and E. G. Moon. 2004. Potential exposure of larval and juvenile delta smelt to dissolved pesticides in the Sacramento-San Joaquin Delta, California. *American Fisheries Society Symposium* 39:229-241.
- Lauren, D. J., and D. G. McDonald. 1985. Effects of copper on branchial ionregulation in the rainbow trout, *Salmo gairdneri* Richardson. *Journal of Comparative Physiology B*, 155:635-644.
- Luskova, V. 1998. Factors affecting haematological indices in free-living fish populations. *Acta Veterinaria Brno* 67:249-255.
- McCormick, S. D., and H. A. Bern. 1989. In vitro stimulation of  $\text{Na}^+/\text{K}^+/\text{ATPase}$  activity and ouabain binding by cortisol in coho salmon gill. *American Journal of Physiology* 256:R707-715.
- Modra, H., Z. Svobodova and J. Kolarova. 1998. Comparison of differential leukocyte counts in fish of economic and indicator importance. *Acta Veterinaria Brno* 67:215-226.
- Moyle, P. B., B. Herbold, D. E. Stevens, and L. W. Miller. 1992. Life history and status of delta smelt in the Sacramento-San Joaquin Estuary, California. *Transactions of the American Fisheries Society* 121:67-77.
- Roger, J. T., J. G. Richards, and C. M. Wood. 2003. Ionregulatory disruption as the toxic mechanism for lead in the rainbow trout (*Oncorhynchus mykiss*). *Aquatic Toxicology* 64:215-234.
- Stine, C. B., A. S. Kane, and A. M. Baya. 2010. Mycobacteria isolated from Chesapeake Bay fish. *Journal of Fish Diseases* 33:39-46.
- Swanson, C., R. C. Mager, S. I. Doroshov, and J. J. Cech. 1996. Use of salts, anesthetics, and polymers to minimize handling and transport mortality in Delta smelt. *Transactions of the American Fisheries Society* 125:326-329.
- Teh, S. J. 2007. Final report of histopathological evaluation of starvation and/or toxic effects on pelagic fishes: pilot study of the health status of 2005 adult Delta smelt in the upper San Francisco Estuary. Department of Anatomy, Physiology, and Cell Biology, School of Veterinary Medicine, University of California, Davis, USA.
- Yasutake, W. T., and J. H. Wales. 1983. Microscopic anatomy of salmonids: an atlas. United States Department of the Interior Resource Publication 150. Washington, D.C., USA.

Received 26 August 2010

Accepted 19 October 2010

Associate Editor was P. Kalvass