

**State of California
The Resources Agency
DEPARTMENT OF FISH AND GAME**

**2006 REPORT
TRINITY RIVER TRIBUTARIES
STEELHEAD SPAWNING SURVEY REPORT**



Prepared by

**Andrew M. Hill
Northern Region
California Department of Fish & Game**

**Anadromous Fisheries Resource Assessment Monitoring Program
May 2008**

TABLE OF CONTENTS

	Page
Abstract	1
Introduction	1
Study Objectives	3
Study Area	4
Methods	7
Sampling Frame	7
Private Property Permission	7
Timing	8
Redd Identification	8
Results	10
Discussion	13
Recommendations	17
Acknowledgements	18
Literature Cited	19

List of Tables

Table 1	Tributary pass schedule	8
Table 2	Redd identification criteria	9
Table 3	Data recorded on each redd	10
Table 4	Trinity River winter-run steelhead spawning survey summary results, March-May 2006	11
Table 5	Redd area measurements by tributary	11
Table 6	Cumulative steelhead redds observed by date, redd type, and temperature	12
Table 7	Summary results of work by D.A. LaFaunce (1964). A steelhead spawning survey of the upper Trinity River system.	15
Table 8	Results of steelhead spawning surveys conducted by D.W. Rogers 1971 and 1972	15

List of Figures

Figure 1	Map showing key geographic locations for the Klamath Mountains Geological Province	2
Figure 2	Map of Trinity basin with selected spawner survey tributaries....	6

List of Appendices

Appendix 1	Reach location and total distance	22
Appendix 2	Redd counts from previous California Department of Fish and Game winter steelhead spawning surveys	23
Appendix 3	Survey beginning and ending water temperatures	23
Appendix 4	Total wild and hatchery steelhead returns to the Trinity River Hatchery	24

**2006 REPORT
TRINITY RIVER TRIBUTARIES
WINTER-RUN STEELHEAD SPAWNING SURVEY REPORT**

Prepared by

**Andrew Hill
Northern Region
California Department of Fish & Game**

ABSTRACT

This report documents the results of spawning surveys conducted by the California Department of Fish and Game on selected Trinity River tributaries from March through May of 2006. This is a continuation of spawning surveys on selected tributaries and serves to create an index of spawning steelhead abundance by enumerating redds. Between March and May 2006, we observed a total of 22 redds in 23.8 kilometers of surveyed habitat. Overall redd density for all tributaries surveyed was 0.92 redds/kilometer. The highest redd density occurred in East Fork of Hayfork Creek (2.65 redds/km), while zero redds were observed in four tributaries: South Fork Indian, Soldier, Deadwood, and Rusch Creeks. During the course of the surveys, crews encountered 16 adult steelhead, in only Deadwood and East Fork of Hayfork Creeks.

INTRODUCTION

The current state of knowledge regarding steelhead (*Oncorhynchus mykiss*) spawning in the Trinity basin is limited. Steelhead found within the Trinity basin have been classified by the National Marine Fisheries Service as the Klamath Mountains Province (KMP) Steelhead based on geologic boundaries. These boundaries extend from the Cape Blanco area in southern Oregon to the Klamath River Basin in northern California (Figures 1) (Busby et al. 1994). This includes the Klamath, Trinity, Rogue, Elk, and Smith River Basins. In 2001, the KMP Steelhead were determined not warranted for listing by the Endangered Species Act (ESA) due to the fact they are not a separate evolutionarily significant unit (ESU) of the steelhead species. An ESU is a “reproductively isolated population” which “represents an important component in the evolutionary legacy of the species” (Waples 1991). The determination for this ESU was made by combining the winter populations, fall populations, summer populations and nonanadromous populations within the KMP boundaries. The summer populations were labeled depressed, but the winter populations were labeled healthy and the ESA listing was deferred.




Figure 1. Map showing key geographic locations for the Klamath Mountains Geologic Province (modified from Irwin 1966 and Walker & MacLeod 1991).

Most prior spawner surveys within the Klamath Mountains Province concentrated on salmon and were therefore terminated prior to steelhead spawning. Prior surveys have been conducted on main-stem Trinity River tributaries in 1964, 1971, and 1972 to monitor the effect of Lewiston Dam on steelhead populations. Most recently, steelhead spawning surveys were conducted in South Fork Trinity River tributaries in 1990 - 1995 under the California Department Fish and Game's Trinity River Project. This is the sixth year of spawning surveys conducted by the Anadromous Fisheries Monitoring Assessment Program (formerly the Steelhead Research and Monitoring Program) on

selected Trinity River tributaries which started in 2000. Traditional basin-wide estimates of steelhead abundance provide little information on the distribution of steelhead spawning. Surveys conducted to enumerate successful steelhead spawning and habitat utilization in tributaries will help to assess this critical component of life history.

Steelhead in the Trinity basin can be split into three races based upon spatial and temporal segregation: summer, fall, and winter which are all included in the same ESU (Busby et al. 1994). Summer-run fish enter freshwater in April through September and over-summer in deep pools prior to entering smaller tributary streams during the first November rain. They continue to migrate upstream through January, and spawn in January and February. (Barnhardt, 1986). Fall-run fish, referred to as summer run-B in systems such as the Rogue, enter freshwater in September and October and spawn from January through April (Currier, personal communication). Winter-run steelhead enter the mouth of the Klamath and migrate upstream from November 1st through April 30th (Barnhardt, 1986). Winter-run steelhead spawning begins in early March and continues through May (Fukushima and Lesh, 1998). Historically, Moffitt and Smith (1950) observed, prior to the completion of Trinity Dam, that spawning of winter-run steelhead began in the upper Trinity drainage in the last part of February, peaking in late March and early April, with some scattered spawning continuing through early June. Previous spawning surveys of Trinity tributaries by the Department of Fish and Game from 2000-2005 showed that spawning in main-stem tributaries peaked by April 1st, approximately two to three weeks prior to peaks in the South Fork basin in April (Garrison, 2002).

Study Objectives

1. Quantify the number of steelhead redds in selected tributaries.
2. Assess spawning habitat conditions.
3. Create index for future comparison of redd numbers. Selected tributaries are included in future surveys for comparison and possible trend analysis.
4. Determine temporal and spatial spawning distribution of steelhead in Trinity River tributaries.
5. Verify and assess barriers to steelhead migration on surveyed tributaries.

Study Area

The area covered by these spawning surveys includes all anadromous tributaries of the Trinity basin upstream of the New River, including the South Fork of the Trinity River (Figure 2). A stratified random sampling design was used to select tributaries within the basin. To develop a sampling universe, all anadromous tributaries within the named basins were identified. The entire basin was then stratified into two sub-basins, the South Fork and the main-stem, each of which was sampled approximately evenly. The following eight Trinity River tributaries were surveyed from their confluence to an upstream migrational barrier except where noted.

South Fork Trinity River sub-basin tributaries

Smoky Creek was surveyed from the South Fork Trinity River confluence to a waterfall barrier 4.1 km upstream. Access is only available through private property owned by Jon Ostrat near Silver Creek. Smoky Creek's remoteness has spared it from most recent land management activities, leaving the creek in a rather pristine condition. Stream condition inventories of Smoky Creek by the US Forest Service in 1998 stated the quality and quantity of gravel in Smoky Creek was not found to be a critically limiting factor to steelhead, although high redd gravel embeddedness values could impede fry emergence. Overall, they concluded that the Smoky Creek watershed is one of the premier tributary systems of the South Fork Trinity River, with ample salmonid habitat, cool water temperatures, relatively low sediment recruitment and an ample riparian and structural cover component (Garrison, 1999).

Eltapom Creek was surveyed from the South Fork Trinity River confluence to a waterfall barrier 1.26 km upstream. Access is only available by crossing the South Fork Trinity River (SFTR), off of Forest Highway 311. A raft is recommended and sometimes necessary for crossing the SFTR at higher flows, especially in March and early April. Eltapom Creek is often referred to as the gem of the South Fork Trinity River; it has excellent spawning gravel, sufficient holding pools, and a dense riparian corridor. Although very short in length, it consistently shows high redd densities and fish counts.

East Fork of Hayfork Creek (EF Hayfork) was surveyed from its confluence with Hayfork Creek to Byron Gulch approximately 6.77 km upstream. There is no permanent barrier on EF Hayfork. EF Hayfork has been heavily impacted by historic mining, evidenced by large piles of mine tailings that stand above the channel. Even through much of the boulder/cobble framework needed to retain gravel has been removed, plentiful spawning gravel and suitable habitat flourishes. Major anadromous tributaries to EF Hayfork include Potatoe Creek and North Fork East Fork Hayfork Creek.

Rusch Creek was surveyed from its confluence with Hayfork Creek approximately 2.1 km upstream. From Hayfork, California, the confluence is accessed by taking Tule Creek road to Forest Route 3N08, then right on 2N02 to 32N11 road. Walk down to the confluence from the road and survey up to the shuttle vehicle.

Mainstem Trinity River sub-basin tributaries

Deadwood Creek was surveyed from its confluence with the Trinity River to a waterfall barrier 3.82 km upstream. Access is available from Deadwood Road. Deadwood is the uppermost tributary to the Trinity River below Lewiston Dam. Deadwood Creek has a steep high energy channel in the lower kilometer, which flattens out into a section of sinuous, complex spawning habitat with adequate large wood and a dense riparian corridor. One aesthetic problem is that Deadwood has become a dumping ground for trash, cars, and used appliances; these litter the banks of the creek in several areas, but have not yet led to any perceived or observable acute pollution problems. Five Counties Salmonid Conservation Program has initiated several restoration projects involving sediment control and fish passage since 2005 along Deadwood Creek.

South Fork of Indian Creek (SF Indian) was surveyed from its confluence with Indian Creek to a waterfall barrier 1.49 km upstream. Access is available via Reading Creek Rd. and by an unnamed SPI logging road. SF Indian has a bedrock dominated high energy channel, with no anadromous tributaries. Some spawning habitat is available in the lower reach, but gravel availability in the upper reach is sporadic. SF Indian has an abundance of deep pools and a thick riparian corridor.

Maxwell Creek was surveyed from its confluence with the Trinity to a 8.25 foot waterfall barrier approximately 2.5 km upstream. The creek is most easily accessed by Dutch Creek Road and a short hike over from the Dutch Creek watershed. The majority of Maxwell Creek consists of a steep V-shaped canyon with plentiful riparian vegetation, in-stream cover and moderate quantities of suitable spawning gravels.

Soldier Creek was surveyed from its confluence with the Trinity River to a culvert barrier approximately 2.2 km upstream. The confluence is accessed by taking Dutch Creek road towards Evans Bar and turning left on a dirt road before the culvert. The upstream barrier is accessed by taking Dutch Creek road to Nation Forest route 33N47. Soldier Creek has abundant vegetation growth and a steep stream gradient with little suitable spawning gravels. Five Counties Salmonid Conservation Program has reconstructed two culverts on Soldier Creek for restoring fish passage.




Figure 2. Map of Trinity basin with selected spawner survey tributaries.

METHODS

Sampling Frame/Tributary Selection

The sampling frame was developed by scouring U.S. Forest Service habitat typing files located in the Hayfork and Weaverville Forest Service Fisheries offices. Tributaries located in the Six Rivers National Forest were confirmed with the local Forest Service zone fisheries biologist (L. Morgan, personal communication). Most habitat typing data from the Forest Service is 15-30 years old; some barriers are classified as semi-permanent, i.e. log-jams, short cascade fields. We are currently verifying and expanding our sampling universe when time allows.

The same panel of selected tributaries is surveyed every year, and other tributaries are either randomly selected or selected if restoration work is going to be performed on it. For example, Soldier creek is in process of having two culverts rebuilt for better fish passage, so it was chosen.

Tributaries are surveyed on foot by surveyors walking in the tributary creeks from the confluence of the creek with another water body upstream to the anadromous barrier or the end of the survey section. Surveyors wear polarized glasses and visually survey the stream for steelhead redds. Encountered redds are measured and all data from Table 3 is collected and written on data sheets.

Private Property Permission

Permission to survey across private property is obtained from all landowners prior to any surveys being conducted. Specific parcels to be surveyed across are identified using Trinity County Assessor maps. All landowners are notified by mail and asked to return a postcard allowing the Department permission to survey the named tributary across their property with the condition that crews stay below the high-water mark. Additional permission is ascertained in cases where access to the tributary across a landowners property is necessary. Letters verifying permission are sent out annually in late January or early February. Sierra Pacific Industries (SPI) is the largest private landowner in Trinity County and has been most cooperative in allowing permission on all SPI lands.

Timing

All tributaries are surveyed every three to four weeks from March through May. Main-stem tributaries are surveyed first due to historically earlier spawning when compared to the South Fork basin. Survey reaches are surveyed sequentially from confluence to headwaters whenever possible. Some timing adjustment was necessary due to snow, rain events, and problems with funding for technicians. Table 1 (below) lists pass dates per reach and tributary.

Table 1. Tributary pass schedule.

Tributary	Pass Date				
	Pass 1	Pass 2	Pass 3	Pass 4	Pass 5
Deadwood	3/13/2006	3/28/2006	4/17/2006	4/28/2006	5/12/2006
South Fork Indian	3/24/2006	4/21/2006	5/15/2006		
Smoky	5/16/2006				
East Fork Hayfork/R1	3/21/2006	4/04/2006	4/26/2006	5/09/2006	5/25/2006
East Fork Hayfork/R2	3/22/2006	4/06/2006	4/26/2006	5/10/2006	5/26/2006
Eltapom	5/11/2006	5/24/2006			
Rusch	5/01/2006	5/18/2006			
Soldier	3/23/2006	4/25/2006	5/04/2006	5/23/2006	
Maxwell	3/15/2006	4/20/2006	5/03/2006	5/22/2006	

Redd Identification

Crews are trained in proper redd identification prior to the beginning of the season. Ultimately, an experienced crew leader is present to make all “tough calls” in terms of redd identification. In-experienced technicians often overlook redds or have trouble distinguishing steelhead redds from scour hydraulics or lamprey and resident trout redds. The following criteria (Table 2) is used to insure proper identification of steelhead redds; not all criteria must necessarily be satisfied in order for a redd to be called a redd.

Table 2. Redd identification criteria.

Criteria	Explanation
Location	Most redds are located in pool tail-outs or riffles; Briggs (1953) found that most redds occupied the transitional area between pools and riffles.
Size	Hunter (1973) found the area of average steelhead redds to be 4.4 meter ² , although, redds are often smaller when spawning habitat is limited or constrained by channel morphology.
Structure	Redd should consist of a pit and mound (tail-spill), with the mound downstream of the pit. Steelhead redds can be easily differentiated from lamprey redds, as lamprey redds lack a mound or tail-spill.
Substrate size	Steelhead prefer to spawn in gravel 0.6-10.2 cm in diameter. (Smith, 1973).
Gravel sorted	The substrate of freshly constructed redds is usually well sorted, with larger gravel positioned anterior compared to smaller gravel.
% fines	Redds should not be overly embedded with fine substrate, as the mechanics of redd construction should wash away fine sediment.
Water velocity	There must be adequate velocity to insure oxygenation of eggs. Bovee (1978) found optimum velocity for steelhead spawning at 2 feet/sec.
Pit/tailspill mechanics	Redds should be properly spatially positioned, so that the pit is upstream of the tail-spill and gravel excavated from pit could form tail-spill.
Lack of algae or detritus	New constructed redds should be free of algal formation (i.e. periphyton) and detritus. Detritus often accumulates in the pit of older redds.
Presence of fish on redd	Presence of an actively spawning pair of fish indicates probable construction of a successful redd. Test digging can often be confused for successful completion of a redd.

Table 3 contains the data recorded on all redds encountered during the course of the survey. GPS coordinates were taken using a Garmin 12XL receiver utilizing the NAD 27 datum. All redd measurements were taken using a water-proof tape measure. During measurements, extreme caution was taken to avoid disturbing redds. Redds currently under construction (fish on redd) were not measured at that time, to avoid disturbing spawning. These redds were measured on the subsequent pass. All encountered redds are flagged with date, redd number, position, and recorders initials, to prevent double counting, and to allow future evaluation.

Table 3. Data recorded on each redd.

Data Field	Description
Redd I.D.#	3 digit code with the first digit being the reach no. and the second two being the consecutive redd no. for that reach e.g. R101=reach 1, redd no.1
GPS coordinates	Lat/Long waypoint of redd location
Pit length	Pit length measured parallel to the flow
Pit width	Pit width measured perpendicular to the flow
Depth 1	Depth from substrate to bottom of the pit
Depth 2	Depth from water surface to bottom of the pit
Pit substrate	Dominant substrate in the pit
Tail spill length	Tail spill length measured parallel to the flow
Tail spill width 1	Tail spill width perpendicular to the flow at 1/3 of the distance down from the upstream end
Tail spill substrate	Dominant substrate in the tail spill
Habitat type	Habitat type where redd is located
Redd type	Condition of redd: 1=well defined recently completed 2=well defined but not new 3=not well defined 4=older and difficult to identify, may be questionable
Comments	Redd location description and information on redd condition

RESULTS

This 2006 season, a total of 22 redds were observed in eight tributary creeks during surveys encompassing 23.8 kilometers of habitat. Summary redd and fish observation data by tributary is provided in Table 4, and a summaries of previous year's surveys are provided in Table 7, Table 8, and Appendix 2. Evidence of steelhead spawning was observed in only two creeks; Deadwood and East Fork of Hayfork Creeks. An attempt was made to identify presence or absence of an adipose fin on all live fish observed. Of the 16 steelhead observed during this season's surveys, only 7 (43.7%) could properly be identified as of wild or hatchery origin. In Deadwood Creek, one wild adult steelhead was observed and one hatchery male carcass. This contrasts to East Fork of Hayfork Creek which had five wild adults.

Only one fish was observed actively constructing a redd (1 of 22). This percentage may seem low, but with a low sampling periodicity (about once per month), and assuming it takes three days to construct a redd (Hannon, Healey and Deason, 2003), you would expect to see only 10% of the redds being actively constructed.

Table 4. Trinity River winter-run steelhead spawning survey summary results, March-May 2006.

Tributary	Mileage (km)	Redds	Redds/km	Adult Steelhead
Smoky	4.1	1	0.24	0
East Fork Hayfork	6.8	18	2.65	12
South Fork Indian	1.5	0	0	0
Maxwell	2.5	1	0.40	0
Soldier	1.7	0	0	0
Rusch	2.1	0	0	0
Eltapom	1.3	2	1.5	0
Deadwood	3.8	0	0	4
Totals	23.8	22	0.92	16

Redd location was characterized by habitat type for all redds observed. Not surprisingly, steelhead preferred to spawn in pool tail-outs (50%), followed in preference by runs (27%), and riffles (23%). No redds were located in other types of habitat, most likely due to the need for suitable flow. The average depth of water over the pit for all redds observed was 13.1 inches (33.27cm). The average depth of excavation (material removed from pit to create tail-spill) was 5.1 inches (12.96 cm).

All redds were measured utilizing methods put forth by Gallagher (2002). Table 5 (below) provides summary area measurements for all redds observed by tributary. Overall, the average total area of all redds observed during the survey was 12.9 ft². The smallest redd observed during the survey occurred on Eltapom Creek in a predominantly boulder/ gravel pool tail and measured 7.92 ft². The largest redd was also observed in East Fork of Hayfork Creek and measured 27.88 ft².

Table 5. Redd area measurements by tributary.

Tributary	N=	Mean pit area (ft²)	Mean tailspill Area (ft²)	Mean total redd area (ft²)	Mean pit depth in substrate (inches)	Mean pit depth in water (inches)	Dominant redd location
E.F. Hayfork	18	9.21765	9.92555	19.14355	7.1	12.5	Pool tail
Eltapom	2	4.41667	3.75000	8.16667	5.5	21.0	Run/ Pool tail
Smoky	1	4.87500	6.61111	11.48611	2.0	9.0	Run
Maxwell	1	2.33333	10.27778	12.61111	8.0	12.0	Riffle
Overall	22	5.2107	7.641	12.8511	5.1	13.1	Pool tail

With so few redds observed it's difficult to say much about the main-stem spawning peaks, but no redds were observed in main-stem tributaries after March 15, 2006 (Table 6). Spawning in the Hayfork basin peaked by the end of March and no new redds were observed after April 6, 2006 (Table 6). Surveys could not be done in March or April in South Fork Trinity River tributaries until May, but one out of the three redds observed was redd type 1. This indicates that the spawning is later in South Fork Trinity River tributaries and they were probably finished all spawning activity by the time surveyors were able to access the creeks. Typically, main-stem Trinity River tributary spawning peaks are before those in the South Fork Trinity River tributaries (Garrison 2005).

Table 6. Cumulative steelhead redds observed by date, redd type, and temperature (Celsius)

Creek	Drainage basin	Date	Number of redds	Redd type	Average survey temperature
Maxwell	Main-stem Trinity River	3/15	1	1	5
E.F. Hayfork	Hayfork Creek, South Fork Trinity	3/21	8	1	3
E.F. Hayfork	Hayfork Creek, South Fork Trinity	3/22	7	1	5
E.F. Hayfork	Hayfork Creek, South Fork Trinity	4/4	1	1	3.5
E.F. Hayfork	Hayfork Creek, South Fork Trinity	4/6	2	1	3.5
Eltapom	South Fork Trinity	5/11	2	3.5	7.5
Smoky	South Fork Trinity	5/16	1	1	9.5

Water temperature was recorded at the beginning and end of each survey. Temperatures during peak spawning activity this year were between 2°C to 6°C. Daily recorded temperatures for each tributary surveyed are listed in Appendix 3. Temperatures ranged from 2°C on a cold morning in March, to 11°C on a May afternoon. Redds in all tributaries except for Eltapom were classified as redd type 1, and all redds were discovered when average survey temperatures were less than 10°C.

DISCUSSION

Redd surveys serve as a good, but partially incomplete means of monitoring steelhead spawning escapement. These surveys are most appropriate when other means of estimating adult escapement or spawning success are not appropriate or impossible to conduct. In the Trinity basin, problems do occur which limit the ability to estimate the abundance of winter-run steelhead. High flows and the extended length of adult steelhead migration make weir estimates partial at best. Several weirs were constructed to estimate winter-run steelhead run-size by the Department in 1986-1996 at Sandy Bar and at Forest Glen on the South Fork of the Trinity River (CDFG, 1990-95). Efforts were finally terminated after multiple blow-outs due to high flows.

The availability of spawning habitat could possibly inhibit spawning and resulting production of steelhead, especially in smaller order tributaries, where gravel availability is scarce. The amount of suitable stream substrate for spawning varies with the size (order) of the stream and species of salmonid using it, as Boehne and House (1983) learned from their study of two coastal and two Cascade Range watersheds in Oregon. First and second order streams were rarely used by anadromous salmonids; the larger anadromous steelhead, Coho, and Chinook salmon spawned in a few third-order streams, but most were found in fourth- and fifth-order streams. As stream order increased, gradient decreased but stream length, width, and depth increased. The amount of spawning gravel per kilometer of stream was greatest in fourth order coastal watersheds and fifth order Cascade Range watersheds. Precursory examinations of gravel in surveyed tributaries have found that in the Trinity basin gravel retention could be impeded by the following factors: effects of historic mining and the 1964 flood. Retention of gravel is often problematic, even in fourth- and fifth-order streams due to incision of the channel as a result of historic mining. This incision causes loss of channel sinuosity, and increases channel energy, especially during high flow events. Without the complexity associated with a sinuous channel, little large wood or boulder/cobble framework is available to sort and retain gravels. Further complicating the problem is the long lasting effects of the 1964 flood; this 100-year flood aggraded spawning beds up to eight feet deep with unsuitable substrate; most tributaries continue to down-cut through this aggregation, and have yet to reach channel equilibrium 40 years later.

No apparent trend is evident when examining redd survey data collected during previous field seasons. Some tributary creeks appear to annually fluctuate more than others, and some appear to show a trend of declining numbers of redds since the 1990's (Appendix 2). 2002 redd surveys document the highest numbers of redds than other years. The best data possible trend data is from the three creeks which were surveyed every one of these years; East Fork of Hayfork Creek, Eltapom, and Smoky Creeks. East Fork of Hayfork creek steelhead redd counts fluctuate the most from 64 redds in 2002 to 0 redds in year 2000 with no upward or downward trend. Both Eltapom and Smoky Creek fluctuate annually with a downward trend in redd counts.

Studies conducted in Washington and Idaho have both approximated average steelhead redd area at 47ft² (Hunter 1973, Reiser and White 1981). Gallagher and Gallagher in 2005 found steelhead redds in several anadromous streams in Mendocino County to average 19.2ft² with a standard deviation of 1.5 ft². Redds in Trinity basin appear to be smaller than those constructed by steelhead elsewhere which may be an indication that wild steelhead in the upper Trinity River basin may be smaller than those closer to the ocean. Different substrate compositions in the Trinity Basin compared to coastal watersheds may also account for this difference in redd size. This year the average redd area measured during these surveys was 12.85 ft². This is smaller than previous years' measurements of 23.63ft² in 2005 and 13.78 ft² in 2004. These are all smaller than those in Washington and Idaho and may be a result of smaller fish sizes, but the data is not available to make that statement.

Patrick Higgins (personnel communication, 2000) hypothesizes that South Fork Trinity River steelhead begin spawning when water temperatures approach 8⁰C. Hunter (1973) states the range of preferred temperatures for steelhead spawning at 3.9-9.4⁰C. The results of this years' survey support Hunter's statement where most all spawning in the Trinity had been completed prior to water temperatures reaching 10⁰C (Table 6). 100% of the redds discovered this year were discovered when average survey temperatures were less than 10⁰C.

Steelhead run size is highly variable from year to year. Redd surveys during 2006 season documented a below average year for Trinity basin steelhead. For comparison, work by D.A. La Faunce in 1964 and D.W. Rogers in 1971 and 1972 set up base-line numbers for natural production of steelhead in the Trinity basin (Table 7&8). Those surveys show that adult steelhead estimates were markedly higher in 1964 than in any of the following years. However, these numbers could also be biased due to the construction of the Trinity Dam and displacement of the returning steelhead from their historic spawning grounds. The dam may have forced the steelhead to exploit new tributaries and produce higher than average redd counts that year. Survey results by D.W. Rogers are more similar to results obtained by these surveys. This may indicate that after the completion of dam work on the Trinity River, the steelhead populations have now reached their carrying capacities in the Trinity River basin at reduced population sizes than before dam construction.

Table 7. Summary results of work by D.A. LaFaunce (1964). A steelhead spawning survey of the upper Trinity River system.

Tributary surveyed	Distance surveyed (km.)	Redds observed	Redds/km.
Deadwood Creek	1.66	27	16.26
S.F. Indian Creek	0.37	4	10.80
Soldier Creek	1.70	21	35.70
Maxwell Creek	0.34	6	2.04

Table 8. Results of steelhead spawning surveys conducted by D.W. Rogers (1971).

Tributary surveyed	Distance surveyed (km.) 1971	Redds observed 1971	Redds/km. 1971	Distance surveyed (km.) 1972	Redds observed 1972	Redds/km. 1972
Deadwood Creek	3.7	0	0	3.7	0	0
S.F. Indian Creek	1.85	3	1.62	0.85	0	0
Soldier Creek	2.72	1	0.37	No survey	No survey	No survey
Maxwell Creek	1.53	1	0.65	No survey	No survey	No survey

Results of this spawning survey have important fisheries management implications; fisheries managers use escapement data to analyze the ability of a stock to sustain recreational fisheries. Proposals have recently been submitted to the Fish and Game Commission to increase the bag limit on the Trinity and to allow the take of wild steelhead. The Department must make management recommendations based upon the best available science documenting the status of the steelhead in the basin. The results of this project are currently the only data available examining population status and trends of winter-run steelhead in the Trinity Basin. The Department used this justification to recommend not allowing the take of wild steelhead on the Trinity River. Should spawning surveys show a prolonged increase in escapement throughout the basin, additional fisheries and decreased angling restrictions (increased opportunities) could be considered.

Problems are commonplace and often complicate and sometimes prevent redd surveys from occurring. Possible problems include adequate survey frequency; redd

discrimination by species, tributary sample selection, access, and private property permission. Some of these problems create bias within the data, while others prevent the proper coverage of a selected tributary.

One primary problem that affects sample design, as well as proper and even coverage, is access. Most of the Trinity basin is composed of rugged mountainous terrain with little road coverage. Existing roads are often poorly maintained logging roads, which rarely lead to the confluence of a selected tributary. Some tributaries lie within wilderness areas, where no roads exist, and hiking in to survey is the only possibility. Access problems are further compounded by extreme winter conditions such as snow storms and high river flows. Some tributaries, such as Smoky Creek, in South Fork Trinity basin are inaccessible by road until late April due to heavy snow-pack. High flows in the South Fork Trinity River can also limit access across the river to tributaries such as Eltapom Creek.

High flows often have a negative effect on survey periodicity and quality. These high flows often prevent surveys by limiting travel through the stream corridor and impeding visibility with whitewater. Rain further impedes a surveyor's ability to detect redds by breaking the smooth surface of the water, making underwater terrain features nearly invisible. High flow events during the spring survey can also scour redds making them indiscernible by the time surveyors can get to them. The spring of 2006 had high volume of rain and therefore high flows in April and May which had a negative effect on the numbers of redds observed.

Discrimination of redds created by different fish species is a problem which often complicates redd surveys in systems where several species of fish co-exist and spawn during similar time frames. Several fish species temporally co-exist in the Trinity basin, several of which have similar spawning time frames; coho salmon (*Oncorhynchus kisutch*) enter the watershed in November and December and spawn in January and February. Similarly, Pacific Lamprey (*Lampetra tridentata*) migrate into the system in the fall and winter, and spawn during the spring months. Small trout exhibiting a resident life-history also co-exist in the system and spawn during the spring. Several measures are taken by crews to ensure proper classification of steelhead redds. All redds with no substantial tail-spill or a pit diameter of less than 12 inches are considered resident trout or lamprey redds. Resident trout tend to utilize smaller substrate in areas with less apparent velocity. Lamprey redds are distinguished by a small circular pit and no tail-spill, but could be identified as steelhead redds by inexperienced surveyors as witnessed by D.W. Rogers in 1971. In the Trinity basin, coho redds are infrequently confused with steelhead redds due to their earlier spawning (January/February vs. March-May).

Surveys utilizing multiple technicians inherently suffer from problems with inter-observer variation, both with observer efficiency and the subjective nature of identifying redds. This can be minimized by pairing experienced with inexperienced technicians, sufficient training, and frequent quality control trials. This is currently not recognized to

be a problem, as all crews contain at least one experienced member.

Prior to the beginning of the first season in 2000, permission from private property owners was obtained on all tributaries to be surveyed. Tributaries with excessive refusal of landowners were dropped from the sampling frame. For long term management, streams entirely located on public property are chosen over those with multiple private ownerships. Four of the eight tributaries selected for the 2006 survey are located entirely on public land. Two of the eight tributaries are located on Southern Pacific Industries land. SPI has been very cooperative in allowing the department to survey on their land. Access to the remaining two tributaries was obtained by sending letters to the owners asking for permission to survey on their land.

RECOMMENDATIONS

The limits of the anadromy in the Trinity basin need to be verified in order to properly delineate the sampling universe and quantify habitat available to steelhead. Previously, an effort to identify barriers to anadromy was completed by Trinity Fisheries Consulting, but focused solely on road induced barriers (Trinity Fisheries Consulting, 1988). Most other barrier information relies on antiquated US Forest Service habitat typing files, produced in the late 1960s and 1970s.

Coho salmon spawner surveys should be initiated to complement existing steelhead surveys. Few surveys quantifying Coho salmon spawning are currently conducted in the Trinity basin, with the only effort being made the US Forest Service, and only when funding for fisheries technicians allows (L. Everest, personal communication).

More intensive habitat evaluation needs to be included in the survey design. A quick, one pass field extensive evaluation should be made of quantity and quality of available spawning substrate in selected tributaries. Evaluation of spawning habitat was conducted previously by Fish and Game from 1994-1997 on South Fork Trinity tributaries (Borok and Jong, 1997). This effort should be expanded to include the entire sampling universe using a protocol similar to that put forward by Schuett-Hames and Pleus (1996).

Patrick Garrison has recommended that some examination be attempted to look at the relationship between channel incision, removal of boulder framework, historical mining, and availability of spawning habitat (Garrison, 2004). Historical mining in the late 1800s and early 1900s has resulted in the removal of bed-load framework necessary for the retention of suitable spawning gravel. Removal of this framework has further resulted in channel incision, loss of channel sinuosity, and loss of habitat complexity. Evidence of these effects is made apparent by the large piles of mine tailings covering the banks of over half of all Trinity tributaries; some of the piles are over 50 feet high and several

hundred feet wide.

All redds should be evaluated with each pass as season progresses to assess sufficient periodicity of surveys. Last year, all redds were evaluated during the last pass to determine if survey periodicity was sufficient. During the 2006 field season, all redds in creeks which had multiple passes were no longer visible redds by the last pass. It would be informative to have all redds be re-evaluated during each subsequent passes to evaluate whether current survey periodicity is sufficient. Currently it appears that we are sampling longer than needed to evaluate spawning activity.

A more intensive effort should be undertaken to understand the relationship between the number of adult steelhead and the corresponding number of redds. The Oregon Department of Fish and Wildlife has five years of data in the Alsea and Nestucca basins that shows a strong relationship between redd counts and fish numbers ($R^2=0.97$, $p<0.001$). Using this regression as a calibration between adults and redds, they further suggest that redd counts are a good indicator of population size over a range of run-sizes from 35 – 2,131 fish (Susac and Jacobs, 2003).

A reliable funding source needs to be identified to insure that everything is fully operation by March 1st, and that funding for technicians proceeds unfettered through the completion of the season. This should insure a more even sampling effort that spans the entire season.

ACKNOWLEDGEMENTS

A special thanks goes out to all those who help make this project and report possible, most importantly Patrick Garrison. Without his dedication to steelhead in supervising this survey there would be no report. Also, thank you to all of the private property owners, whose cooperation with us is greatly appreciated.

LITERATURE CITED

- Barnhardt, R.A. 1986. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates: steelhead. U.S. Fish and Wildlife Service Biological Report 82. 21 pp.
- Bell, M.C. 1986. Fisheries handbook of engineering requirements and biological criteria. Fish Passage Development and Evaluation Program. US Army Corps of Engineers, North Pacific Division. Portland, OR. 290 pp.
- Boehme, P.L., and R.A. House. 1983. Stream ordering: a tool for land managers to classify western Oregon streams. U.S. Bureau of Land Management, Technical Note OR-3, Portland, Oregon.
- Borok, S. L. and H.W. Jong. 1997. Evaluation of salmon and steelhead spawning habitat quality in the South Fork Trinity River basin, 1997. California Department of Fish and Game. Inland Fisheries Administrative Report No. 97-8. 14 pp.
- Bovee, K.D. 1978. Probability-of-use criteria for the family Salmonidae. In-stream Flow Informational Paper 4. U.S. Fish and Wildlife Service. FWS/OBS-78/07. 79 pp.
- Briggs, J.C. 1953. The behavior and reproduction of salmonid fishes in a small stream. California Fish and Game Bulletin No. 94. 62 pp.
- Busby, P.J., T.C. Wainwright, and R.S. Waples. 1994. Status Review for Klamath Mountains Province Steelhead. NOAA Technical Memorandum NMFS-NWFSC-19
- Carroll, E.W. 1984. An evaluation of steelhead trout and in-stream structures in a California intermittent stream. M.S. Thesis, Humboldt State University, Arcata, CA. 51 pp.
- CDFG. 1991-1996. Annual Reports Trinity Basin Salmon and Steelhead Monitoring Project 1989-1995 seasons. Inland Fisheries Division.
- CDFG. 1998. Strategic plan for management of Klamath Mountains Province steelhead trout. Prepared for the National Marine Fisheries Service by the Resources Agency, 15 pp.
- Elms-Cockrum, T.J. 1997. Salmon spawning ground surveys, 1996. Idaho Department of Fish and Game Report 97-25, Boise, Idaho.
- Fukushima, L., and E.W. Lesh. 1998. Adult and juvenile salmonid migration timing in

- California streams. California Fish and Game 84(3): 133-145.
- Gallagher, S. P. 2001. Results of the 2000-2001 Coho Salmon (*Oncorhynchus kitsch*) and Steelhead (*Oncorhynchus mykiss*) spawning surveys on the Noyo River, California. California State Department of Fish and Game, 1031 South Main, Suite A, Fort Bragg, CA 95437. Draft November 2001. 45pp.
- Gallagher, S.P. 2002. Salmonid spawning survey protocols for 2002-2003. California Department of Fish and Game. Draft. 14 pp.
- Garrison, P.S. 1999. Stream condition inventory report, Somky Creek 1998. Shasta-Trinity National Forest, South Fork management Unit. 21 pp.
- Garrison, P.S. 2002. Trinity River tributaries Winter-run steelhead spawning surveys, 2002 season. Project 1d1. California Department of Fish and Game, P.O. Box 1185, Weaverville, CA 96093. November 2002. 20 pp.
- Garrison, P.S. 2004. Trinity River tributaries Winter-run steelhead spawning surveys, 2003/2004 season. Project 1d1. California Department of Fish and Game, P.O. Box 1185, Weaverville, CA 96093. June 2004. 30 pp.
- Hannon, J. 2000. Compilation of yearly steelhead escapement index counts in six Prince of Wales Island streams, 1994-2000. U.S. Forest Service, Tongass National Forest. Craig, AK.
- Hannon, J., M. Healey, and B. Deason. 2003. American River steelhead (*Oncorhynchus mykiss*) spawning 2001-2003. U.S. Bureau of Reclamation and California Department of Fish and Game. 30 pp.
- Hunter, J.W. 1973. A discussion of game fish in the State of Washington as related to water requirements. Report by the Washington State Department of Game, Fishery Management Division, to the Washington State Department of Ecology, Olympia.
- Irwin, W.P. 1966. Geology of the Klamath Mountains Province. In E.H. Bailey (editor), Geology of northern California, p. 19-38. U.S. Geological Survey, California Division of Mines and Geology, Bulletin 190.
- LaFaunce, D.A. 1964. A Steelhead Spawning Survey of the Upper Trinity River System, 1964. Marine Resources Administrative Report No. 65-4. California Department of Fish and Game, Region 1, Inland Fisheries Branch.
- Moffett, J.W., and S.H. Smith. 1950. Biological Investigations of Fishery Resources of the Trinity River, CA. U.S. Fish and Wildlife Service Special Scientific Report: Fisheries No. 12. 70 pp.

- Riemen, B.E., and D.L. Myers. 1997. Use of redd counts to detect trends in bull trout (*Salvelinus confluentus*) populations. *Conservation Biology* 11:1015-1018.
- Reiser D.W., and T.C. Bjornn. 1979. Habitat requirements of anadromous salmonids. USDA Forest Service, Pacific Northwest Forest and Range Experiment Station, Portland, OR. General Technical Report PNW-96. 54 pp.
- Reiser, D. W., and R. T. Peacock. 1985. A technique for assessing upstream fish passage problems at small-scale hydropower developments. Pages 423-432 in F.W. Olson, R.G. White, and R.H. Hamre, editors. Symposium on small hydropower and fisheries. American Fisheries Society. Bethesda, MD.
- Reiser, D.w., and R.G. White. 1983. Effects of flow fluctuation and redd dewatering on salmonid embryo development and fry quality. Idaho Water and Energy Resources Research Institute, Research Technical Completion Report, Contract DE-AC79-79BP10848, Moscow, ID.
- Rogers, D.W. 1971. A Steelhead Spawning Survey of the Tributaries of the Upper Trinity River and Upper Hayfork Creek Drainage. California Department of Fish and Game, Region 1, Inland Fisheries Branch.
- Rogers, D.W. 1972. A Steelhead Spawning Survey of the Tributaries of the Upper Trinity River and Upper Hayfork Creek Drainage. California Department of Fish and Game, Region 1, Inland Fisheries Branch.
- Susac, G.L., and S.E. Jacobs. 2003. Assessment of the status of Nestucca and Alsea River winter steelhead, 2002. Oregon Department of Fish and Wildlife. Sport fish and Wildlife Restoration Projects F-181-D-03 and F-1662-R-07. 14 pp.
- Schuett-Hames, D. and A. Pleus. 1996. Literature review and monitoring recommendations for salmonid spawning habitat availability. Northwest Indian Fisheries Commission. 32 pp.
- Smith, A.K. 1973. Development and application of spawning velocity and depth criteria for Oregon salmonids. *Transactions of the American Fisheries Society* 102:312-316.
- Trinity Fisheries Consulting. 1988. Analysis of road induced fish migration barriers in the Trinity River basin, a stream crossing inventory. Prepared for the California Department of Fish and Game, Inland Fisheries Division. Standard Agreement No. C-1927. 144 pp.
- Walker, G.W. and N.S. MacLeod. 1991. Geologic map of Oregon: Geomorphic divisions Of Oregon. U.S. Dept. Inter., U.S. Geological Survey

Waples, R.S. 1991. Pacific salmon, *Oncorhynchus* spp., and the definition of “species” under the Endangered Species Act. *Marine Fisheries Review* 53(3): 11-12.

Personal Communications

Loren Everest, December 2003. US Forest Service, Trinity River Management Unit.

Monty Currier, December 2004. California Department of Fish and Game.

Appendix 1. Reach location and total distance.

Stream/ Reach	Start Latitude/Longitude	End Latitude/Longitude	Reach distance (km)	Hike distance (km)	Total Distance (km)
Eltapom	N40°39.656 W123°29.680	N40°39.559 W123°29.084	1.3	1.3	2.6
Soldier	N40°41.355 W123°01.688	N40°41.353 W123°03.628	1.7	no hike	1.7
Maxwell	N40°39.966 W123°00.502	N40°39.031 W123°01.055	2.5	3.4	5.9
Rusch	N40°34.670 W123°16.804	N40°35.210 W123°15.910	2.1	0.25	2.6
East Fork Hayfork R1	N40°29.335 W123°04.142	N40°30.136/ W123°02.144	3.2	no hike	4.2
East Fork Hayfork R2	N40°30.139 W123°02.070	N40°30.564 W122°59.609	3.7	no hike	4.7
Smoky R1	N40°18.869/ W123°12.136	N40°19.908 W123°11.575	3.8	8.2	12
Smoky R2	N40°28.681 W123°01.684	N40°30.198 W123°02.521	3.5	7.3	10.8
South Fork Indian	N40°35.807 W122°49.892	N40°35.289 W122°49.333	1.5	2.3	3.8
Deadwood	N40°43.089 W122°48.091	N40°42.847 W122°45.826	3.8	no hike	3.8

Appendix 2: Redd counts from previous California Department of Fish and Game winter steelhead spawning surveys.

Tributary	1990	1991	1992	1993	1994	1995	2000	2001	2002	2003	2004	2005
Deadwood	-	-	-	-	-	-	0	2	17	14	22	6
South Fork Indian	-	-	-	-	-	-	0	2	7	2	4	0
Smoky	16	12	14	5	5	5	10	3	9	8	8	3
East Fork Hayfork	32	20	4	3	16	2	0	10	64	16	37	13
Rusch	-	-	-	-	-	-	-	-	11	-	-	-
Eltapom	18	8	13	18	25	3	11	5	11	5	17	2

Appendix 3. Survey beginning and ending water temperatures.

Stream Name	Reach	Date	Start Water Temperature (C [□])	End Temperature (C [□])
Soldier Creek	1	3/23	5	5
Soldier Creek	1	4/25	6	7
Soldier Creek	1	5/4	6	7
Soldier Creek	1	5/23	7	8
SF Indian Creek	1	3/24	4	4
SF Indian Creek	1	4/21	5	6
SF Indian Creek	1	5/15	7	8
Deadwood	1&2	3/13	4	4
Deadwood	1&2	3/28	3	4
Deadwood	1&2	4/17	4	6
Deadwood	1&2	4/28	8	7
Deadwood	1&2	5/12	7	8
Rusch	1	5/1	6	8
Rusch	1	5/18	9	10
EF Hayfork	1	3/21	2	4
EF Hayfork	2	3/22	6	4
EF Hayfork	1	4/4	3	4
EF Hayfork	2	4/6	3	4
EF Hayfork	1	4/26	4	7
EF Hayfork	2	4/26	5	7
EF Hayfork	1	5/9	6	8
EF Hayfork	2	5/10	6	7
EF Hayfork	1	5/25	7	8
EF Hayfork	2	5/26	6	5
Smoky	1	5/16	8	11
Eltapom	1	5/11	7	8
Eltapom	1	5/24	8	7
Maxwell	1	3/15	5	5
Maxwell	1	4/20	6	8
Maxwell	1	5/3	7	8

Appendix 4. Total wild and hatchery steelhead returns to the Trinity River Hatchery

Steelhead Returns at the hatchery							
Year	Steelhead	Year	Steelhead	Year	Steelhead	Year	Steelhead
1960	2,071	1977	285	1990	930	2003	10,224
1961	3,526	1978	683	1991	446	2004	5,725
1962	3,243	1979	382	1992	455	2005	8,143
1963	1,687	1980	2,005	1993	885	2006	11,547
1964	894	1981	1,004	1994	411		
1965	6,941	1982	713	1995	705		
1966	992	1983	599	1996	4,012		
1967	135	1984	142	1997	429		
1968	232	1985	461	1998	441		
1969	554	1986	3,780	1999	1,571		
1970	241	1987	3,007	2000	768		
1971	67	1988	817	2001	2,333		
1972	242	1989	4,765	2002	6,008		