

2012 Vegetation Map in Support of the Desert Renewable Energy Conservation Plan

Interim Report (1.1)



By
Vegetation Classification and Mapping Program
California Department of Fish and Game
in collaboration with
Aerial Information Systems, Inc.

For the
Desert Renewable Energy Conservation Planning Program and
California Energy Commission



June 2012

Project Staff

Vegetation Classification and Mapping Program (VegCAMP)

Jim Alford
Rachelle Boul
Mary Jo Colletti
Diana Hickson
Todd Keeler-Wolf
Anne Klein
Aicha Ougzin
Gina Radieve
Cynthia Roye
Rosalie Yacoub

Aerial Information Systems (AIS)

Lisa Cotterman
John Fulton
Arin Glass
Anne Hepburn
Ben Johnson
Debbie Johnson
John Menke
Lisa Morse
Mike Nelson
Ed Reyes
Janet Reyes

Acknowledgments:

Funding for this project was provided by:

California Energy Commission Public Interest Energy Research (PIER) Program
US Bureau of Land Management
California Wildlife Conservation Board
California Department of Fish and Game

This project could not have been accomplished without the assistance of many people. Among others, we thank the following public land managers and resource experts for permitting and facilitating our field data collection visits and for help with the project overall:

California Department of Fish and Game:

Region 6 - Inland Deserts Region: Kim Nicol, Jeff Brandt, Katrina Banda,
Patricia Jimenez, Natasha Krupnak, Gabbi Gatchel, Angela Salcedo
Region 5 - South Coast Region: Mary Meyer, Karen Miner
Biodiversity Data Branch: Kristi Lazar
Habitat Conservation Planning Branch: Junko Hoshi

California Department of Food and Agriculture: Fred Hrusa

California Department of Parks and Recreation: Carrie Bemis

California Energy Commission: Scott Flint, Misa Milliron, Paul Richens, Kristi Chew

Kern County Department of Parks and Recreation: John Fedorsin

Mojave Desert Manager's Group: Fon Duke

National Aeronautics and Space Administration, Goldstone Deep Space Communications
Complex: April Holmgren, Geoffery Hewitt

National Park Service: Tasha LaDoux

Saint Mary's College: Phil Leitner

Tejon Ranch Conservancy: Mike White

US Bureau of Land Management, Barstow Field Office: Anthony Chavez
Ridgecrest Field Office: Marty Dickes, Glenn Harris, Hector Villalobos,
Matt Duarte (SCA Conservation Corps);
California State Office: Christina Lund

US Department of Defense:

Edwards Air Force Base: Mark Bratton, Dan Reinke
Fort Irwin National Training Center: Liana Aker, Clarence Everly, Dave Housman,
Mark Hessing, George Peck
Naval Air Weapons Station China Lake: Tom Campbell, Tim Ludwick, Anna-Maria Easley

US Forest Service, Angeles National Forest: Bob Blount, Steven Goldschmidt

We apologize to anyone we may have left off of this list.

Table of Contents

Abstract	1
Purpose	1
Classification	1
Delineation Rules	3
Reconnaissance	3
Accuracy Assessment	3
Portion of Map Visited	7
Map Attributes	7
Literature cited	14
Appendix A - Rapid Assessment Field Form & Protocol	
Appendix B - Classification Hierarchy	
Appendix C - Hierarchical Field and Mapping Key	
Appendix D - Land Use Mapping Criteria	
Appendix E - Reconnaissance Field Form	
Appendix F - Field Verification Form	

List of Figures

Figure 1: Study area boundaries	2
--	---

Abstract

The California Department of Fish and Game (CDFG) Vegetation Classification and Mapping Program (VegCAMP) and contractor Aerial Information Systems (AIS) created a fine-scale vegetation map of a portion of the western Mojave Desert in California. The mapped area is bounded to the west and south by USDA Ecoregional Subsection 322Ag of the Mojave Desert (Miles and Goudey 1997). To the east, the portion mapped by AIS (approx. 4,202,000 acres) is bounded by the borders of a vegetation map produced in 2004 for the Mojave Desert Ecosystem Program (MDEP). VegCAMP mapped a portion of the area mapped previously for the MDEP (approx. 776,000 acres) using the finer-scale rules and classification that AIS used for the larger portion (see Figure 1). The boundary of VegCAMP's study area was chosen to eliminate an arbitrary hole in AIS's work area based on the MDEP boundary around Ord Mountain, absorb an additional core area for the Mojave Ground Squirrel, include more critical habitat area for the Desert Tortoise, and ensure access to 1-ft. ancillary imagery via ImageConnect®.

The vegetation classification follows Federal Geographic Data Committee (FGDC) and National Vegetation Classification Standards (NVCS). The classification is based on previous survey and classification work including the classification done for MDEP. An additional 98 Rapid Assessment vegetation field surveys were collected in 2011, and some additional mapping classes are based on those unclassified surveys. These classes are considered provisional until better understood (for example, the *Ericameria cooperi* provisional alliance). See Appendix A for the field form and protocol for the Rapid Assessment surveys.

The map was produced using heads up digitizing on a base of true-color and color infrared 2010 1-meter National Agricultural Imagery Program (NAIP) imagery. Supplemental imagery included Bing mapping services and true-color 1-foot aerial imagery available through GlobeXplorer ImageConnect®. The minimum mapping unit (MMU) is 10 acres; exceptions are made for wetlands and certain wash types (which were mapped to a 1 acre MMU) and areas characterized as human land use polygons (which were mapped to a 2.5 acre MMU).

This report and accompanying data are to be released at the end of June 2012. After this time, a review of AIS polygons and fieldwork will be performed to analyze the percentage of mapped polygons that have had some sort of field assessment related to this project. Additionally, some of the polygons that were mapped to higher levels in the NVCS hierarchy will be re-assessed and assigned to a lower level.

Purpose

The purpose of the vegetation map is to assist with the Desert Renewable Energy Conservation Plan (DRECP). This map will provide planners with detailed information to help identify high quality habitat and rare communities. Although the primary purpose of the map is to document vegetation communities, it also provides structural data such as herbaceous, shrub and tree cover, and information about the level of disturbance within the vegetation stand. These are important habitat factors for species covered in the DRECP, including the Mojave Ground Squirrel and Desert Tortoise. Please read the attribute descriptions for caveats.

Supplemental Information

Classification

The map classification is based largely on work done in the area for previous and ongoing projects: Vegetation Mapping of Anza-Borrego Desert State Park and Environs (1998), the Mojave Desert Ecosystem Program's Vegetation Database (2004), Vegetation of Joshua Tree National Park (unpublished draft), and Vegetation Classification and Mapping at Lake Mead National Recreation Area, Mojave National Preserve and Death Valley National Park (in progress). These were compiled and placed within the NVCS hierarchy as it stood in March of 2011. Please see Appendices B and C for the classification and key, respectively.

Delineation Rules

Lines are drawn both to distinguish between types (MapUnits) and to indicate vegetation cover breaks within a type. The following minimum mapping units are observed while making these delineations: 1 acre for wetlands and certain wash types, 2.5 acres for areas characterized as human land use, and 10 acres for all other polygons.

An additional guideline establishes the approximate minimum width of a polygon: washes and riparian stands must be 90 ft. wide and upland vegetation must be 330 ft. wide. The appropriate MMUs must still be observed. This guideline isn't meant to exclude polygons where a small section falls below the minimum width, as long as the greater portion of the polygon meets the stated criteria.

Land use, including towns, mining, agriculture, and individual settlements, was handled using a special set of rules. These were designed to separate larger urban tracts as a whole, and to maintain the ability to have a vegetation type on a lightly used or re-vegetating area. See Appendix D for Land use rules.

Reconnaissance

Between February 2011 and October 2011, AIS staff and CDFG staff each conducted eight field trips throughout the mapping area with 1-2 crews per trip to perform reconnaissance of vegetation types. This reconnaissance allowed better matching of signatures seen on the imagery with the vegetation. Todd Keeler-Wolf of CDFG often accompanied AIS on their trips. During reconnaissance, crews traversed the study areas in vehicles, stopping to assess the vegetation types at various points. GPS points were taken and observations were recorded for vegetation type and cover at that point. Observations were also made for vegetation seen at a distance, with the point of observation determined using a compass and laser rangefinder. Points were frequently taken to mark the transition from one vegetation type to another, to help the photo-interpreter determine the location of the edges of stands. Sometimes one observation contained information about two or more stands, and other times the same stand was assessed in multiple places. Excluding some duplicate observations, approximately 4600 reconnaissance observations were made. See Appendix E for a copy of the form used by CDFG staff during reconnaissance surveys.

Accuracy Assessment

From October 2011 through June 2012, 2387 accuracy assessments (AAs) were made throughout the mapping area. Assessments were done by navigating to polygons allocated from a module of completed vegetation polygons. The allocations were designed to ensure that most, if not all, of the mapped vegetation types were assessed, and the allocated polygons were chosen based on their accessibility by road. The assessments were stand-based, that is, both the type and the extent of the polygons were evaluated where possible. Additionally, all of the cover values and all disturbance attributes except roadedness were assessed, where possible. When a polygon could be divided due to the presence of multiple types, an assessment was done for each type. As with reconnaissance, some of the stands were assessed from a distance using a compass, laser rangefinder, binoculars, and occasionally a spotting scope. ESRI's ArcPad software, loaded on Trimble's Juno SB devices, was used to collect GPS locations, and ArcPad's offset function was used to assist with distance surveys. See Appendix F for a copy of the form used in accuracy assessments.

In the office, data from the field AA forms were entered into an Access database and the vegetation type recorded by the photo-interpreter was scored using the field surveys and accompanying ground photos. Cover and disturbance attributes were not scored, but were provided as feedback to the photo-interpreters.

Each assessment could receive a maximum of 5 points, with points given depending on how closely the photo-interpreter (PI) matched the type reported from the AA. Below is a table showing how these points were assigned.

Code	Reason For Score	Score
A	PI completely correct.	5
B	The PI chose the correct Group OR the next level up in the hierarchy.	4
C	Threshold/transition between PI call and Final call. This was used when cover values of the dominant or indicator species were close to the values that would key to the PI's type (e.g., an AA call of <i>Yucca brevifolia</i> Alliance for a stand with 1% evenly distributed <i>Yucca brevifolia</i> over <i>Larrea tridentata</i> - <i>Ambrosia dumosa</i> would get this score if the PI call was <i>Larrea tridentata</i> - <i>Ambrosia dumosa</i> Alliance with <1% <i>Yucca brevifolia</i>).	4
D	Correct Macro Group OR next level up in hierarchy.	3
E	Based on close ecological similarity. Ecological similarity addresses assessed and mapped calls that contained vegetation with overlapping diagnostic species but were not technically closely related in the NVCS hierarchy. This was common in stands that contain a mix of species of late and early seral vegetation types and also common in zones of overlap between ecoregions.	3
F	Correct Division.	2
G	Some floristic/hydrologic similarity. This addresses cases in which the mapped and the assessed vegetation type had different diagnostic species, but bore some similarity in ecological traits based on predicted and actual setting such as hydrologic regime, overall climate, or successional state.	2
H	Correct only at Lifeform.	1
I	No similarity above Formation and incorrect life form.	0
J	Survey removed because significant change in polygon (e.g., the stand was burned, developed, or cleared since the date of the base imagery).	no score
K	Survey removed because inadequate portion (<10%) of the polygon was viewed by the AA field crew.	no score
L	Survey removed because field/PI data is incomplete, inadequate or confusing (e.g., cover values were not provided for key species in the stand).	no score
M	Supplementary record, not scored (for multiple point assessments where the AA call was the same at multiple points).	no score

The map received an overall score of 82.6%. The table below summarizes the average score per type sampled. A contingency table for the area will be available in the final version of this report.

Vegetation Type	Average Score	# of Samples
<i>Acacia greggii</i>	4.0	24
<i>Achnatherum hymenoides</i>	5.0	2
<i>Achnatherum speciosum</i>	5.0	1
<i>Adenostoma fasciculatum</i>	4.1	25
<i>Adenostoma fasciculatum</i> - <i>Salvia mellifera</i>	4.0	1
<i>Allenrolfea occidentalis</i>	5.0	5
<i>Ambrosia dumosa</i>	3.8	67
<i>Ambrosia salsola</i>	3.3	54
<i>Amsinckia (menziesii, tessellata)</i>	4.9	7
Anthropogenic areas of little or no vegetation	2.8	5
<i>Arctostaphylos glauca</i>	4.0	2
<i>Artemisia tridentata</i> (including <i>A. tridentata</i> spp. <i>parishii</i>)	4.3	25
<i>Atriplex canescens</i>	4.0	43
<i>Atriplex confertifolia</i>	3.7	41
<i>Atriplex hymenelytra</i>	3.5	22

Vegetation Type	Average Score	# of Samples
<i>Atriplex lentiformis</i>	1.7	3
<i>Atriplex polycarpa</i>	3.9	145
<i>Atriplex spinifera</i>	4.0	71
<i>Baccharis salicifolia</i>	4.1	7
Brassica nigra & other mustards	5.0	2
<i>Brickellia desertorum</i>	4.0	2
<i>Brickellia incana</i>	4.2	5
<i>Bromus rubens</i> - <i>Schismus</i> (<i>arabicus</i> , <i>barbatus</i>)	4.9	11
CA annual forb/grass vegetation Gp	4.3	21
CA perennial grassland Gp	1.0	1
Calif. Annual & Perennial Grassland (native component) MU	4.7	27
Calif. Annual & Perennial Grassland MG	3.6	7
<i>Ceanothus crassifolius</i>	5.0	2
<i>Cercocarpus montanus</i>	4.6	11
<i>Chilopsis linearis</i>	4.8	19
<i>Coleogyne ramosissima</i>	4.1	22
<i>Corethrogyne filaginifolia</i>	4.0	1
Desert Pavement Mapping Unit	4.5	27
<i>Distichlis spicata</i>	3.6	5
<i>Encelia farinosa</i>	4.3	19
<i>Encelia virginensis</i>	3.6	24
<i>Ephedra californica</i>	4.3	23
<i>Ephedra nevadensis</i>	3.7	35
<i>Ephedra viridis</i>	3.6	15
<i>Ericameria cooperi</i>	2.9	8
<i>Ericameria linearifolia</i>	4.0	4
<i>Ericameria nauseosa</i>	3.6	51
<i>Ericameria paniculata</i>	4.1	9
<i>Ericameria teretifolia</i>	3.6	25
<i>Eriodictyon</i> (<i>crassifolium</i> , <i>trichocalyx</i>)	3.9	9
<i>Eriogonum</i> (<i>elongatum</i> , <i>nudum</i>)	5.0	2
<i>Eriogonum fasciculatum</i>	3.7	73
<i>Eriogonum wrightii</i>	4.0	1
<i>Eschscholzia</i> (<i>californica</i>)	5.0	7
<i>Forestiera pubescens</i>	4.4	7
<i>Fremontodendron californicum</i>	3.4	5
<i>Geraea canescens</i> - <i>Chorizanthe rigida</i> MU	5.0	2
<i>Grayia spinosa</i>	3.6	60
<i>Gutierrezia sarothrae</i>	3.3	3
<i>Hyptis emoryi</i>	4.3	3
Intermontane deep or well-drained soil scrub Gp	3.2	5
<i>Isocoma acradenia</i>	3.0	1
<i>Juncus arcticus</i> (var. <i>balticus</i> , <i>mexicanus</i>)	3.0	2
<i>Juniperus californica</i>	4.8	48
<i>Krascheninnikovia lanata</i>	3.3	10
<i>Larrea tridentata</i>	4.3	141
<i>Larrea tridentata</i> - <i>Ambrosia dumosa</i>	4.2	172
<i>Larrea tridentata</i> - <i>Encelia farinosa</i>	4.4	12

Vegetation Type	Average Score	# of Samples
Lasthenia californica - Plantago erecta - Vulpia microstachys	5.0	2
Lepidospartum squamatum	4.0	42
Lycium cooperi	3.5	6
massive sparsely vegetated rock outcrop	4.2	13
Medit. CA naturalized annual & perennial grassland Gp	4.3	42
Menodora spinescens	4.0	1
mud hills sparsely vegetated ephemeral herbs (Annual Eriogonum, Plantago, etc.)	4.3	18
North American warm desert dunes and sand flats Gp	5.0	2
Panicum urvilleanum	2.0	1
Peucephyllum schottii	4.0	1
Pinus monophylla	5.0	10
Pinus sabiniana	5.0	1
Platanus racemosa	4.5	8
Pleuraphis rigida Alliance	4.8	5
Populus fremontii	4.5	22
Prosopis glandulosa	4.9	15
Prosopis pubescens	0.0	1
Prunus fasciculata	4.0	27
Prunus ilicifolia	4.0	1
Pseudotsuga macrocarpa	5.0	1
Psoralea spinosa	4.8	11
Purshia tridentata	4.5	11
Quercus berberidifolia	4.0	3
Quercus berberidifolia-Adenostoma fasciculatum	4.0	1
Quercus chrysolepis	4.5	4
Quercus douglasii	5.0	3
Quercus john-tuckeri	4.5	26
Quercus wislizeni	3.5	4
Salazaria mexicana	3.6	16
Salix exigua	3.9	9
Salix laevigata	3.8	6
Salix lasiolepis	4.0	7
Sambucus nigra	5.0	2
Sarcobatus vermiculatus	5.0	1
Schoenoplectus spp. Mapping Unit	4.3	3
Senna armata (not currently defined)	4.0	2
sparsely vegetated playa (ephemeral annuals)	4.8	36
Suaeda moquinii	4.2	36
Tamarix spp.	5.0	18
Typha (angustifolia, domingensis, latifolia)	5.0	8
unvegetated wash and river bottom	5.0	28
Viguiera parishii	5.0	5
Yucca brevifolia	4.7	76
Yucca schidigera	4.3	48

Portion of Map Visited

The final report will include an evaluation of the percent of mapped polygons that had some type of field assessment (rapid assessment, reconnaissance observation, accuracy assessment).

Map Attributes

NVCSName/NVCSLevel: The standardized name/level of the vegetation description used in the National Vegetation Classification System (see <http://biology.usgs.gov/npsveg/nvcs.html>). Since the NVCS does not have categories for human land use or otherwise unvegetated land, those descriptions were drawn from the California Wildlife Habitat Relationship.

MapUnit: The code for a vegetation alliance or group or an unvegetated type like playa; or a code for a land use for polygons that are not natural vegetation (please see the classification list and key).

Conifer Cover (ConCov): Percent bird's-eye cover of conifers within a vegetation stand, broken into the classes in the table below. "Bird's-eye" total cover is what can be seen on an air photo, i.e., it does not include the cover of understory layers obscured by overstory layers.

Code	Range	Example
0	none	There are no conifers in this stand.
1	>0-1%	<i>Juniperus californica</i> , <i>Pinus monophylla</i> , <i>P. sabiniana</i> are widely scattered as emergents. Note: if <i>J. californica</i> is present with >1% <i>Yucca brevifolia</i> , this is a <i>Y. brevifolia</i> alliance.
2	>1-5%	Conifers are sparse and unevenly scattered to dispersed and evenly distributed. If no other tree species have greater cover and the type is 1000 = TEMPERATE FOREST SUBCLASS, then the conifer is usually not an indicator species (the nominal type). The exception is <i>Juniperus californica</i> over scrub or herb understory. For example, <i>Pinus sabiniana</i> at 2% with <i>Quercus douglasii</i> at 11% would be a <i>Quercus douglasii</i> type, but <i>J. californica</i> at 5% over shrubs at 14% would be a <i>J. californica</i> type.
3	>5-15%	This is a common cover class for <i>Juniperus californica</i> in the western Mojave Desert, and an occasional cover class for <i>Pinus monophylla</i> or <i>Pseudotsuga macrocarpa</i> in and near the Transverse Ranges.
4	>15-25%	This is an occasional cover class for <i>Juniperus californica</i> , <i>Pinus monophylla</i> , and <i>Pseudotsuga macrocarpa</i> .
5	>25-50%	This class is unusual, except perhaps in the most shaded/mesic setting of lower canyons in the Transverse Range.
6	>50-75%	This cover class is unlikely in the mapping area.
7	>75-100%	This cover class is highly unlikely in the mapping area.
9	NA	Conifer cover is not applicable when the MapUnit is 9210, 9220, 9300, 9310, 9800, 9801, 9803, 9804, 9805.

Hardwood Cover (HdwdCov): Percent bird's-eye cover of hardwoods within a vegetation stand, broken into the following classes:

Code	Range	Example
0	none	This class is typical for most desert upland away from the Transverse or Tehachapi ranges.
1	>0-1%	The stand may include isolated <i>Platanus racemosa</i> , <i>Populus fremontii</i> , <i>Salix gooddingii</i> , or <i>Salix laevigata</i> in riparian scrub.
2	>1-5%	A low cover of evenly distributed riparian trees is indicated. If <i>Populus fremontii</i> , <i>Platanus racemosa</i> , <i>Salix gooddingii</i> or other tree willows are evenly distributed over marginally higher cover of riparian shrubs, the stand may be labeled as the appropriate tree alliance type.

Code	Range	Example
3	>5-15%	This class is typical of open <i>Quercus douglasii</i> or <i>Quercus lobata</i> stands in the Transverse or Tehachapi ranges, or of riparian woodlands with <i>Populus fremontii</i> , <i>Salix gooddingii</i> , etc.
4	>15-25%	This class would be an uncommonly high density of upland oak types or riparian woodlands.
5	>25-50%	This class would be an unusually high cover of upland oak, or riparian woodlands dominated by hardwoods.
6	>50-75%	This would indicate a quite uncommon, locally dense riparian stand of trees.
7	>75-100%	Cover this high is not expected in the study area
9	NA	Hardwood cover is not applicable when the MapUnit is 9210, 9220, 9300, 9310, 9800, 9801, 9803, 9804, 9805.

Joshua tree cover (YUBRcover): Percent bird's-eye cover of *Yucca brevifolia* within a vegetation stand, broken into the following classes:

Code	Range	Example
0	none	There are no visible <i>Yucca brevifolia</i> in the stand, although widely scattered juveniles <3m tall may be included.
1	present, but <1% and unevenly distributed	This is common in desert shrublands of <i>Larrea tridentata</i> - <i>Ambrosia dumosa</i> , <i>Atriplex polycarpa</i> , <i>Atriplex canescens</i> , <i>Coleogyne ramosissima</i> , etc. <i>Yucca brevifolia</i> tree signature may be visible, but individuals are not evenly distributed and are widely dispersed.
2	present, between 1 and 5% and evenly distributed	This class commonly denotes an open, tree-size <i>Yucca brevifolia</i> woodland, usually evenly distributed with higher shrub cover.
3	> 5%, generally dense clonal stands or occasionally in dense woodlands of spreading tree morphs (rare)	This density is usually only found at higher or wetter sites within the study area; it can occasionally involve non-clonal stands at highest density, as near the southwest side of Ft Irwin.
9	NA	Joshua tree cover is not applicable when MapUnit is 9210, 9220, 9300, 9310, 9800, 9801, 9803, 9804, 9805.

Total Tree Cover (TreeCov): Percent bird's-eye cover of trees (including Joshua Trees) within a vegetation stand, broken into the following classes:

Code	Range	Example
0	none	Most desert shrubland lacking emergent <i>Yucca brevifolia</i> falls into this class.
1	>0-1%	<i>Larrea tridentata</i> - <i>Ambrosia dumosa</i> , <i>Atriplex polycarpa</i> , <i>Atriplex spinifera</i> , <i>Coleogyne ramosissima</i> , etc. with isolated emergent tree-size <i>Juniperus californica</i> , <i>Pinus monophylla</i> , <i>Yucca brevifolia</i> are examples of this class, as well as wetlands with isolated tree <i>Salix</i> or <i>Populus</i> .
2	>1-5%	Most <i>Yucca brevifolia</i> or open <i>Juniperus californica</i> woodlands over a well-developed shrub cover are included. Desert washes with scattered <i>Chilopsis linearis</i> , <i>Psoralea spinosa</i> are other examples.
3	>5-15%	The highest density <i>Yucca brevifolia</i> (usually the short clonal type in west Mojave), well developed <i>Juniperus californica</i> woodland, and open <i>Quercus lobata</i> , <i>Q. douglasii</i> woodland in the Transverse and Tehachapi ranges fall in this class.
4	>15-25%	<i>Quercus lobata</i> , <i>Q. douglasii</i> , <i>Q. chrysolepis</i> woodlands in concavities in the Transverse and Tehachapi ranges are examples.

Code	Range	Example
5	>25-50%	This may be high-cover <i>Pseudotsuga macrocarpa</i> (with <i>Quercus chrysolepis</i>) stands in concavities in the Transverse Range on the edge of the ecoregion, or locally dense stands of <i>Populus fremontii</i> or other riparian trees.
6	>50-75%	This class is unlikely in the study area.
7	>75-100%	This class is highly unlikely in study area
9	NA	Tree cover is not applicable when the MapUnit is 9210, 9220, 9300, 9310, 9800, 9801, 9803, 9804, 9805

Shrub Cover (ShrubCov): Percent bird's eye cover of shrubs within a vegetation stand, broken into the following classes:

Code	Range	Example
0	none	There is no perennial shrub signature, ie: extensive cliffs and outcrops, extensive dunes or sand sheets, current agriculture, urban areas, etc.
1	>0-1%	These types are not defined by shrubs; they do not have an even distribution of shrubs (wetland or upland herbaceous vegetation).
2	>1-5%	Shrubs are widely distributed on harsh substrates, e.g., <i>Larrea tridentata-Ambrosia dumosa</i> or <i>Encelia farinosa</i> on steep rocky slopes, or old inactive alluvial surfaces.
3	>5-15%	Modal <i>Larrea tridentata-Ambrosia dumosa</i> , <i>Atriplex</i> spp. upland vegetation, active vegetated washes (e.g., <i>Ephedra californica</i> , <i>Ericameria paniculata</i> , <i>Ambrosia salsola</i>) fall in this range.
4	>15-25%	High cover <i>Larrea tridentata-Ambrosia dumosa</i> (as in west Mojave near Inyokern with high density <i>Ambrosia dumosa</i>), well developed <i>Coleogyne ramosissima</i> on moderate rocky slopes, well defined <i>Atriplex polycarpa</i> in washes and swales in the western portion of the study area are examples.
5	>25-50%	High cover chaparral of <i>Quercus john-tuckeri</i> , <i>Adenostoma fasciculatum</i> , <i>Eriogonum fasciculatum</i> , <i>Arctostaphylos glauca</i> , <i>Fremontodendron californicum</i> , and other chaparral or coastal sage scrub types may be found in the foothills of the Transverse, Tehachapi or Sierra Nevada ranges. Small stands of <i>Salix</i> or <i>Prosopis</i> associated with springs, stable reservoirs or other water bodies are in this range. This cover class is rare in the study area.
6	>50-75%	This is extremely rare in the mapping area.
7	>75-100%	Incidental, no stands in this range are known.
9	NA	Shrub cover is not applicable when the MapUnit is 9210, 9220, 9300, 9310, 9800, 9801, 9803, 9804, 9805.

Herbaceous Cover (HerbCov): Percent bird's-eye cover of herbaceous plants within a vegetation stand, broken into the following classes:

Code	Range	Example
1	0-2%	Steep slopes with low shrub cover (shrub cover class 2) and no herbaceous signature, ancient non-active alluvial fans.
2	>2-15%	Most standard <i>Larrea tridentata-Ambrosia dumosa</i> stands, wash stands of <i>Ambrosia salsola</i> , <i>Ephedra californica</i> , <i>Chilopsis linearis</i> without any surface moisture are expected to have herb covers in this class.
3	>15-40%	Most wetland woody and herbaceous polygons, and degraded <i>Larrea tridentata</i> or <i>Larrea tridentata-Ambrosia dumosa</i> have visible and evenly dispersed largely non-native herb cover.
4	>40%	Herb cover in this range is only found in dense wetlands, ie: local stands of <i>Typha</i> , <i>Juncus</i> , <i>Schoenoplectus</i> , etc.
9	NA	Herb cover is not applicable when the MapUnit is 9210, 9220, 9300, 9310, 9800, 9801, 9803, 9804, 9805.

Exotics: Level of impact by exotic invasive species, broken into the following classes

Code	Range	Example
0	no evidence of exotics visible	This is expected on coarse rocky slopes, with no tawny or reddish <i>Schismus</i> or <i>Bromus rubens</i> signature, desert pavement, very steep bouldery slopes, etc.
1	patches of exotics visible, but cover not significant (relative cover to total <33%)	General <i>Larrea tridentata</i> - <i>Ambrosia dumosa</i> without high roadedness or degraded understory usually fall in this class. <i>Schismus</i> or <i>Bromus</i> may be visible in relatively small discrete patches (<1/2 of the substrate signature), but will not present an extensive signature.
2	exotics (particularly herbaceous) significant and cover may exceed dominant vegetation strata (relative cover <66%)	A "haze" of <i>Schismus</i> (tawny) or <i>Bromus rubens</i> (reddish brown) is uniform in the understory of shrubby or treed overstory, as in degraded <i>Larrea tridentata</i> (lacking much <i>Ambrosia dumosa</i>), or ochre "haze" of <i>Brassica</i> is found in the sandy soil of <i>Larrea tridentata</i> - <i>Ambrosia dumosa</i> , or a brown to dark gray signature of <i>Salsola</i> is associated with a native woody overstory (e.g., <i>Suaeda</i> , <i>Atriplex</i> spp.) in saline or alkaline soils.
3	stand characterized by exotics (vegetation type is "exotic") (relative cover >66%)	The overstory is dominated or characterized by non-native exotics (either woody or herbaceous) without an adequate cover of characteristic native species to define it as a native vegetation type.
9	NA	Exotics is not applicable when the MapUnit is 9300, 9310, 9320, 9800, 9801, 9803, 9804, 9805.

Roadedness: Level of impact by paved and unpaved roads, OHV trails, railroads, etc. Impact is defined by proportion of any polygon of vegetation that is roadless. This has the advantage of helping to identify roadless areas—but the disadvantage of being scale independent. A polygon with a road more or less bisecting it will have a rating of 2, regardless of size; a very large polygon with this rating might still contain a big roadless area. This is broken into the following classes:

Code	Range	Example
0	none visible	
1	low: at least 2/3 (67% to 100%) of the vegetation polygon area is un-roaded by any type of road or OHV track	<p>ROADEDNESS 1:</p>
2	moderate: between 1/3 and 2/3 (33% to 66%) of the vegetation polygon is intersected by any type of road or OHV track	<p>ROADEDNESS 2:</p>

3	high: <33% of the vegetation polygon lacks roads of any kind	
9	NA	Roadedness is not applicable when the MapUnit is 9210, 9220, 9300, 9310, 9800, 9801, 9803, 9804, 9805, 9320 (non OHV).

Development: Level of impact by structures (buildings, tanks, paved parking lots, trailers, utility and mining structures) This is for areas where low mmu settlement cannot be pulled out, or the development does not meet the criteria of a settlement. NOTE: This attribute also includes anthropogenic debris (junked vehicles, major trash dumping, collapsed structures, etc.), the removal of which could result in a vegetation stand that would otherwise be in very good to pristine ecological condition.

Code	Range	Example
0	none visible	There are no noticeable junk piles, isolated homes or structures within polygon.
1	low: less than 2% of polygon affected	Junk piles, structures, cement pads, etc. are widely spaced at very low density.
2	moderate: between 3%- 5% of the polygon affected	Multiple examples of dispersed junk, houses, other structures are visible throughout the polygon. There may be a dense concentration of development within a single or few parts of a vegetation polygon, but no developed concentration larger than 2.5 acres (the minimum threshold for pulling out a single "settlement" or 9300 polygon).
3	high: > 5% of polygon affected	Multiple examples are evenly distributed in a vegetated polygon, but none are large enough to pull out as 9300.
9	NA	Development is not applicable when the MapUnit is 9210, 9220, 9801.

AnthropogenicAlteration: Level of impact on vegetation by anthropogenic clearing through tillage, scraping, grazing, etc. Fire effects are not considered in this category. This captures past disturbances in the landscape which are still visible through their impact on vegetation when that impact is not significant enough to be a "line-former." This attribute can also be used when small-scale effects are present that may cause a break in cover class, but this break affects an area less than the MMU rule for the vegetation type in question.

Code	Range	Example
0	none visible	No "ghost" lines of tilling, differential effects of enclosure/exclosure fencing, effects of grazing/browsing, etc. are visible.
1	less than 33% of polygon is affected and/or impact is seen but does not affect vegetation density (as broken down here) or type	Less than 1/3 of a vegetation polygon has visible evidence of clearing, prior agricultural activity or other effects.
2	between 33%-66% of the polygon is affected	A vegetation polygon has >1/3 but <2/3 visible effects of clearing, prior agricultural or other effects.

Code	Range	Example
3	>66% of polygon affected	A vegetation polygon has >2/3 visible effects of clearing, prior agricultural or other effects.
9	NA	Anthropogenic alteration is not applicable when the MapUnit is 9801.

HydroModifier (yes/no): Is the vegetation stand modified by a hydrologic impediment? This is used to tag stands of desert vegetation that have their extent directly impacted by restricted sheet flow or active channel flow crossing under a road, railroad, levee, etc. Examples include polygons where: 1) washes have expanded on the upslope side or contracted on the downslope side of the impediment (typically a berm or levee), or 2) railroad or highway berms have eliminated natural sheet flow downslope across alluvial fans and bajadas, or other slopes. The effect must create a line-forming break such as in vegetation type, shrub cover, tree cover, or herbaceous cover. The modifier is only attributed to the polygon down-slope of the impediment. Drainage ditches conveying flow off the side of a road (though often visible on imagery) are not considered unless they make a line-forming break in the vegetation.

Notes: Text field for additional information.

LandUse: If a polygon is either designated as a water or land use type (MapUnit >9000) or the polygon represents a stand with sub-mmua land use area within it, then a land use code is assigned to the polygon. See Appendix D for further information.

Land use code	Description
0	Not Assigned, Not Assessed
1000	Urban
1436	Water Transfer
2100	Non-woody row and field crops
2200	Orchards and Vineyards
9800	Undifferentiated water
9810	Water Impoundment Feature

MethodID: For areas that were mapped by CDFG (see Mappers field below), this field identifies what type of field data (if any were available) supported the vegetation type decision for that polygon. Some of the stands were assessed from a distance using a compass, laser rangefinder, binoculars, and occasionally a spotting scope. For reconnaissance, multiple stands were often assessed from a single location. [MethodID is currently filled in only for areas delineated by CDFG.]

Method Code	Description
1	Rapid Assessment (current project)
2	Releve
3	Field Verification or Accuracy Assessment
4	Photo Interpretation
5	Adjacent stand information or photo
6	Reconnaissance (current project)
7	Other information
8	Older plot data
9	Older recon data

Mappers: Indicates whether a polygon was delineated and attributed by Aerial Information Systems (AIS), or the California Department of Fish and Game's Vegetation Classification and Mapping Program (CDFG).

NRFVconfidence: No regional field verification. For areas that were mapped by CDFG (see Mappers field above), this field identifies photo-interpreted polygons in large, contiguous areas that were inaccessible and in which the confidence in correctly identifying the vegetation type was therefore low. These are marked with "low" in this field.

Conservation: The status of a plant community based on rarity and habitat value. This field is based on the interaction between rarity ranking and the site quality attributes assigned to each polygon. State and globally assigned rarity ranks have already been assigned (see vegCAMP website: http://dfg.ca.gov/biogeodata/vegcamp/natural_comm_background.asp and http://dfg.ca.gov/biogeodata/vegcamp/natural_comm_list.asp) and locally estimated rarity within the Mojave and Sonoran desert ecoregions of California (Miles and Goudey 1997) was assigned by the VegCamp team based on relative representation of the vegetation unit within the mapping area. If a vegetation type was considered rare (S3 or lower) in the state, it automatically also received a locally rare code within the mapping area. However, some vegetation types which are common throughout the state could be considered locally rare if they were represented by a relatively small number of mapped occurrences compared to their overall state distribution. These codified rarity levels were then subjected to analysis based on attributes that pertain to the quality of the individual polygons, such as Development, Roadedness, Exotics, and Anthropogenic Alteration. A ranked prioritization system is used to give ratings as defined below.

Conservation	Description
Rare Vegetation Community S1 S2	Communities with a state rarity rank of S1 or S2 are assigned to the highest conservation level.
Locally Rare or Important Vegetation Community High Quality	Communities that have a state rarity rank of S3 or that are identified as locally rare or important, and are relatively undisturbed are assigned to this level. Four disturbance attributes were considered to define "relatively undisturbed." The values for Exotics, Development and Anthropogenic Alterations must all be less than 2. The value for Roadedness must be less than 2 for small polygons (<50 acres) or less than 3 for large polygons (>=50 acres).
Locally Rare or Important Vegetation. Community	Communities that have a state rarity rank of S3 or that are identified as locally rare or important, but do not have the lowest disturbance level are placed in this category.

CalVegName/CalVegCode: A crosswalk to the CalVeg vegetation system. Note that there may be a one-to-many relationship between CalVeg and NVCS. See <http://www.fs.usda.gov/detail/r5/landmanagement/resourcemanagement/?cid=stelprdb5347192>.

CWHRtype/CWHRcode: A crosswalk to the California Wildlife Habitat Relationships system. Note that there is usually a one-to-many relationship between CWHR and NVCS. See <http://www.dfg.ca.gov/biogeodata/cwhr/>.

GlobalRank/StateRank: The global/state rarity rank of the plant community mapped (only for alliances and associations).

Global Rank	State Rank	Description
G1	S1	There are fewer than 6 viable occurrences and/or less than 2000 acres of the community worldwide/statewide.

Global Rank	State Rank	Description
G2	S2	There are 6-20 viable occurrences and/or 2000-10,000 acres of the community worldwide/statewide.
G3	S3	There are 21-100 viable occurrences and/or 10,000-50,000 acres of the community worldwide/statewide.
G4	S4	There more than 100 viable occurrences and/or greater than 50,000 acres of the community worldwide/statewide.
G5	S5	The community is demonstrably secure due to secure worldwide and statewide abundance.

CACode: California Natural Community Codes - unique code assigned to alliances and associations.

NVSMG: The standardized name for the macrogroup within the National Vegetation Classification System. See <http://biology.usgs.gov/npsveg/nvcs.html>.

EcologicalSystems: A classification system used to create the Gap map for California. They are “groups of plant community types that tend to co-occur within landscapes with similar ecological processes, substrates and/or environmental gradients” (Comer et al. 2003). See <http://www.natureserve.org/library/usEcologicalsystems.pdf> and <http://gapanalysis.usgs.gov/gaplandcover/vision/>.

UID: Unique identifier for each polygon

Acres and Hectares: GIS-calculated area measurements of each mapped polygon.

Literature cited

Barbour, M.G., T. Keeler-Wolf, and A. Schoenherr, editors. 2007. *Terrestrial vegetation of California*, 3rd edition. University of California Press, Berkeley, CA.

Beatley, J.C..1976. *Vascular plants of the Nevada Test Site and central-southern Nevada: ecological and geographical distributions*. National Technical Information Service. U.S. Department of Commerce, Springfield, VA.

Comer, P., D. Faber-Langendoen, R. Evans, S. Gawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schulz, K. Snow, and J. Teague. 2003. *Ecological Systems of the United States: A Working Classification of U.S. Terrestrial Systems*. NatureServe, Arlington, Virginia. Available from: <http://www.natureserve.org/library/usEcologicalsystems.pdf>

Evens, J.M., K. Sikes, and D. Stout. 2011. Vegetation Classification and Mapping at Lake Mead National Recreation Area, Mojave National Preserve and Death Valley National Park, Annual Report. Natural Resource Data Series NPS/MOJN/NRDS—2011/XXX. National Park Service, Fort Collins, Colorado.

Faber-Langendoen, D., J. Drake, S. Gawler, M. Hall, G. Kittel, S. Menard, C. Nordman, M. Pyne, M. Reid, L. Sneddon, K. Schulz, J. Teague, M. Russo, K. Snow, P. Comer. 2012. *Macrogroups and Groups for the U.S. National Vegetation Classification*. NatureServe, Arlington, VA. + Appendices.

Faber-Langendoen, D., D.L Tart, and R.H.Crawford. 2009. Contours of the revised U.S. National Vegetation Classification standard. *Bulletin of the Ecological Society of America* **90**:87-93.

FGDC. 2008. National Vegetation Classification Standard, Version 2 FGDC-STD-005-2008 (version 2). Vegetation Subcommittee, Federal Geographic Data Committee, FGDC Secretariat, U.S. Geological Survey. Reston, VA. 55 pp. + Appendices.

Jennings, M.D., D. Faber-Langendoen, O.L. Loucks, R.K. Peet, and D. Roberts. 2009. Standards for associations and alliances of the U.S. National Vegetation Classification. *Ecological Monographs* 79: 173-199.

Keeler-Wolf, T. and J. Evens. 2006. Vegetation Classification of the Santa Monica Mountains National Recreation Area and Environs in Ventura and Los Angeles Counties, California. Version 1—Association Level and Specific Alliances. A report submitted to the National Park Service. California Department of Fish and Game and The California Native Plant Society, Sacramento, CA.

Keeler-Wolf, T., K. Lewis, and C. Roye. 1998. Vegetation Mapping of Anza-Borrego Desert State Park and Environs. California Department of Fish and Game, Sacramento, CA Available from: <http://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=18246>

Klein, A. and J. Evens, 2005.. Vegetation alliances of western Riverside County, California. Unpublished report, revised 2006, prepared for California Department of Fish and Game Habitat Conservation Division. California Native Plant Society. Sacramento, CA.

Lichvar, R., S. Sprecher, D. Charlton, G. Gustina, M. Ericsson, and J. Campbell. 2004. Ecosystem Classification and Relationships For Pleistocene Lake Thompson Bed, Mojave Desert, California. US Army Corps of Engineers®, Engineer Research and Development Center, ERDC/CRREL TR-04-21. Prepared for EDWARDS AIR FORCE BASE, CALIFORNIA

Miles, S.R., and C.B. Goudey, 1997. Ecological subregions of California. Technical Report R5-EM-TP-005, USDA Forest Service, Pacific Southwest Research Station, San Francisco, CA.

Peet, R.K. 2008. A decade of effort by the ESA Vegetation Panel leads to a new federal standard.. *Bulletin of the Ecological Society of America* 89:210-211.

Thomas, K., T. Keeler-Wolf, J. Franklin, and P. Stine. 2004. *Desert Ecosystem Program: Central Mojave Vegetation Database*. Final Report , U.S. Geological Survey, Western Ecological Research Center and Southwest Biological Science Center, Sacramento, CA. Colorado Plateau Field Station, Flagstaff, AZ. Available from: <http://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=13890>

Appendix A

Rapid Assessment Field Form & Protocol

CNPS and CDFG Combined Vegetation Rapid Assessment and Relevé Field Form

Relevé or Rapid Assessment (circle one)

(Desert Version Revised March 7, 2011)

For Office Use:	Final database #:	Final vegetation type name:	Alliance Association
-----------------	-------------------	-----------------------------	----------------------

I. LOCATIONAL/ENVIRONMENTAL DESCRIPTION

Polygon/Stand #:	Air photo:	Date:	Name(s) of surveyors (circle recorder):
------------------	------------	-------	---

GPS wypt #: _____ GPS name: _____ Datum: _____ or NAD83. Bearing, left axis at SW pt _____ (degrees) of Long / Short side
 UTME _____ UTMN _____ Zone: 10 / 11 (circle one) Error: ± _____ ft / m / pdop

GPS within stand? Yes / No If No, cite from waypoint to stand, distance _____ (meters) & bearing _____ (degrees)

Elevation: _____ ft / m Camera Name/Photograph #'s: _____

Stand Size (acres): <1, 1-5, >5 | Plot Size (m²): 10 / 100 / 400 / 1000 | Plot Shape _____ x _____ ft / m or Circle Radius _____ ft / m
 Exposure, Actual °: _____ NE NW SE SW Flat Variable All | Steepness, Actual °: _____ 0° 1-5° 5-25° > 25

Topography: Macro: top upper mid lower bottom | Micro: convex flat concave undulating
 Geology code: _____ Soil Texture code: _____ | Upland or Wetland/Riparian (circle one)

% Surface cover: _____ (Incl. outcrops) (>60cm diam) (25-60cm) (7.5-25cm) (2mm-7.5cm) (Incl sand, mud)
 H20: _____ BA Stems: _____ Litter: _____ Bedrock: _____ Boulder: _____ Stone: _____ Cobble: _____ Gravel: _____ Fines: _____ =100%

% Current year bioturbation _____ Past bioturbation present? Yes / No | % Hoof punch _____

Fire evidence: Yes / No (circle one) If yes, describe in Site history section, including date of fire, if known.

Site history, stand age, comments:

Disturbance code / Intensity (L,M,H): _____ / _____ / _____ / _____ / _____ / _____ / _____ "Other" _____ / _____

II. HABITAT AND VEGETATION DESCRIPTION

Tree DBH : T1 (<1" dbh), T2 (1-6" dbh), T3 (6-11" dbh), T4 (11-24" dbh), T5 (>24" dbh), T6 multi-layered (T3 or T4 layer under T5, >60% cover)

Shrub: S1 seedling (<3 yr. old), S2 young (<1% dead), S3 mature (1-25% dead), S4 decadent (>25% dead)

Herbaceous: H1 (<12" plant ht.), H2 (>12" ht.) Desert Riparian Tree/Shrub: 1 (<2ft. stem ht.), 2 (2-10ft. ht.), 3 (10-20ft. ht.), 4 (>20ft. ht.)

Desert Palm/Joshua Tree: 1 (<1.5" base diameter), 2 (1.5-6" diam.), 3 (>6" diam.) % NonVasc cover: _____ Total % Veg cover: _____

% Cover - Conifer tree / Hardwood tree: _____ / _____ Regenerating Tree: _____ Shrub: _____ Herbaceous: _____

Height Class - Conifer tree / Hardwood tree: _____ / _____ Regenerating Tree: _____ Shrub: _____ Herbaceous: _____

Height classes: 01=<1/2m 02=1/2-1m 03=1-2m 04=2-5m 05=5-10m 06=10-15m 07=15-20m 08=20-35m 09=35-50m 10=>50m

Species, Stratum, and % cover. Stratum categories: T=Tree, S = Shrub, H= Herb, E = SEedling, A = SApling, N= Non-vascular.

% cover intervals for reference: <1%, 1-5%, >5-15%, >15-25%, >25-50%, >50-75%, 75%.

Strata	Species	% cover	C	Strata	Species	% cover	C

Unusual species: _____

III. INTERPRETATION OF STAND

Field-assessed vegetation alliance name: _____

Field-assessed association name (optional): _____

Adjacent alliances/direction: _____ / _____, _____ / _____

Confidence in alliance identification: L M H Explain: _____

Phenology (E,P,L): Herb Shrub Tree Other identification or mapping information: _____

Is poly >1 type: Yes / No If yes, explain: _____

**CALIFORNIA NATIVE PLANT SOCIETY / DEPARTMENT OF FISH AND GAME PROTOCOL
FOR COMBINED VEGETATION RAPID ASSESSMENT
AND RELEVÉ SAMPLING FIELD FORM
(Modified for DRECP)**

Introduction

This protocol describes the methodology for both the relevé and rapid assessment vegetation sampling techniques as recorded in the combined relevé and rapid assessment field survey form dated March 22, 2010. The same environmental data are collected for both techniques. However, the relevé sample is plot-based, with each species in the plot and its cover being recorded. The rapid assessment sample is based not on a plot but on the entire stand, with 12-20 of the dominant or characteristic species and their cover values recorded. For more background on the relevé and rapid assessment sampling methods, see the relevé and rapid assessment protocols at www.cnps.org.

Selecting stands to sample:

To start either the relevé or rapid assessment method, a stand of vegetation needs to be defined. A stand is the basic physical unit of vegetation in a landscape. It has no set size. Some vegetation stands are very small, such as alpine meadow or tundra types, and some may be several square kilometers in size, such as desert or forest types. A stand is defined by two main unifying characteristics:

- 1) It has compositional integrity. Throughout the site, the combination of species is similar. The stand is differentiated from adjacent stands by a discernable boundary that may be abrupt or indistinct.
- 2) It has structural integrity. It has a similar history or environmental setting that affords relatively similar horizontal and vertical spacing of plant species. For example, a hillside forest originally dominated by the same species that burned on the upper part of the slopes, but not the lower, would be divided into two stands. Likewise, sparse woodland occupying a slope with very shallow rocky soils would be considered a different stand from an adjacent slope with deeper, moister soil and a denser woodland or forest of the same species.

The structural and compositional features of a stand are often combined into a term called homogeneity. For an area of vegetated ground to meet the requirements of a stand, it must be homogeneous (uniform in structure and composition throughout).

Stands to be sampled may be selected by evaluation prior to a site visit (*e.g.*, delineated from aerial photos or satellite images), or they may be selected on site during reconnaissance (to determine extent and boundaries, location of other similar stands, etc.).

Depending on the project goals, you may want to select just one or a few representative stands of each homogeneous vegetation type for sampling (*e.g.*, for developing a classification for a vegetation mapping project), or you may want to sample all of them (*e.g.*, to define a rare vegetation type and/or compare site quality between the few remaining stands).

For the rapid assessment method, you will collect data based on the entire stand.

Selecting a plot to sample within in a stand (for relevés only):

Because many stands are large, it may be difficult to summarize the species composition, cover, and structure of an entire stand. We are also usually trying to capture the most information as efficiently as possible. Thus, we are typically forced to select a representative portion to sample.

When sampling a vegetation stand, the main point to remember is to select a sample that, in as many ways possible, is representative of that stand. This means that you are not randomly selecting a plot; on the contrary, you are actively using your own best judgment to find a representative example of the stand.

Selecting a plot requires that you see enough of the stand you are sampling to feel comfortable in choosing a representative plot location. Take a brief walk through the stand and look for variations in species composition and in stand structure. In many cases in hilly or mountainous terrain look for a vantage point from which you can get a representative view of the whole stand. Variations in vegetation that are repeated throughout the stand should be included in your plot. Once you assess the variation within the stand, attempt to find an area that captures the stand's common species composition and structural condition to sample.

Plot Size

All relevés of the same type of vegetation to be analyzed in a study need to be the same size. Plot shape and size are somewhat dependent on the type of vegetation under study. Therefore, general guidelines for plot sizes of tree-, shrub-, and herbaceous communities have been established. Sufficient work has been done in temperate vegetation to be confident the following conventions will capture species richness:

- Herbaceous communities: 100 sq. m plot
- Special herbaceous communities, such as vernal pools, fens: 10 sq m plot
- Shrublands and Riparian forest/woodlands: 400 sq. m plot
- Open desert and other shrublands with widely dispersed but regularly occurring woody species: 1000 sq. m plot
- Upland Forest and woodland communities: 1000 sq. m plot

Plot Shape

A relevé has no fixed shape, though plot shape should reflect the character of the stand. If the stand is about the same size as a relevé, the plot boundaries may be similar to that of the entire stand. If we are sampling streamside riparian or other linear communities, our plot dimensions should not go beyond the community's natural ecological boundaries. Thus, a relatively long, narrow plot capturing the vegetation within the stand, but not outside it would be appropriate. Species present along the edges of the plot that are clearly part of the adjacent stand should be excluded.

If we are sampling broad homogeneous stands, we would most likely choose a shape such as a circle (which has the advantage of the edges being equidistant to the center point) or a square (which can be quickly laid out using perpendicular tapes).

Definitions of fields in the protocol

Relevé or Rapid Assessment Circle the method that you are using.

LOCATIONAL/ENVIRONMENTAL DESCRIPTION

Polygon/Stand #: Number assigned either in the field or in the office prior to sampling. It is usually denoted with a four-letter abbreviation of the sampling location and then a four-number sequential number of that locale (e.g. CARR0001 for Carrizo sample #1). The maximum number of letters/numbers is eight.

Air photo #: The number given to the aerial photo in a vegetation-mapping project, for which photo interpreters have already done photo interpretation and delineations of polygons. If the sample site has not been photo-interpreted, leave blank.

Date: Date of the sampling.

Name(s) of surveyors: The full names of each person assisting should be provided for the first field form for the day. On successive forms, initials of each person assisting can be recorded. Please note: The person recording the data on the form should circle their name/initials.

GPS waypoint #: The waypoint number assigned by a Global Positioning System (GPS) unit when marking and storing a waypoint for the sample location. Stored points should be downloaded in the office to serve as a check on the written points and to enter into a GIS.

For relevé plots, take the waypoint in the southwest corner of the plot or in the center of a circular plot.

GPS name: The name/number assigned to each GPS unit. This can be the serial number if another number is not assigned.

Datum: (NAD 83) The standard GPS datum used is NAD 83. If you are using a different datum, note it here.

Bearing, left axis at SW pt (note in degrees) of Long or Short side: For square or rectangular plots: from the SW corner (= the GPS point location), looking towards the plot, record the bearing of the axis to your left. If the plot is a rectangle, indicate whether the left side of the plot is the long or short side of the rectangle by circling “long” or “short” side (no need to circle anything for circular or square plots). If there are no stand constraints, you would choose a circular or square plot and straight-sided plots should be set up with boundaries running in the cardinal directions. If you choose a rectangular plot that is not constrained by the stand dimensions, the short side should run from east to west, while the long side should run from north to south.

UTM coordinates: Easting (UTME) and northing (UTMN) location coordinates using the Universal Transverse Mercator (UTM) grid. Record in writing the information from a GPS unit or a USGS topographic map.

UTM zone: Universal Transverse Mercator zone. Zone 10 is for California west of the 120th longitude, zone 11 is for California east of 120th longitude, which is the same as the straight portion of California’s eastern boundary.

Error: ± The accuracy of the GPS location, when taking the UTM field reading. Please record the error units by circling feet (ft), meters (m), or positional dilution of precision (pdop). If your GPS does not determine error, insert N/A in this field.

Is GPS within stand? Yes / No Circle “Yes” to denote that the GPS waypoint was taken directly within or at the edge of the stand being assessed for a rapid assessment, or circle “No” if the waypoint was taken at a distance from the stand (such as with a binocular view of the stand).

If No, cite from waypoint to stand, distance (note in meters) & bearing (note in degrees): An estimate of the number of meters and the compass bearing from the GPS waypoint to the stand.

Elevation: Recorded from the GPS unit or USGS topographic map. Please circle feet (ft) or meters (m).

Photograph #s: Write the name or initials of the camera owner, JPG/frame number, and direction of photos (note the roll number if using film). *Take four photos in the main cardinal directions (N, E, S, W) clockwise from the north, from the GPS location.* If additional photos are taken in other directions, please note this information on the form.

Stand Size: Estimate the size of the entire stand in which the sample is taken. As a measure, one acre is about 4000 square meters (approximately 64 x 64 m), or 208 feet by 208 feet. One acre is similar in size to a football field.

Plot Size: If this is a relevé, circle the size of the plot.

Plot Shape: Record the length and width of the plot and circle measurement units (i.e., ft or m). If it is a circular plot, enter radius (or just put a check mark in the space).

Exposure: (Enter actual ° and circle general category): With your back to the general uphill direction of the slope (i.e., by facing downhill of the slope), read degrees of the compass for the aspect or the direction you are standing, using degrees from north, adjusted for declination. Average the reading over the entire stand, even if you are sampling a relevé plot, since your plot is representative of the stand. If estimating the exposure, write “N/A” for the actual degrees, and circle the general category chosen. “Variable” may be selected if the same, homogenous stand of vegetation occurs across a varied range of slope exposures. Select “all” if stand is on top of a knoll that slopes in all directions or if the same, homogenous stand of vegetation occurs across all ranges of slope.

Steepness: (Enter actual ° and circle general category): Read degree slope from a compass or clinometer. If estimating, write “N/A” for the actual degrees, and circle the general category chosen. Make sure to average the reading across the entire stand even if you are sampling in a relevé plot.

Topography: First assess the broad (**Macro**) topographic feature or general position of the stand in the surrounding watershed, that is, the stand is at the top, upper (1/3 of slope), middle (1/3 of slope), lower (1/3 of slope), or bottom. **Circle all of the positions that apply for macrotopography.**

Then assess the local (**Micro**) topographic features or the lay of the area (e.g., surface is flat or concave). **Circle only one of the microtopographic descriptors.**

Geology: Geological parent material of site. If exact type is unknown, use a more general category (e.g., igneous, metamorphic, sedimentary). *See code list for types.*

Soil Texture: Record soil texture that is characteristic of the site (e.g., coarse loamy sand, sandy clay loam). *See soil texture key and code list for types.*

Upland or Wetland/Riparian (circle one): Indicate if the stand is in an upland or a wetland. There are only two options. Wetland and riparian are one category. Note that a site need not be officially delineated as a wetland to qualify as such in this context (e.g., seasonally wet meadow).

% Surface cover (abiotic substrates). It is helpful to imagine “mowing off” all of the live vegetation at the base of the plants and removing it – you will be estimating what is left covering the surface. **The total should sum to 100%.** Note that non-vascular cover (lichens, mosses, cryptobiotic crusts) is not estimated in this section.

% Water: Estimate the percent surface cover of running or standing water, ignoring the substrate below the water.

% BA Stems: Percent surface cover of the plant basal area, i.e., the basal area of stems at the ground surface. Note that for most vegetation types BA is 1-3% cover.

% Litter: Percent surface cover of litter, duff, or wood on the ground.

% Bedrock: Percent surface cover of bedrock.

% Boulders: Percent surface cover of rocks > 60 cm in diameter.

% Stone: Percent surface cover of rocks 25-60 cm in diameter.

% Cobble: Percent surface cover of rocks 7.5 to 25 cm in diameter.

% Gravel: Percent surface cover of rocks 2 mm to 7.5 cm in diameter.

% Fines: Percent surface cover of bare ground and fine sediment (e.g. dirt) < 2 mm in diameter.

% Current year bioturbation: Estimate the percent of the sample or stand exhibiting soil disturbance by fossorial organisms (any organism that lives underground). Do not include disturbance by ungulates. Note that this is a separate estimation from surface cover.

Past bioturbation present? Circle Yes if there is evidence of bioturbation from previous years.

% Hoof punch: Note the percent of the sample or stand surface that has been punched down by hooves (cattle or native grazers) in wet soil.

Fire Evidence: Circle Yes if there is visible evidence of fire, and note the type of evidence in the “Site history, stand age and comments section,” for example, “charred dead stems of *Quercus berberidifolia* extending 2 feet above resprouting shrubs.” If you are certain of the year of the fire, put this in the Site history section.

Site history, stand age, and comments: Briefly describe the stand age/seral stage, disturbance history, nature and extent of land use, and other site environmental and vegetation factors. Examples of disturbance history: fire, landslides, avalanching, drought, flood, animal burrowing, or pest outbreak. Also, try to estimate year or frequency of disturbance. Examples of land use: grazing, timber harvest, or mining. Examples of other site factors: exposed rocks, soil with fine-textured sediments, high litter/duff build-up, multi-storied vegetation structure, or other stand dynamics.

Disturbance code / Intensity (L,M,H): List codes for potential or existing impacts on the stability of the plant community. Characterize each impact each as **L** (=Light), **M** (=Moderate), or **H** (=Heavy). For invasive exotics, divide the total exotic cover (e.g. 25% *Bromus diandrus* + 8% *Bromus madritensis* + 5% *Centaurea melitensis* = 38% total exotics) by the total % cover of all the layers when added up (e.g. 15% tree + 5% low tree + 25% shrub + 40% herbs = 85% total) and multiply by 100 to get the % relative cover of exotics (e.g. 38% total exotics/85% total cover = 45% relative exotic cover). L = 0-33% relative cover of exotics; M = 34-66% relative cover, and H = > 66% relative cover. See code list for impacts.

II. HABITAT AND VEGETATION DESCRIPTION

California Wildlife-Habitat Relationships (CWHR)

For CWHR, identify the size/height class of the stand using the following tree, shrub, and/or herbaceous categories. These categories are based on functional life forms.

Tree DBH: Circle one of the tree size classes provided when the tree canopy closure exceeds 10 percent of the total cover, or if young tree density indicates imminent tree dominance. Size class is based on the average diameter at breast height (dbh) of each trunk (standard breast height is 4.5ft or 137cm). When marking the main size class, make sure to estimate the mean diameter of all trees over the entire stand, and weight the mean if there are some larger tree dbh's. The “**T6 multi-layered**” dbh size class contains a multi-layered tree canopy (with a size class T3 and/or T4 layer growing under a T5 layer and a distinct height separation between the classes) exceeding 60% total cover. Stands in the T6 class need also to contain at least 10% cover of size class 5 (>24” dbh) trees growing over a distinct layer with at least 10% combined cover of trees in size classes 3 or 4 (>11-24” dbh).

Shrub: Circle one of the shrub size classes provided when shrub canopy closure exceeds 10 percent (except in desert types) by recording which class is predominant in the survey. Shrub size class is based on the average amount of crown decadence (dead standing vegetation on live shrubs when looking across the crowns of the shrubs).

Herb: Circle one of the herb height classes when herbaceous cover exceeds 2 percent by recording the predominant class in the survey. Note: *This height class is based on the average plant height at maturity, not necessarily at the time of observation.*

Desert Palm/Joshua Tree: Circle one of the palm or Joshua tree size classes by averaging all the stem-base diameters (*i.e.* mean diameter of all stem-base sizes). Diameter is measured at the plant's base above the bulge near the ground.

Desert Riparian Tree/Shrub: Circle one of the size classes by measuring mean stem height (whether tree and/or shrub stand).

Overall Cover of Vegetation

Provide an estimate of cover for the following categories below (based on functional life forms). Record a specific number for the total aerial cover or "bird's-eye view" looking from above for each category, estimating cover for the living plants only. Litter/duff should not be included in these estimates. The porosity of the vegetation should be taken into consideration when estimating percent cover (how much of the sky can you see when you are standing under the canopy of a tree, or how much light passes through the canopy of the shrub layer?).

To come up with a specific number estimate for percent cover, first use generalized cover classes as reference aids such as the CWHR cover classes (<2%, 2-9%, 10-24%, 25-39%, 40-59%, 60-100%) or the modified Braun-Blanquet cover-abundance scale (<1%, 1-5%, >5-15%, >15-25%, >25-50%, >50-75%, >75%). While keeping these intervals in mind, you can then refine your estimate to a specific percentage for each category below.

% Total NonVasc cover: The total cover of all lichens, bryophytes (mosses, liverworts, hornworts), and cryptogammic crust on substrate surfaces including downed logs, rocks and soil, but not on standing or inclined trees or vertical rock surfaces.

% Total Vasc Veg cover: The total cover of all vascular vegetation taking into consideration the porosity, or the holes, in the vegetation. This is an estimate of the absolute vegetation cover, disregarding overlap of the various tree, shrub, and/or herbaceous layers and species.

% Cover by Layer

% Conifer Tree /Hardwood Tree: The total foliar cover (considering porosity) of all live tree species, disregarding overlap of individual trees. Estimate conifer and hardwood covers separately.

Please note: These cover values should not include the coverage of regenerating tree species (*i.e.*, tree seedlings and saplings).

% Regenerating Tree: The total foliar cover of seedlings and saplings, disregarding overlap of individual recruits. See seedling and sapling definitions below.

%Shrub: The total foliar cover (considering porosity) of all live shrub species disregarding overlap of individual shrubs.

%Herbaceous: The total cover (considering porosity) of all herbaceous species, disregarding overlap of individual herbs.

Height Class by Layer

Modal height for conifer tree /hardwood tree, shrub, and herbaceous categories: Provide an estimate of height for each category listed. Record an average height value per each category by

estimating the mean height for each group. Please use the following height intervals to record a height class: 01 =< 1/2m, 02=1/2-1m, 03 = 1-2 m, 04 = 2-5 m, 05 = 5-10 m, 06 = 10-15 m, 07 = 15-20 m, 08 = 20-35 m, 09 = 35-50 m, 10 => 50m.

Species List and Coverage

For rapid assessments, list the 10-20 species that are dominant or that are characteristically consistent throughout the stand. These species may or may not be abundant, but they should be constant representatives in the survey. When different layers of vegetation occur in the stand, make sure to list species from each stratum. As a general guide, make sure to list at least 1-2 of the most abundant species per stratum.

For relevés, list all species present in the plot, using the second species list page if necessary.

For both sample types, provide the stratum:

T = Tree. A woody perennial plant that has a single trunk.

S = Shrub. A perennial, woody plant, that is multi-branched and doesn't die back to the ground every year.

H = Herb. An annual or perennial that dies down to ground level every year.

E = SEedling. A tree species clearly of a very young age that is < 1" dbh.

A = SApling. 1" - <6" dbh and young in age, OR small trees that are < 1" diameter at breast height, are clearly of appreciable age, and kept short by repeated browsing, burning, or other disturbance.

N = Non-vascular. Includes moss, lichen, liverworts, hornworts, cryptogamic crust, and algae.

Be consistent and don't break up a single species into two separate strata. The only time it would be appropriate to do so is when one or more tree species are regenerating, in which case the Seedling and/or Sapling strata should be recorded for that species. These may be noted on the same line, e.g.:

Strata	Species	%Cover	C
T/E/A	Quercus douglasii	40/<1/<1	

If a species collection is made, it should be indicated in the collection column with a "C" (for collected). If the species is later keyed out, cross out the species name or description and write the keyed species name in pen on the data sheet. Do not erase what was written in the field, because this information can be used if specimens get mixed up later. If the specimen is then thrown out, the "C" in the collection column should be crossed out. If the specimen is kept but is still not confidently identified, add a "U" to the "C" in the collection column (CU = collected and unconfirmed). In this case the unconfirmed species epithet should be put in parentheses [e.g. *Hordeum (murinum)*]. If the specimen is kept and is confidently identified, add a "C" to the existing "C" in the collection column (CC = Collected and confirmed).

Use Jepson Manual nomenclature. Write out the genus and species of the plant. Do not abbreviate. When uncertain of an identification (which you intend to confirm later) use parentheses to indicate what part of the determination needs to be confirmed. For example, you could write out *Brassica (nigra)* if you are sure it is a *Brassica* but you need further clarification on the specific epithet.

Provide the % absolute aerial cover for each species listed. When estimating, it is often helpful to think of coverage in terms of the following cover intervals at first:

<1%, 1-5%, >5-15%, >15-25%, >25-50%, >50-75%, >75%.

Keeping these classes in mind, then refine your estimate to a specific percentage. All species percent covers may total over 100% because of overlap.

Include the percent cover of snags (standing dead) of trees and shrubs. Note their species, if known, in the “Stand history, stand age and comments” section.

For rapid assessments, make sure that the major non-native species occurring in the stand also are listed in the space provided in the species list with their strata and % cover. For relevés, all non-native species should be included in the species list.

Also for relevés, you can record the <1% cover in two categories: r = trace (i.e., rare in plot, or solitary individuals) and + = <1% (few individuals at < 1% cover, but common in the plot).

Unusual species: List species that are locally or regionally rare, endangered, or atypical (e.g., range extension or range limit) within the stand. This field will be useful to the Program for obtaining data on regionally or locally significant populations of plants.

INTERPRETATION OF STAND

Field-assessed vegetation alliance name: Name of alliance or habitat following the most recent CNPS classification system or the Manual of California Vegetation (Sawyer J.O., Keeler-Wolf T., and Evens, J. 2009). Please use scientific nomenclature, e.g., *Quercus agrifolia* forest. An alliance is based on the dominant or diagnostic species of the stand, and is usually of the uppermost and/or dominant height stratum. A dominant species covers the greatest area. A diagnostic species is consistently found in some vegetation types but not others.

Please note: The field-assessed alliance name may not exist in the present classification, in which case you can provide a new alliance name in this field. If this is the case, also make sure to state that it is not in the MCV under the explanation for “Confidence in alliance identification.”

Field-assessed association name (optional): Name of the species in the alliance and additional dominant/diagnostic species from any strata, as according to CNPS classification. In following naming conventions, species in differing strata are separated with a slash, and species in the uppermost stratum are listed first (e.g., *Quercus douglasii/Toxicodendron diversilobum*). Species in the same stratum are separated with a dash (e.g., *Quercus lobata-Quercus douglasii*).

Please note: The field-assessed association name may not exist in the present classification, in which you can provide a new association name in this field.

Adjacent Alliances/direction: Identify other vegetation types that are directly adjacent to the stand being assessed by noting the dominant species (or known type). Also note the distance away in meters from the GPS waypoint and the direction in degrees aspect that the adjacent alliance is found

(e.g., *Amsinckia tessellata* / 50m, 360° N *Eriogonum fasciculatum* /100m, 110°).

Confidence in Identification: (L, M, H) With respect to the “field-assessed alliance name”, note whether you have L (=Low), M (=Moderate), or H (=High) confidence in the interpretation of this alliance name.

Explain: Please elaborate if your “Confidence in Identification” is low or moderate. Low confidence can occur from such things as a poor view of the stand, an unusual mix of species that does not meet the criteria of any described alliance, or a low confidence in your ability to identify species that are significant members of the stand.

Phenology: Indicate early (E), peak (P) or late (L) phenology for each of the strata.

Other identification problems or mapping issues: Discuss any further problems with the identification of the assessment or issues that may be of interest to mappers. Note if this sample represents a type that is likely too small to map. If it does, how much of the likely mapping unit

would be comprised of this type. For example: “this sample represents the top of kangaroo rat precincts in this general area, which are surrounded by vegetation represented by CARR000x; this type makes up 10% of the mapping unit.”

Is polygon >1 type: Yes / No (circle one): *In areas that have been delineated as polygons on aerial photographs/imagery for a vegetation-mapping project*, assess if the polygon is mapped as a single stand. “Yes” is noted when the polygon delineated contains the field-assessed alliance and other vegetation type(s), as based on species composition and structure. “No” is noted when the polygon is primarily representative of the field-assessed alliance.

If yes, explain: If “Yes” above, explain the other vegetation alliances that are included within the polygon, and explain the amount and location that they cover in the polygon.

Appendix B

Classification Hierarchy

Classification Hierarchy 6-27-2012

1000 = TEMPERATE FOREST SUBCLASS

1100 = California Forest and Woodland Macrogroup MG009

- 1110 = Californian broadleaf forest and woodland Group
 - 1111 = *Quercus douglasii* (Blue oak woodland) Alliance
 - 1112 = *Quercus lobata* (Valley oak woodland) Alliance
 - 1113 = *Quercus chrysolepis* (Canyon live oak forest) Alliance
 - 1114 = *Quercus wislizeni* (Interior live oak woodland) Alliance
 - 1115 = *Juglans californica* (California walnut groves) Alliance
 - 1116 = *Aesculus californica* (California buckeye groves) Alliance
- 1120 = Californian evergreen coniferous forest and woodland Group
 - 1121 = *Pinus sabiniana* (Foothill pine woodland) Alliance
 - 1122 = *Juniperus californica* (California juniper woodland) Alliance

1200 = Californian-Vancouverian Montane and Foothill Forest Macrogroup MG023

- 1210 = Californian montane conifer forest Group
 - 1211 = *Pseudotsuga macrocarpa* (Bigcone Douglas-fir) Alliance

1300 = Intermountain Basins Pinyon-Juniper Woodland Macrogroup MG026

- 1310 = Western Great Basin montane conifer woodland Group
 - 1311 = *Pinus monophylla* (Singleleaf pinyon woodland) Alliance

1400 = Southwestern North American Riparian, Flooded and Swamp Forest Macrogroup MG036

- 1410* = Southwestern North American riparian evergreen and deciduous woodland Group
 - 1411* = *Populus fremontii* (Fremont cottonwood forest) Alliance
 - 1412* = *Salix laevigata* (Red willow thickets) Alliance
 - 1413* = *Salix gooddingii* (Black willow thickets) Alliance
 - 1414* = *Platanus racemosa* (California sycamore woodlands) Alliance
 - 1415* = *Washingtonia filifera* (California fan palm oasis) Alliance
- 1420* = Southwestern North American riparian/wash scrub Group
 - 1421* = *Baccharis emoryi* (Emory's baccharis thickets) Provisional Alliance
 - 1422* = *Baccharis salicifolia* (Mulefat thickets) Alliance
 - 1423* = *Baccharis sergiloides* (Broom baccharis thickets) Alliance
 - 1424* = *Salix exigua* (Sandbar willow thickets) Alliance
 - 1425* = *Forestiera pubescens* (Desert olive patches) Alliance
 - 1426* = *Sambucus nigra* (Blue elderberry stands) Alliance
 - 1427* = *Salix lasiolepis* (Arroyo willow thickets)
- 1430* = Southwestern North American introduced riparian scrub Group
 - 1431* = *Arundo donax* (Giant reed breaks) Semi-natural Stands
 - 1432* = *Tamarix* spp. (Tamarisk thickets) Semi-natural Stands

1500 = Western Cordilleran Montane-Boreal Riparian Scrub and Forest MG034

- 1510* = Vancouverian riparian deciduous forest Group
 - 1511* = *Alnus rhombifolia* (White alder groves) Forest Alliance

* = 1acre MMU ** = 2.5acre MMU

2000 = MESOMORPHIC SHRUB AND HERB CLASS

2100 = California Chaparral Macrogroup MG043

- 2110 = Californian xeric chaparral Group
 - 2111 = *Arctostaphylos glauca* (Bigberry manzanita chaparral) Alliance
 - 2112 = *Adenostoma fasciculatum* (Chamise) Alliance
 - 2113 = *Ceanothus crassifolius* (Hoary leaf ceanothus chaparral) Alliance
 - 2114 = *Fremontodendron californicum* (flannelbush scrub) Alliance
 - 2115 = *Adenostoma fasciculatum* – *Salvia mellifera* (Chamise-black sage chaparral) Alliance
- 2120 = Californian pre-montane chaparral Group
 - 2121 = *Arctostaphylos glandulosa* (Eastwood manzanita) Alliance
 - 2122 = *Ceanothus leucodermis* (Chaparral whitethorn) Alliance
- 2130 = Californian mesic chaparral Group
 - 2131 = *Cercocarpus montanus* (Birchleaf mountain mahogany) Alliance
 - 2132 = *Quercus berberidifolia* (Scrub oak chaparral) Alliance
 - 2133 = *Quercus berberidifolia* – *Adenostoma fasciculatum* (Scrub oak-chamise chaparral) Alliance
 - 2134 = *Prunus ilicifolia* (Holly leaf cherry chaparral) Alliance

2200 = California Coastal Scrub Macrogroup MG044

- 2210 = Central and south coastal California seral scrub Group
 - 2211 = *Gutierrezia californica* (California match weed patches) Provisional Alliance
 - 2212 = *Lotus scoparius* (Deer weed scrub) Alliance
 - 2213 = *Lupinus albifrons* (Silver bush lupine scrub) Alliance
 - 2214 = *Ericameria linearifolia* (Narrowleaf goldenbush scrub) Provisional Alliance
 - 2215 = *Eriodictyon (crassifolium, trichocalyx)* (Thick leaf and hairy yerba santa scrub) Provisional Alliance
 - 2216 = *Malacothamnus fasciculatus* (Bush mallow scrub) Alliance
 - 2217 = *Eriogonum elongatum, nudum* (Longstem buckwheat) Provisional Alliance
 - 2218 = *Corethrogyne filaginifolia* (Common sandaster scrub)
- 2220 = Central and South Coastal Californian coastal sage scrub Group
 - 2221 = *Eriogonum fasciculatum* (California buckwheat scrub) Alliance
 - 2222 = *Eriogonum wrightii* (Wright's buckwheat patches) Alliance
 - 2223 = *Salvia mellifera* (Black sage scrub) Alliance

2300 = California Annual and Perennial Grassland Macrogroup MG045

- 2305 = California annual and perennial grassland Mapping Unit (Native component)
- 2310 = California annual forb/grass vegetation Group
 - 2311 = *Eschscholzia (californica)* (California poppy fields) Alliance
 - 2312 = *Amsinckia (menziesii, tessellata)* (Fiddleneck fields) Alliance
 - 2313 = *Lasthenia californica - Plantago erecta - Vulpia microstachys* (California goldfields - Dwarf plantain - Six-weeks fescue flower fields) Alliance
 - 2314 = *Monolopia (lanceolata)-Coreopsis (calliopsidea)* (*Monolopia* and Tickseed) Alliance
 - 2315 = *Plagiobothrys nothofulvus* (Popcorn flower fields) Alliance
- 2320 = California perennial grassland Group
 - 2321 = *Nassella cernua* (Nodding needle grass grassland) Provisional Alliance
 - 2322 = *Nassella pulchra* (Purple needle grass grassland) Alliance
- 2330 = Mediterranean California naturalized annual and perennial grassland Group

* = 1acre MMU ** = 2.5acre MMU

- 2331 = *Brassica nigra* and other mustards (Upland mustards) Semi-natural Stands
- 2332 = *Bromus rubens* - *Schismus (arabicus, barbatus)* (Red brome or Mediterranean grass grasslands) Semi-natural Stands
- 2333 = *Lolium perenne* (Perennial rye grass fields) Semi-natural Stands
- 2334 = *Pennisetum setaceum* (Fountain grass swards) Semi-natural Stands

3000 = TEMPERATE AND BOREAL SHRUBLAND AND GRASSLAND SUBCLASS (3000)

3100 = Western North American Temperate Grassland and Meadow Macrogroup MG048

- 3110 = Vancouverian and Rocky Mountain naturalized annual grassland Group
- 3111 = *Bromus tectorum* (Cheatgrass grassland) Semi-natural Stands
- 3120 = Western dry upland perennial grassland Group
- 3121 = *Elymus multisetus* (Big squirreltail patches) Provisional Alliance
- 3122 = *Poa secunda* (Curly or one-sided blue grass grassland) Alliance

3200 = Western Cordilleran Montane Shrubland and Grassland Macrogroup MG049

- 3210 = Western Cordilleran montane deciduous scrub Group
- 3211* = *Ribes quercetorum* (Oak gooseberry thickets) Provisional Alliance
- 3220 = Western Cordilleran montane moist graminoid meadow Group
- 3221 = *Muhlenbergia richardsonis* (Mat muhly meadows) Provisional Alliance

3300 = Warm Interior Chaparral Macrogroup MG051

- 3310 = Western Mojave and Western Sonoran Desert borderland chaparral Group
- 3311 = *Ceanothus greggii* (Cup leaf ceanothus chaparral) Alliance
- 3312 = *Quercus john-tuckeri* (Tucker oak chaparral) Alliance
- 3313 = *Quercus palmeri* (Palmer oak) Alliance
- 3314 = *Quercus cornelius-mulleri* (Muller oak chaparral) Alliance

3400* = Western North American Freshwater Marsh Macrogroup MG073

- 3410* = Arid West freshwater emergent marsh Group
- 3411* = *Phragmites australis* (Common reed marshes) Alliance
- 3412* = *Schoenoplectus* spp. (Bulrush) Mapping Unit
- 3413* = *Schoenoplectus acutus* (Hardstem bulrush marsh) Alliance
- 3414* = *Schoenoplectus californicus* (California bulrush marsh) Alliance
- 3415* = *Typha (angustifolia, domingensis, latifolia)* (Cattail marshes) Alliance

3500* = Western North America Vernal Pool Macrogroup MG074

- 3510* = Californian mixed annual/perennial freshwater vernal pool/swale/plain bottomland Group
- 3511 = *Deinandra fasciculata* (Clustered tarweed fields) Alliance

3600* = Western North America Wet Meadow and Low Shrub Carr Macrogroup MG075

- 3610* = Californian warm temperate marsh/seep Group
- 3611* = *Juncus arcticus* (var. *balticus, mexicanus*) (Baltic and Mexican rush marshes) Alliance
- 3612* = *Leymus triticoides* (Creeping rye grass turfs) Alliance
- 3613* = *Muhlenbergia rigens* (Deer grass beds) Alliance

3700* = Warm Semi-Desert/Mediterranean Alkali-Saline Wetland Macrogroup MG083

- 3710* = Southwestern North American alkali marsh/seep vegetation Group
- 3711* = *Spartina gracilis* (Alkali cordgrass marsh) Alliance
- 3712* = *Sporobolus airoides* (Alkali sacaton grassland) Alliance
- 3713 = *Anemopsis californica* (Yerba mansa meadows) Alliance

* = 1acre MMU ** = 2.5acre MMU

3714 = *Juncus cooperi* (Cooper's rush marsh) Alliance
3715 = *Schoenoplectus americanus* (American bulrush marsh) Alliance

3720* = Southwestern North American salt basin and high marsh Group
3721* = *Allenrolfea occidentalis* (Iodine bush scrub) Alliance
3722* = *Atriplex lentiformis* (Quailbush scrub) Alliance
3723 = *Atriplex spinifera* (Spinescale scrub) Alliance
3724* = *Frankenia salina* (Alkali heath marsh) Alliance
3725* = *Suaeda moquinii* (Bush seepweed scrub) Alliance
3726* = *Distichlis spicata* (Salt grass flats) Alliance
3727* = *Salicornia depressa* (Pickleweed flats) Herbaceous Alliance
3728 = *Isocoma acradenia* (Alkali goldenbush) Alliance
3729 = *Atriplex parryi* (Parry's saltbush) Alliance

4000 = WARM SEMI-DESERT SCRUB AND GRASSLAND SUBCLASS

4100 = Mojavean–Sonoran Desert Scrub Macrogroup MG088

4110 = Lower Bajada and Fan Mojavean–Sonoran desert scrub Group
4111 = *Ambrosia dumosa* (White bursage scrub) Alliance
4113 = *Atriplex polycarpa* (Allscale scrub) Alliance
4114 = *Encelia farinosa* (Brittle bush scrub) Alliance
4115 = *Larrea tridentata* - *Ambrosia dumosa* (Creosote bush - white bur sage scrub) Alliance
4118 = *Larrea tridentata* - *Encelia farinosa* (Creosote bush - brittle bush scrub) Alliance
4119 = *Larrea tridentata* (Creosote bush scrub) Alliance
4121 = *Tidestromia oblongifolia* (Arizona honey sweet sparse scrub) Provisional Alliance
4122 = *Pleuraphis rigida* (Big galleta shrub-steppe)
4123 = *Brickellia desertorum* (Desert brickellbush scrub) Alliance

4150 = Arizonan upland Sonoran desert scrub Group
4151 = *Viguiera parishii* (Parish's goldeneye scrub)

4200* = Madrean Warm Semi-Desert Wash Woodland/Scrub Macrogroup MG092

4210* = Mojavean semi-desert wash scrub Group
4211 = *Ephedra californica* (California joint fir scrub) Alliance
4212* = *Lepidospartum squamatum* (Scale broom scrub) Alliance
4213* = *Ericameria paniculata* (Blackstem rabbitbrush) Alliance
4214* = *Prunus fasciculata* (Desert almond) Alliance
4215* = *Brickellia incana* (Woolly brickellia) Provisional wash scrub
4216 = *Ambrosia salsola* (Cheesebush scrub) Alliance
4217* = *Artemisia tridentata* spp. *parishii* (Parish's sagebrush) Provisional Alliance

4220* = Sonoran-Coloradan semi-desert wash woodland/scrub
4221* = *Pluchea sericea* (Arrow weed thickets) Alliance
4222* = *Prosopis glandulosa* (Mesquite bosque, mesquite thicket) Alliance
4223* = *Prosopis pubescens* (Screwbean mesquite bosques) Alliance
4224* = *Chilopsis linearis* (Desert willow woodland)
4225* = *Psoralea argophylla* (Smoke tree woodland)
4226* = *Acacia greggii* (Catclaw acacia thorn scrub)
4227* = *Parkinsonia florida*-*Olneya tesota* (Blue palo verde-ironwood woodland) Alliance
4228* = *Hyptis emoryi* (Desert lavender scrub) Alliance

5000 = COOL SEMI-DESERT SCRUB AND GRASS SUBCLASS

* = 1acre MMU ** = 2.5acre MMU

5100 = Cool Semi-Desert Alkali-Saline Flats Macrogroup MG093

- 5110 = Shadscale-saltbush cool semi-desert scrub Group
- 5111 = *Atriplex canescens* (Fourwing saltbush scrub) Alliance
- 5112 = *Atriplex confertifolia* (Shadscale scrub) Alliance

5200 = Cool Semi-desert wash and disturbance scrub Macrogroup MG095

- 5210 = Intermontane seral shrubland Group
- 5211 = *Encelia (actoni, virginensis)* (Acton's encelia & Virgin River brittle brush scrub) Alliance
- 5212 = *Ericameria nauseosa* (Rubber rabbitbrush scrub) Alliance
- 5214 = *Gutierrezia sarothrae* (Broom snake weed scrub) Provisional Alliance
- 5215 = *Ericameria cooperi* (Cooper's goldenbush) Provisional Alliance
- 5216 = *Dendromecon rigida* (Bush poppy scrub) Alliance

5300 = Western North America Tall Sage Shrubland and Steppe Macrogroup MG096

- 5310 = Inter-Mountain West mesic tall sagebrush shrubland and steppe Group
- 5311 = *Artemisia tridentata* (Big sagebrush) Alliance

5400 = Inter-Mountain Dry Shrubland and Grassland Macrogroup MG098

- 5410 = Intermontane deep or well-drained soil scrub Group
- 5411 = *Grayia spinosa* (Spiny hop sage scrub) Alliance
- 5412 = *Krascheninnikovia lanata* (Winterfat scrubland) Alliance
- 5413 = *Ephedra nevadensis* (Nevada jointfir) Alliance
- 5414 = *Lycium andersonii* (Anderson's boxthorn scrub) Alliance
- 5415 = *Salazaria mexicana* (Bladder sage scrub)
- 5416 = *Ericameria teretifolia* (Needleleaf rabbitbrush scrub) Alliance
- 5417 = *Ephedra viridis* (Mormon tea scrub) Alliance
- 5418 = *Lycium cooperi* (Cooper's boxthorn scrub) Provisional Alliance
- 5420 = Mojave and Great Basin upper bajada and toeslope Group
- 5421 = *Coleogyne ramosissima* (Black brush scrub) Alliance
- 5422 = *Purshia tridentata* (Bitter brush scrub) Alliance
- 5423 = *Yucca brevifolia* (Joshua tree woodland) Alliance
- 5424 = *Yucca schidigera* (Mojave yucca scrub)
- 5425 = *Menodora spinescens* (Greenfire scrub) Alliance
- 5430 = Southern Great Basin semi-desert grassland Group
- 5431 = *Achnatherum speciosum* (Desert needlegrass grassland) Alliance
- 5432 = *Pleuraphis jamesii* (James' galleta shrub-steppe) Alliance
- 5433 = *Achnatherum hymenoides* (Indian rice grass grassland) Alliance
- 5440 = Intermountain shallow/calcareous soil scrub Group
- 5441 = *Cercocarpus ledifolius* (Curl leaf mountain mahogany scrub) Alliance

5500 = Cool Semi-Desert Alkali-Saline Wetlands MG082

- 5510 = Great Basin cool semi-desert alkali basin Group
- 5511* = *Sarcobatus vermiculatus* (Greasewood scrub) Alliance

6000 = NORTH AMERICAN WARM SEMI-DESERT CLIFF, SCREE AND ROCK VEGETATION DIVISION

6100 = North American Warm Semi-Desert Cliff, Scree, and Other Rock Vegetation Macrogroup MG117

- 6110 = North American warm desert bedrock cliff and outcrop Group
- 6111 = *Atriplex hymenelytra* (Desert holly scrub) Alliance
- 6112 = *Ephedra funerea* (Death Valley joint fir scrub) Alliance
- 6113 = Mud Hills sparsely vegetated ephemeral herbs Mapping Unit
- 6114* = Unvegetated wash and river bottom Mapping Unit
- 6115 = Massive sparsely vegetated rock outcrop Mapping Unit

* = 1acre MMU ** = 2.5acre MMU

Appendix B – Classification Hierarchy

- 6116* = Sparsely vegetated playa (Ephemeral annuals) Mapping Unit
- 6117 = Desert pavement Mapping Unit
- 6118 = *Peucephyllum schottii* (Desert fir) Alliance
- 6119 = *Geraea canescens-Chorizanthe rigida* (Desert gold-Spiny herb) Mapping Unit
- 6120 = North American warm desert dunes and sand flats Group
 - 6121 = *Dicoria canescens - Abronia villosa* (Desert dunes) Alliance
 - 6122 = *Panicum urvilleanum* (Desert panic grass patches) Alliance
 - 6123 = *Wislizenia refracta* (Spectacle fruit special stands) Alliance

9000 = MISCELLANEOUS CLASSES

- 9200 = Agriculture (within the current 5-year cycle) (includes nurseries)
 - 9210** = Woody Agriculture (orchards, vineyards)
 - 9220** = Row Agriculture

- 9300** = Built-up & Urban Disturbance
 - 9310 = Urban Window
 - 9320** = Anthropogenic areas of little or no vegetation

- 9500 = Exotic Trees
 - 9501 = Eucalyptus

- 9800* = Water
 - 9801* = Perennial Stream Channel (Open Water)
 - 9803* = Small Ponds and Natural Lakes
 - 9804* = California Aqueduct (Open Water)
 - 9805** = Water Impoundment Feature

* = 1acre MMU ** = 2.5acre MMU

Appendix C

Hierarchical Field and Mapping Key

Hierarchical Field and Mapping Key to the 2012 DRECP Vegetation Map

This key is developed for areas mapped between February 2011 and July 2012 in the Desert Renewable Energy Conservation Plan (DRECP) footprint (Figure 1). It only includes those areas of the planning area and is intended for use as a guide to identification of field-based and image interpretation-based vegetation assessments.

Due to the high diversity of the vegetation communities in the area, this is a complex key. You will need to collect or refer to plant composition data that includes not only those species that are dominant but also those "indicator," or characteristic species, whose presence may cause the plot to key to another vegetation type. If you are using this key for mapping rules please also note that some of the types are typically below the accurate detectability for mapping in this project.

Terms and Concepts Used Throughout the Key (please read prior to use):

Stand: The basic physical unit of plant communities in a landscape. It has no set size. Some vegetation stands are very small, such as certain wetland types, and some may be several square kilometers in size, such as certain forest types. A stand is defined by two main unifying characteristics:

1. It has compositional integrity. Throughout the stand, the combination of species is similar. The stand is differentiated from adjacent stands by a discernible boundary that may be abrupt or occur indistinctly along an ecological gradient.
2. It has structural integrity. It has a similar history or environmental setting that affords relatively similar horizontal and vertical spacing of plant species. For example, a hillside forest originally dominated by the same species that burned on the upper part of the slopes but not the lower would be divided into two stands. Likewise, a sparse woodland occupying a slope with very shallow rocky soils would be considered a different stand from an adjacent slope with deeper, moister soil and a denser woodland or forest of the same species.

The structural and compositional features of a stand are often combined into a term called homogeneity. For an area of vegetated ground to meet the requirements of a stand, it must be homogeneous at the scale being considered. The DRECP mapping project has a variable Minimum Mapping Unit (MMU) size. For fine scale features such as wetlands, riparian, or playas it is 0.4 ha (1 ac) and for upland vegetation it is 4 ha (10 ac). Certain types of upland vegetation are only found in washes (e.g., associations of *Larrea tridentata* alliance with *Ambrosia salsola*, or *Atriplex polycarpa* alliance). In these cases, these vegetation types are mapped at a MMU of 1 ha (2.5 ac) in order to maintain the wash vegetation features separate from surrounding upland. Special types are mapped to 0.4 ha (1 ac). The foothills of the Transverse Ranges are mapped to 1 ha (2.5 ac), as are land use polygons, in keeping with state-wide mapping standards.

Alliance: Plant communities based on dominant/diagnostic species of uppermost or dominant stratum. Part of the United States National Vegetation Classification (USNVC) hierarchy.

Association: The most botanically detailed plant community designation based on dominant species and multiple co- or subdominant indicator species from any strata. Part of the USNVC hierarchy.

Plant community nomenclature: Species separated by "-" are within the same stratum; species separated by "/" are in different strata. The number that precedes some plant community names is the Mapping Code used for labeling plant community polygons for the associated GIS-based plant community map.

Cover: The primary metric used to quantify the importance/abundance of a particular species or a particular vegetation layer within a stand. It is measured by estimating the aerial extent of the living

plants, or the bird's-eye view looking from above, for each category. Cover in this mapping project uses the concept of "porosity" or foliar cover rather than "opacity" or crown cover. Thus, field crews and aerial photo interpreters are trained to estimate the amount of shade produced by the canopy of a plant or a stratum by taking into account the amount of shade it casts excluding the openings it may have in the interstitial spaces (e.g., between leaves or branches). This is assumed to provide a more realistic estimate of the actual amount of shade cast by the individual or stratum which, in turn, relates to the actual amount of light available to individual species or strata beneath it. However, as a result cover estimates can vary substantially between leaf-on versus leaf-off conditions.

*If there is a species present in high cover for which no type exists in the key, there are two options. First, the plot can key to another species that is present in high cover. For example, a plot with 6 percent cover *Senna armata* and 4 percent *Ambrosia salsola* would key to *Ambrosia salsola*, since there is no *Senna armata* type defined in the study area. If this is not a reasonable option, the plot can be designated "unable to key." Plots that are unable to key may be candidates for new vegetation types especially if similar stands are seen to repeat in the landscape. In addition to *Senna armata*, several other woody species may dominate a stand but may not be keyable here, including: *Tetradymia* spp. (including *T. stenolepis* and *T. fasciculatus*), *Lepidium fremontii*, and *Fraxinus velutina*. Comments are inserted in the likely places in the key to address not as yet formally designated vegetation types, which may be dominated by such species.*

Absolute cover: The actual percentage of the surface area of the plot that is covered by a species or physiognomic group (trees, shrubs, herbaceous), as in "creosote bush covers 10 percent of the plot." Absolute cover of all species or physiognomic groups, when added together, may total greater than 100 percent, because this is not a proportional number and plants can overlap each other. For example, a plot could have 25 percent tree cover, 40 percent shrub cover, and 50 percent herbaceous cover.

Relative cover: The percentage of the surface area of the plot that is covered by one species or physiognomic group (trees, shrubs, herbaceous) as compared or relative to the amount of surface of the plot covered by all species or groups. Thus, 50 percent relative cover means that half of the total proportion of cover of all species or physiognomic groups is composed of the single species or group in question. Relative cover values are a proportional number that, when added together, total 100 percent for each sample or stand. For example, a creosote bush–burro bush vegetation plot with 5 percent cover creosote bush and 5 percent cover burro bush estimated using absolute cover would translate to 50 percent relative cover of creosote bush and 50 percent relative cover of burro bush.

Dominance: Dominance refers to the preponderance of vegetation cover in a stand of uniform composition and site history. It may refer to cover of an individual species as in "dominated by creosote bush," or it may refer to dominance by a physiognomic group, as in "dominated by shrubs." When we use the term in the key a species is dominant if it is in at least 70 percent of the stands of this type, with at least 50 percent relative cover in each stand, however, see "dominance by layer," below.

Strongly dominant: 60 percent+ relative cover. A species in the dominant life form stratum has 60 percent or greater relative cover.

Codominant: Each species has 30 percent–60 percent relative cover. Codominance refers to two or more species in a stand with near equal cover. In general, codominance can occur among species that have between 30 and 60 percent relative cover each. **To be codominant species** should be in at least 70 percent of the stands of this type, with at least 30 percent relative cover in each stand. For example in a desert scrub stand with 5% *Larrea tridentata*, 3% *Ambrosia dumosa*, and 4 % *Ephedra nevadensis* (total 13% shrub cover), technically only the *Larrea* ($5/13 = 39\%$ relative cover) and the *Ephedra* ($4/13 = 31\%$ relative cover) would be codominant, even though the stand would key out to *Larrea tridentata* - *Ambrosia dumosa* alliance (see rules for *Larrea tridentata* – *Ambrosia dumosa* in key below.).

Consistent/Characteristic/Diagnostic species: Should be present in at least 75 percent of the stands of the type, with no restriction on cover.

Abundant species: Should be present in at least 50 percent of the samples, with an average of at least 30 percent relative cover in all samples.

Dominance by layer: Tree, shrub, and herbaceous layers are considered physiognomically distinct. A vegetation type is considered to belong to a certain physiognomic group if it is dominated by one layer. Layers are prioritized in order of height. The tallest layer, if it meets a criterion in the "characterized" definitions (see below) is said to dominate, and the type is usually named at the alliance level by the characteristic species of the tallest layer. Average covers within the dominant layer reflect the "modal" concept of the characteristics of a particular vegetation type. For example, a higher average cover of woody plants within a stand not recently affected by disturbance reflects a mode of general availability of water, nutrition, and equitable climate, while lower average cover under similar conditions would reflect lower availability of these things.

Layer dominance concepts are relative to higher levels in the classification that are driven by regional climate (usually from Macrogroup to Formation levels). This is an important concept in the mapping area where, for example, desert shrublands meet California Mediterranean climate shrublands or montane woodlands. Although rules within the Mojave-Sonoran desert may discuss *Yucca brevifolia* having a threshold membership rule of >1% cover, evenly distributed, when *Y. brevifolia* occurs over a much more dense and evenly distributed sclerophyllous shrub cover of *Adenostoma fasciculatum* or *Quercus john-tuckeri*, that rule does not apply and that stand would key to the best characteristic species of the shrub layer. In order to be keyed to a *Y. brevifolia* alliance, such a stand would have to contain at least 10% cover of *Y. brevifolia* over the sclerophyll layer, since for wetter non-desert environments the rule for tree layer dominance is $\geq 10\%$ tree cover. This also applies to situations where *Pinus monophylla* occurs over chaparral as in portions of the foothills of the Transverse Ranges, since *P. monophylla* alliance is diagnostic of the Intermountain Singleleaf Pinyon - Western Juniper Woodland Macrogroup. Although *P. monophylla* may only need to be > 1% cover in desert vegetation, it would need to be > 10% when present in predominantly sclerophyllous Mediterranean scrub, which regularly has > 25% shrub cover in a stand.

Plant Dispersion (sociability) in semi deserts and sparsely vegetated landscapes:

For all desert vegetation, an even distribution of species in the dominant layer is an important factor in correctly identifying alliances and associations. Whatever the dominant overstory layer, the diagnostic species in that layer should be evenly distributed across the stand being assessed. This applies to riparian stands characterized by willows, cottonwoods, and palms; desert scrub with species such as creosote bush, saltbush, or encelia; or herb-land/grassland stands with species such as *Coreopsis*, *Amsinckia*, *Eriogonum*, *Pleuraphis*, or *Bromus*. Therefore, when using this key in the field or with high resolution aerial imagery, it is important to assess not just the estimated cover of the diagnostic species in their layer(s), but also reflect upon their dispersion within the stand.

Relatively even spacing throughout the stand is important particularly in vegetation with low total cover since an even distribution of the diagnostic species is a much better indicator than overall cover, as this may only vary a few percent between diagnostic and associated species. Irregular distribution of species suggests a history of recent disturbance and makes precise determination more difficult. In some cases due to disturbance such as recent fire or clearing, desert vegetation may not be identifiable to alliance and can only be described at higher levels such as group or macrogroup. In other cases early seral vegetation also has diagnostic opportunistic species such as *Ambrosia salsola*, *Ericameria cooperi*, *E. nauseosa*, or *Encelia actoni*. They may quickly colonize and form stands that will begin, if left undisturbed for several years, to shift to a more stable and structurally diverse stand of a different alliance. The key is arranged in the general order of the USNVC hierarchy for situations like this; it reflects such ecological relationships.

Increaser: A plant species (usually shrubs as used in this key) that tends to increase in numbers and in relative cover following fire, prolonged grazing, or other disturbance.

Estimating cover using actual percentages, rather than cover classes, is preferable, because it gives the fullest picture of the vegetation present. It enables later review of the data to confirm choice of plant community and may help to describe new vegetation types and answer future management or research questions. If a less rigorous and faster approach is needed, for example, if the project is not primarily a vegetation project, the following cover classes are compatible with the key and may be used:

1. <1 percent
2. 1–5 percent
- 3a. 6–15 percent
- 3b. 16–25 percent
4. 26–50 percent
5. 51–75 percent
6. 76–100 percent

All references to percent cover in the key are to absolute cover unless specified in a particular section as relative cover.

Diagnostic species: A species typically found in the dominant stratum of a vegetation type often lending its name to that association or alliance due to its constancy and reliable presence throughout most similar stands.

Significant: A species which has 1 to 5 percent absolute cover.

Important: A species or species group with >1 percent absolute cover. A species is considered important if it has greater than 1 percent absolute cover. This term is contrasted with dominant to mean that the species is always present in greater than 1 percent cover but not always dominant. All significant species are therefore, important, but species with higher importance may also be codominant or dominant.

Sparse: Used to describe individual layers of vegetation (tree, shrub, herb, or subdivisions of them) where the cover is less than 2 percent absolute cover.

Sparse vegetation: Neither vascular plants nor nonvascular organisms provide a consistent structural component or play an important role in ecological processes on the site. For the desert this is usually below 2% absolute cover in combination with an irregular uneven distribution across the landscape.

Woody plant: Is any species of plant that has noticeably woody stems. It does not include herbaceous species with woody underground portions such as tubers, roots, or rhizomes.

Tree: A one-stemmed woody plant that normally grows to be greater than 5 meters tall. In some cases, trees may be multiple stemmed following ramifying after fire or other disturbance, but the size of mature plants is typically greater than 5 meters. Undisturbed individuals of these species are usually single stemmed. Certain species that resemble shrubs most of the time in our study area but may be trees in other areas (e.g., *Juniperus californica*) are, out of state-wide tradition, called trees, even though conversely tall shrubs such as *Quercus john-tuckeri* may be equally as tall, but are never called trees. It behooves one to memorize which species are “traditionally” placed in one life-form or another. We use the NRCS Plants Database to do this.

Tree-characterized vegetation: Trees are evenly distributed throughout the stand. In the Mediterranean climate margins of the desert, as in the San Gabriel, San Bernardino, Sierra Pelona, Liebre, or Tehachapi Mountains, trees have typically ≥ 10 percent cover, providing a consistent structural component. In the true desert, short trees such as pinyon pine, Joshua tree, or California

juniper are diagnostic at lower than 10% cover. Typically 2 or even 1 percent, of evenly distributed diagnostic tree species is all that is needed, if one or both of the following criteria are met: (1) trees influence the distribution or population dynamics of other plant species; (2) trees play an important role in ecological processes within the stand.

Forest: In the USNVC, a forest is defined as a tree-dominated stand of vegetation with 60 percent or greater cover of trees. Most forest alliances tend to have average cover of trees > 60%, but individual stands under certain conditions may drop lower than 60 percent.

Woodland: In the USNVC, woodland is defined as a tree-dominated stand of vegetation with between 25 percent and 60 percent cover of trees. The same notion of "modality" that applies to forest types also applies here and to the sparsely wooded category.

Sparsely wooded: These are stands with conspicuous trees at generally at least 10% absolute cover but less than 25 percent cover, and may occur over shrubs as the dominant canopy (sparsely wooded shrubland) or herbaceous cover (sparsely wooded herbaceous).

Emergent: A plant (or vegetation layer) is considered emergent if it has a low cover and rises above a layer with has most of the cover in the stand. For example, individual *Quercus lobata* trees may comprise an emergent tree layer of 5 percent over a dense layer of *Artemisia californica* shrubs; the stand would be considered within the *Artemisia californica* Shrubland Alliance because the total tree cover is < 10% and the shrub cover is > 10%. Further, medium to tall shrubs are not considered emergent over shorter shrubs, but short trees are considered emergent over tall shrubs. For desert vegetation, which is inherently sparse, the threshold is lower. Trees such as pinyon pine, California juniper, and Joshua tree can be evenly distributed at as low as 1-2% and instead be considered the dominant members of the canopy because the shrub or herb vegetation cover is usually low.

Joshua Tree and "Microphyll" "woodland". *Yucca brevifolia* is iconic and typically substantially taller than most other woody plants of the Mojave Desert. It is considered a tree even in its short clonal form that is typical of the western desert margins. Even though woodlands outside of the desert are rarely considered such when the tree cover is less than 10%, a stand can be considered Joshua tree woodland when Joshua trees are evenly distributed and have at least 1% cover.

Colorado Desert microphyll woodland species such as *Olneya tesota* and *Parkinsonia florida*, even at their best "woodland" development, are rarely over 10% cover throughout a stand though they are visually and structurally conspicuous. Even at 5% absolute cover they define the stands, especially since any other component woody species typically have less cover than these trees.

Shrub: Usually a multi-stemmed woody plant that is between 0.2 meter and 5 meters tall. Definitions are blurred at the low and high ends of the height scales. At the tall end, shrubs may approach trees based on disturbance frequencies (*e.g.*, old-growth resprouting chaparral species such as *Cercocarpus montanus*, *Fremontodendron californica*, *Prunus ilicifolia*, and so forth, may frequently attain "tree size"). At the short end, woody perennial herbs or subshrubs of various species are often difficult to categorize into a consistent life-form.

Sub-shrub: A multi-stemmed plant with noticeably woody stems less than 0.5 meter tall.

Shrub-characterized vegetation: Shrubs (including sub-shrubs) are evenly distributed throughout the stand, providing a consistent (even if sparse) structural component, the stand cannot be characterized as a tree stand, and one or both of the following criteria are met: (1) shrubs influence the distribution or population dynamics of other plant species; (2) shrubs play an important role in ecological processes within the stand.

Herbaceous plant: Any species of plant that has no main woody stem development and includes grasses, forbs, and perennial species that die back each year.

Herb-characterized vegetation: Herbs are evenly distributed throughout the stand, providing a consistent (even if sparse) structural component, and play an important role in ecological processes within the stand, and the stand cannot be characterized as a tree or shrub stand.

Nonvascular vegetation: Nonvascular organisms provide a consistent (even if sparse) structural component and play an important role in ecological processes within the stand.

Botanical nomenclature: Note that we use the NRCS PLANTS database in vegetation mapping as our standard for botanical names. [In the Final version of this report, users of the Jepson Manual 2nd Edition (Baldwin et al 2012) will find a list of name changes at the end of this key].

Use of the Key

This key is constructed in a traditional dichotomous style, with couplets of opposing choices. Exceptions are in the most diverse groups where dichotomies split the main subdivisions of alliances and then a small number of associations may be simply listed below the subdivisions with diagnostic characteristics. The key is blind to any artificial division between trees, shrub, and herbaceous dominated vegetation. Instead it follows more closely the new National Vegetation Classification hierarchy (Faber-Langendoen et al. 2009) promoted by the Ecological Society of America's Vegetation Panel and the Federal Geographic Data Committee (Peet et al. 2008).

To underscore the relationships of different vegetation in the Western Mojave, bolded text (in addition to Alliance and mapping unit names) has been inserted to show hierarchical position at important ecological breaking points given in the key. Aside from the main concepts of Alliance and Association previously mentioned above, the other hierarchy units are (from highest to lowest) are: **1. Class, 2. Subclass, 3. Formation, 4. Division, 5. Macrogroup, and 6. Group.**

The vegetation map includes some mapping units that are either human created or are not vegetation at all. These may be individuated from the natural vegetation below by using this short key, below.

KEY TO NATURAL AND ANTHROPOGENIC LAND COVER

1. Landscape unit consists of water bodies

9800 = Water

2. Open waters of naturally flowing streams and rivers...

9801 = Perennial Stream Channel (Open Water)

2'. Water either naturally contained and not flowing, or contained by anthropogenic means (canals, reservoirs or aqueducts)

3. Water contained by earthen dams and/or natural landscape features...

9803 = Small Dammed Ponds and Natural Lakes

3'. Water contained entirely by pavement, concrete, or by human constructed earthen walls...

4. Water in the conveyance known as the California Aqueduct...

9804 = California Aqueduct (Open Water)

4'. Water surrounded on all sides by human-constructed walls...

9805 = Water Impoundment Feature

1.' Landscape is not water...

5. Landscape is heavily modified by human activity. This constitutes such things as recently planted agricultural crops, broad extents of residential and industrial buildings, and areas that have been physically denuded of vegetation. In the vegetation mapping classification these are generically termed Miscellaneous Map Classes (the “9000” codes). In the National Vegetation Classification System these would include in part the classes called Hortomorphic (gardens and other plantings for aesthetic purposes) and Agromorphic (all forms of agriculture including annual row crops, vineyards, orchards, and timber plantations)...

6. Landscape is vegetated by human-plantings for aesthetic purposes, food, fiber, or building materials production...

7. Vegetation is confined to active agriculture (defined as planted and maintained for no more than 5 years earlier than the 2010 NAIP base imagery date. May be dominated by annual or perennial types...

9200 = Agriculture (within the current 5-year cycle) (includes nurseries)

8. Vegetation defined by woody vegetation such as orchards and vineyards.

9210 = Woody Agriculture (orchards, vineyards)

8'. Vegetation defined by annual herbaceous vegetation.

9220 = Row Agriculture

7'. Vegetation confined to aesthetic horticultural plantings of trees not grown for harvest for food or other products...

9500 = Exotic Trees

9. Vegetation consisting of planted trees of the genus *Eucalyptus*...

9501 = Eucalyptus

9'. Vegetation consisting of planted trees of other species...
(unit not defined for this project)

6'. Landscape is unvegetated or consists of a matrix of development and small patches of natural or human planted vegetation...

9300 = Built-up & Urban Disturbance

10. Landscape unit is not vegetated but cleared by humans...

9320 = Anthropogenic areas of little or no vegetation

10'. Landscape is defined by a minimum of 10 acres containing a matrix of buildings and small natural or anthropogenic vegetated or unvegetated opening between buildings. Commonly called suburban areas or housing developments...

9310 = Urban Window

5'. Landscape is sparsely to very well-vegetated with naturally growing (not planted and heavily tended) vegetation ...

Use key to natural and semi-natural vegetation, below.

KEY TO NATURAL AND SEMINATURAL VEGETATION

1. Vegetation largely absent and no species are evenly distributed. Vegetation not uniformly distributed across a landscape surface, generally less than 5% cover, not composed of evenly-spaced trees or shrubs, or not characterized by herbaceous species most of the time. Following seasons of exceptional precipitation, herbaceous annual species may be abundant and evenly distributed...

Class 6 Lithomorphic Vegetation (Nonvascular and Sparse Vascular Rock Vegetation)

Subclass 6.C Semi-Desert Nonvascular and Sparse Vascular Vegetation

Formation 6.C.1 Warm Semi-Desert Cliff, Scree, and Rock Vegetation

Division 6.C.1.a North American Warm Semi-Desert Cliff, Scree, and Rock Vegetation

6100 = North American Warm Semi-Desert Cliff, Scree, and Other Rock Vegetation

Macrogroup MG117

6110 = North American warm desert bedrock cliff and outcrop Group

2. Landscape characterized by open dunes, dune aprons, or sand flats. Vegetation is generally sparse to very open (< 2- 10% cover) except for annual blooms in favorable years. [Group - North American warm desert dunes and sand flats.] May include the following possible alliances, although local indicator species may be different (e.g., *Cryptantha angustifolia*, *Camissonia claviformis*, and *Oenothera deltoides*)...

3. *Dicoria canescens* or *Abronia villosa* are characteristically present in stands, but are not necessarily dominants, depending upon the year and the phenology of these annual plants. Skeletons of *Dicoria*, *Oenothera deltoides*, *Abronia villosa* and other psammophytic annuals are usually present and uniform woody plant cover is < 2% absolute cover...

6121 = *Dicoria canescens* - *Abronia villosa* (Desert dunes) Alliance

3'. *Dicoria canescens* or *Abronia villosa* (or other annual herbs) not characteristically present or present in less cover and less uniform distribution than taller perennial sand-loving herbs or grasses...

4. Stands are characterized by even, sparse distribution of the stoloniferous dune panic-grass (*Panicum urvilleanum*). Restricted locally to broad sandy riverbed and adjacent low dunes of the Mojave River from Hinkley to Camp Cady...

6122 = *Panicum urvilleanum* (Desert panic grass patches) Alliance

4'. Stands dominated by the tall perennial herb *Wislizenia refracta*. Found associated with low dunes adjacent to Palen Dry Lake, often adjacent to open stands of *Atriplex canescens* or *Suaeda moquinii* scrubland...

6123 = *Wislizenia refracta* (Spectacle fruit special stands)

2'. Landscape characterized by desert pavement, pediment, badlands, playa margins, or outcrops with ≤ 5 % cover of perennial plant species, none of which are particularly evenly distributed. During exceptional years there may be significant annual ephemeral species cover (which may be evenly distributed). Species are not always present, but on non-sand substrate can include *Plantago ovata*, *Cryptantha angustifolia*, *Chorizanthe rigida*, and *Geraea canescens*, etc...

5. Stands are characterized by very low shrub cover, and are low in total absolute cover (usually 1-5%) of all layers...

6. *Atriplex hymenelytra* > 1% cover and no other woody species with equal or higher cover. May occur on hot rocky slopes, dry bajadas, or alkaline badlands and playa edges. Stands are local in the extreme north of the mapped area near Ridgecrest and Trona on alkali basin sediments and more extensive in the Calico Mountains and in the Alvord Mountains on volcanic ash and flows emanating from the southern and eastern sides of these ranges. Stands are also known from the altered volcanic hills southeast of Barstow and west of Daggett. Stands are generally considered "sparsely vegetated" (mostly < 2% shrub cover). However, some stands can have up to 10% shrub cover under certain circumstances. Stands that are codominated by *Atriplex confertifolia* are *Atriplex confertifolia* alliance...

6111 = *Atriplex hymenelytra* (Desert holly scrub) Alliance

6'. Vegetation dominated or characterized by a low total cover of *Ephedra funerea* or *Peucephyllum schottii*, with no other indicator shrub species present in greater cover or dispersion...

7. An uncertain alliance found in calcareous mountains, or rarely, other shallow volcanic slopes in the study area. Often accompanied by *Echinocactus*, *Ferocactus*, or *Echinocereus* spp. One or two reconnaissance points have been taken for this alliance in the eastern portion of our study area. It has not been mapped...

6112 = *Ephedra funerea* (Death Valley joint fir scrub) Alliance

7'. Stands typical of steep massive outcrops of basalt or calcareous rocks (cliffs and scree) at lower elevations throughout the study area. The bright green arborescent shrub (which may resemble *Larrea* at a distance, beware) *Peucephyllum schottii* is characteristic throughout the stand, but is often only 1-5% total cover...

6118 = *Peucephyllum schottii* (Desert fir) Alliance

5.' Stands usually do not support even a sparse even distribution of shrubs. Cover is either apparently lacking entirely, or in good precipitation years, distinguished by ephemeral blooms of annual plants. Substrates vary from hills, mountains, playas and riverbeds...

8. The substrate is low-lying with little topography; either desert playa or wash and river bed channels of sand, cobbles, silt, clay, or salty or alkaline mineral deposits...

9. This mapping unit is distinguished by largely unvegetated sands and gravels in the active centers of washes throughout the study area. Depending upon site history and recent flooding events these bare "river-wash" channels can change rapidly and regularly from unvegetated to vegetated by annual natives, to vegetated by shrub species indicative of washes of different flooding frequencies and intensities. Mapped usually as scattered shrubs and herbs with < 2% average cover and uneven distribution...

6114 = Unvegetated wash and river bottom Mapping Unit

9'. This mapping unit defines silty, clay, or salt crust playa (dry lake) surfaces throughout the study area. Most of the time playas, whether salty, silty, or clay, are usually < 2% vegetative cover. However, annuals such as *Monolepis nuttalliana*, *Atriplex elegans*, *A. phyllostegia*, and others may occur in good El Niño years in relatively high cover. These are still mapped as this unit. Characteristics include flat or cracked substrate and no obvious slope. Reflectance may be gray to white to light brown...

6116 = Sparsely vegetated playa (Ephemeral annuals) Mapping Unit

8'. The substrate is part of an elevated landscape that usually has hills, slopes, outcrops, or ravines...

10. The surface of the stand is usually gradually sloping, but often characterized by small cobbles or rock fragments, the result of long-term weathering of alluvial fans/bajadas at the bases of mountains...

11. Extremely sparse, dark, and unvegetated to a large degree even in good rainfall years, photo signature is usually distinctive and often looks like asphalt pavement. No evenly spaced shrubs are present. Look for evidence of old dark alluvial surfaces with no shrub cover and gradual to moderate slopes. If the slopes are steep, choose 6115 and if highly dissected and fine-textured choose 6113. Best examples are in the eastern part of the mapping area where these are extensive and diagnostic of this part of the Colorado Desert. Farther west the rainfall average is higher and the quality and extent of this landform are less pronounced. Good examples occur adjacent to Pinto Basin and some also occur in Dale Lake basin. Modeling for mapping suggests using this mapping unit only in the driest part of the mapping area, generally east of the longitude of Barstow and Lucerne Valley...

6117 = Desert pavement Mapping Unit

11'. Surfaces usually have evidence of some herbaceous cover with the indicator herbs *Chorizanthe rigida*, and/or *Geraea canescens* either as residual skeletons or in good rainfall years visually obvious displays characterized by these native annuals. Moving westward from the eastern portions of the mapping project, old alluvial surfaces tend to be better vegetated. Those that resist shrub colonization tend to at least have good stands of *Geraea canescens-Chorizanthe rigida* and sometimes other annuals in good El Niño years. Stands that have other types of herbaceous vegetation including *Lasthenia gracilis*, *Amsinckia* spp., or non-native annual herbs such as *Bromus rubens* should be mapped to their respective alliances or semi-natural stands and not as desert pavement or this alliance...

6119 = *Geraea canescens-Chorizanthe rigida* (Desert gold-Spiny herb) Provisional Alliance

10'. Stands defined by (usually) steeply to moderately sloping landforms that can be considered hills, badlands, mountains, or outcrops of resistant rock. Vegetation is sparse or at best characterized by sporadic occurrences of ephemeral blooms of annuals following good rainfall...

12. This mapping unit is usually sparsely vegetated with < 2% shrub cover or herb cover. Substrate is composed of unconsolidated and uncemented fine, sometimes alkaline, sediments. Strata variations make for different but typically sparsely vegetated slopes. Averaging all cover across these landscapes is usually what constitutes > 10 acre mapping polygons, which consist of a matrix of small patches of shrubs or herbs at 2% interspersed with larger areas of little or no measurable cover of herbs or shrubs. Topography is often rugged and eroded ("badlands"), however, in some areas, such in the northern portion of the study area, the same species may occur on relatively flat terrain adjacent to edges of playas and in broad valleys. In many years these areas are largely unvegetated. Some have scattered *Atriplex hymenelytra*, *Atriplex confertifolia*, *Stanleya pinnata* and other woody species. In El Niño years clay slopes are

heavily covered with annual *Eriogonum* species, probably most commonly *E. inflatum* (many sizes of this plant). This and other species of *Eriogonum*, along with *Plantago ovata*, *Chorizanthe* species, and sometimes *Lepidium lemmonii*, *Coreopsis calliopsidea*, and other species can lend noticeable color to these exposures...

**6113 = Mud Hills sparsely vegetated ephemeral herbs
Mapping Unit**

12'. This mapping unit is defined by extensive solid blocks of resistant rock of any type. In our area these may be volcanic extrusives such as basalt or rhyolite; igneous intrusives such as granodiorite, gabbro, or quartz monzonite; or sedimentary sandstones or limestones, etc. Large unfractured bedrock or boulders are typical, with narrow crevices in different densities. Overall shrub and herb cover tends to be under 5%, making it difficult to distinguish any particularly dominant species. Mapping units of this type may include small (< 10 acre) stands of *Ephedra viridis*, *Atriplex polycarpa*, *Encelia farinosa*, *Ericameria cuneatus*, *E. teretifolia*, *Salazaria mexicana*, *Eriogonum fasciculatum*, and other alliances. Separate this from mud hills by erosional patterns (should see individual rock outcrops, boulders, etc) in this unit...

**6115 = Massive sparsely vegetated rock outcrop
Mapping Unit**

1'. Vegetation easily visible and $\geq 5\%$ total cover (sometimes as low as 2% if evenly distributed) and characterized by trees, shrubs and/or herbs that are evenly distributed across the stand.

13. Vegetation dominated by broad-leaved or needle-leaved trees (or if in riparian, larger winter deciduous shrubs, such as *Salix* spp.) generally over 10% absolute cover. In the California desert, tree vegetation is typically taller than the average creosote-bush (e.g., > 3 m tall). Characteristic species of the dominant layer are either winter deciduous or evergreen and do not exhibit xeromorphic characteristics such as thorns, drought-deciduousness, succulent stems or microphyllous leaves/leaflets. If growing in the desert ecoregion, growing in arroyos, washes, canyon bottoms, springs, seeps, or other areas that receive and retain more moisture than ambient desert settings...

1.C. Temperate Forest and Woodland Subclass

14. Vegetation is dominated by broad-leaved or needle-leaved trees or tall shrubs (such as willows). Not in uplands or dry washes, but in areas where moisture is present at least under the surface in the warmer months (near permanent surface or subsurface moisture). The majority of stands in the study area are riparian and the only truly flooded stands appear at the margins of permanent reservoirs in Palmdale and the western Antelope Valley...

Formation 1.C.3. Temperate Flooded and Swamp Forest

Division 1.C.3.c Western North America Warm Temperate Flooded and Swamp Forest

**1400 = Southwestern North American Riparian, Flooded and Swamp Forest
Macrogroup MG036 (M036 Warm Mediterranean & Desert Riparian, Flooded & Swamp Forest)**

15. Stands are dominated or characterized by riparian winter deciduous, broad-leaved trees or tall shrubs, including *Populus fremontii*, *Platanus racemosa*, and/or a species of *Salix*. If the tree canopy is between 5 and 10 percent absolute cover, the shrub canopy should not be more than double the tree canopy (otherwise, see 15'). Note that all diagnostics in this

macrogroup and group are considered as such even as saplings when similar in size to mature individuals of shrubby *Salix*, (e.g., *S. lasiolepis* and *S. exigua*.) Thus, if the stand has > threshold cover for indicators of this group as saplings, even if there is similar or greater cover of shrub willow species, the stand would key to the tree type...

1410 = Southwestern North American riparian evergreen and deciduous woodland Group

16. *Populus fremontii* is dominant or codominant with >5% absolute cover in the tree canopy. Stands occur along streams, springs, and valleys with a subsurface water supply, and may be mapped to small clumps less than 1 acre. *P. fremontii* occurs with *Salix*, *Forestiera*, *Baccharis*, etc. Stands codominated by tree willows such as *Salix gooddingii* or *S. laevigata* will key here. *P. fremontii* > 5%, *Platanus*, *Salix laevigata*, each usually < 5%; *S. gooddingii* may be codominant, shrubby *Salix lasiolepis*, *S. exigua*, or *Baccharis* spp. may be present at low to high cover in the understory...

1411 = *Populus fremontii* (Fremont cottonwood forest) Alliance

16'. *Populus fremontii* is not dominant or codominant...

17. *Salix laevigata* is the sole dominant in the overstory layer with at least 10% cover. Arroyo willow (*Salix lasiolepis*) may occur as a sub- or codominant in the shrub or low tree layer. If present, *S. lasiolepis* < *S. laevigata* and *Platanus racemosa*, *Populus* spp., and *S. gooddingii* are all trace. Usually small stands associated with isolated springs and seeps, may be associated with *P. fremontii* or shrubby riparian stands of *Forestiera*, *S. exigua*, etc...

1412 = *Salix laevigata* (Red willow thickets) Alliance

17'. *Salix laevigata* is not dominant...

18. *Salix gooddingii* is the strongly dominant tree, while other tall woody shrubs may be subdominant. *S. gooddingii* > trace; if present, *Platanus racemosa* and *Populus* spp. are both trace. Most stands are associated with the Mojave River or are in permanently wet areas in the Antelope Valley. Often adjacent to *P. fremontii* stands...

1413 = *Salix gooddingii* (Black willow thickets) Alliance

18'. *Salix gooddingii* is not dominant...

19. *Platanus racemosa* is characteristic of the riparian tree canopy. *Platanus racemosa* > trace; if present, *Populus* spp. < *P. racemosa*. Along major stream courses, often associated with other stands of trees or shrubs within this macrogroup. Individuals of *P. racemosa* occur as far north as Victorville in the Mojave, but actual stands are limited to within a few miles of the edge of the ecoregion as at Rock Creek, Little Rock Creek, and several drainages

emerging from the Tehachapi Mountains. Stands are common in the Cajon Pass area...

1414 = *Platanus racemosa* (California sycamore woodlands) Alliance

19'. Either *Alnus rhombifolia* or *Washingtonia filifera* dominant...

20. Stands characterized by even distribution of the California Fan Palm (*Washingtonia filifera*), associated with springs and moist canyon bottoms in a few places. Other riparian tree species (*Populus fremontii*, *Salix laevigata*, or *Prosopis glandulosa*) may be associated with them and may be codominant. Stands in the mapping area are mostly introduced in the recent past with the likely exception of Twentynine Palms Oasis and the Fortynine Palms Canyon area...

1415 = *Washingtonia filifera* (California fan palm oasis) Alliance

20'. *Alnus rhombifolia* is present and evenly distributed in the riparian tree layer. If *Platanus* present it is < 1% cover and not evenly distributed; *Platanus* > 1% keys to *Platanus* alliance. Occurs at extreme edge of study area near Valyermo and possibly other creeks that emerge from the Transverse Ranges...

1500 = Western Cordilleran Montane-Boreal Riparian Scrub and Forest MG034 (M034 Rocky Mountain & Great Basin Flooded & Swamp Forest)

1510 = Vancouverian riparian deciduous forest Group

1511 = *Alnus rhombifolia* (White alder groves) Forest Alliance

15'. Stands are dominated or characterized by native or non-native riparian shrubs. None of the above (couplet 15) tree species are present at significant cover or dispersion...

21. Native species of *Baccharis* spp., *Sambucus*, *Forestiera*, *Salix exigua* or *S. lasiolepis* are dominant or codominant. *Populus fremontii* and other *Salix* species may intermix at low cover and uneven distribution and tree willows or other riparian trees are < 10% cover in the stand and/or are not evenly distributed...

1420 = Southwestern North American riparian/wash scrub Group

22. A *Baccharis* species is characteristic of the overstory shrub layer (three choices below)...

Baccharis emoryi covers > 3% of a single stand and exceeds any other shrub in cover. Shrubs are like big *B. sergiloides* and tend to occur in lower elevation ditches or washes, not in granitic mountains near springs (where *B. sergiloides* is usually found). Mappable stands are rare in our area and only a few stands were detected in the field. This species occurs more commonly in the Colorado River Valley north of Blythe...

1421 = *Baccharis emoryi* (Emory's baccharis thickets) Provisional Alliance

Shrublands characterized by the dominance of *Baccharis salicifolia* usually > 50% relative cover in shrub layer. An emergent and sparse tree layer of willows or other species may also be present. Found in upper arroyos on alluvial fans emerging from San Gabriel or San Bernardino Mountains, along the Mojave River at least to Victorville, where it is adjacent to *Populus fremontii*, *Salix gooddingii*, *Tamarix*, or other riparian stands...

1422 = *Baccharis salicifolia* (Mulefat thickets) Alliance

Baccharis sergiloides dominant and characteristic; stands are usually small and associated with rocky granitic arroyos and narrow bouldery drainages adjacent to springs and seeps. Usually in desert mountains or ephemeral creeks in foothills of the Tehachapi, Liebre, or Transverse Ranges...

1423 = *Baccharis sergiloides* (Broom baccharis thickets) Alliance

22'. A *Salix*, *Forestiera*, or *Sambucus* species is dominant in the shrub layer...

23. A *Salix* is dominant (two choices below)...

Salix lasiolepis relative cover is over 50% and no other willows are dominant or subdominant. Considered a shrub even though it may be taller than 5 m, it may be accompanied by *Baccharis salicifolia* or other riparian shrubs. Small stands occur adjacent to freshwater streams and drainages in the western portion of the mapping area, usually below MMU except along Mojave River in Victorville or at the margin of the desert ecoregion as near Cajon Pass. *S. lasiolepis* > trace; if present, *S. laevigata* < *S. lasiolepis* and *P. racemosa*, *Populus spp.*, and *S. gooddingii* all trace...

1427 = *Salix lasiolepis* (Arroyo willow thickets)

Salix exigua is characteristically present as a dominant or codominant shrub with >5% cover. It forms an open to continuous canopy along riparian corridors. It often forms narrow strips along major creeks and rivers and along ditches and reservoir edges. Other willow species may be present as sub-dominants with low cover. *Salix exigua* dominates stands (> 50% relative cover in shrub layer)...

1424 = *Salix exigua* (Sandbar willow thickets) Alliance

23'. *Forestiera* or *Sambucus* are dominant (two choices below)...

Forestiera pubescens is the dominant shrub species in the canopy, usually occurs locally around permanent water or subsurface moisture. Stands occur in the western part of the mapping area adjacent to alkaline flats or on steeper slopes and along ravines in Sierra Pelona. Stands also occur in mostly montane foothill areas around isolated springs or in bottoms of narrow canyons in the foothills of the Sierra Nevada, Ord Mountains, and the El Paso Mountains. Compared to the *Salix exigua* and *Populus fremontii* alliance, *Forestiera pubescens* Intermittently Flooded Shrubland appears to prefer slightly drier conditions as upslope from flowing water. Stands are usually dense with a sparse understory...

1425 = *Forestiera pubescens* (Desert olive patches) Alliance

Sambucus nigra is dominant in the overstory (although other shorter shrubs such as *Artemisia tridentata* ssp. *tridentata* and *Eriodictyon* may be equal or somewhat higher in cover). Although considered a shrub, elderberry usually takes the form of a small tree, which forms open, well-spaced stands with a shorter shrub and herb understory. Mappable stands occur only in the margins of the study area including moist bottomlands adjacent to Mojave River near Hesperia, south to Mormon Rocks and Cajon Pass...

1426 = *Sambucus nigra* (Blue elderberry stands) Alliance

21'. Stands are dominated by non-native *Tamarix* (*chinensis*, *ramosissima*, etc.) or the tall reed *Arundo donax*.

1430 = Southwestern North American introduced riparian scrub Group (two choices below)...

24. *Arundo donax* dominates as clonal clumps in moist areas. A few small stands occur in moist areas along ditches or occasionally in lines along property boundaries or planted as windbreaks. It may not occur in study area as true semi-natural stands. The characteristic signature of *Arundo* should be sufficient to pull out small planted stands (exotic plantings as part of development polygons), but the semi natural 1431 mapping unit is reserved for areas that are not planted or are at least expanding)...

1431 = *Arundo donax* (Giant reed breaks) Semi-natural Stands

24'. Vegetation strongly dominated (>60% relative cover) by tall shrubby invasive *Tamarix* spp. (either *T. ramosissima*, *T. chinensis*, or other similar species, not including the less invasive, taller *T. aphylla*) over other native tall shrubs and/or low trees to be considered as alliance. *T. aphylla* (Athyl) trees are usually not invasive and remain in their originally planted arrangement (hortomorphic)...

1432 = *Tamarix* spp. (Tamarisk thickets) Semi-natural Stands

14'. Vegetation characterized by trees not of wetland or moist low lying areas, although they may grow in washes, arroyos, playas and other intermittently flooded situations...

25. Forest and woodlands characterized by broad-leaved evergreen trees, sometimes with dwarfed stems and small, sclerophyllous leaves (in Mediterranean climates); or various combinations of broad-leaved deciduous, broad-leaved evergreen and needle-leaved evergreen conifer trees. Canopy may range from woodland or forest structure. Winters are mild (mostly frost-free), and may be the rainiest season, springs are temperate humid, summers are hot-dry, and autumn is often dry. Characteristic tree distributions are centered in Mediterranean California climate, may be broadleaf evergreen or deciduous or may be coniferous species. Not characteristic of cooler parts of California (mountain or northern affinities)...

1.B.1. Warm Temperate Forest Formation (F018)

1100 = California Forest and Woodland Macrogroup MG009

26. Broadleaf evergreen or winter deciduous trees of California Mediterranean climate zone. Includes mostly oak trees of the genus *Quercus* in our area, but also includes small extralimital stands of *Aesculus californica* and *Juglans californica*...

1110 = Californian broadleaf forest and woodland Group

27. One or more *Quercus* species are the primary overstory canopy tree, or oaks share dominance with conifers (four choices below)...

Quercus douglasii is usually dominant to codominant locally in the tree layer. If mixed with *Quercus lobata*, *Pinus sabiniana*, *Juniperus californica*, or *Quercus chrysolepis*; must be > 60% relative cover. Generally only at the base of Liebre Mountains and in the

extreme westernmost Tehachapi Mountains near Gorman. Stands are mixed with *Juniperus californica* and *Quercus john-tuckeri* along with emergent *Pinus sabiniana* and a mixture of transmontane and cismontane shrubs and herbs. Stands mostly occur in sheltered locations such as bases of slopes or on north-facing exposures...

1111 = *Quercus douglasii* (Blue oak woodland) Alliance

Quercus lobata is dominant to codominant. May mix with *Quercus douglasii*, *Q. chrysolepis*, or *Pinus sabiniana*, but must be least 30% relative cover in canopy. Generally only at the base of the Liebre Mountains and in the extreme westernmost Tehachapi Mountains near Gorman. Stands are usually in deeper, more mesic soils than *Q. douglasii* stands and are scattered in concave to sheltered (flat) slope positions. Best stands occur in Liebre Mountains between Gorman and Lone Pine Canyon Road...

1112 = *Quercus lobata* (Valley oak woodland) Alliance

Quercus chrysolepis is dominant to codominant in the tree overstory. Only along the Transverse Ranges, on generally steep north-facing or concave exposures. If co-occurring with other oaks (*Q. lobata* or *Q. wislizeni*), *Q. chrysolepis* must be at least 30% cover. If co-occurring with *Pinus monophylla*, *Q. chrysolepis* must be > 60% relative cover. If *Pseudotsuga macrocarpa* is codominant to dominant, then key to *P. macrocarpa* alliance.

1113 = *Quercus chrysolepis* (Canyon live oak forest) Alliance

Quercus wislizeni is dominant or codominant at >30% relative cover, with other tree species in the overstory. *Q. douglasii* and *Quercus chrysolepis*, if present, occur at low cover (generally < 30% relative canopy cover). Stands are limited to the north-facing base of the Liebre and San Gabriel Mountains as far east as Cajon Pass and Silverwood Lake. Some in the Silverwood Lake stands are in low valleys or on terraces adjacent to true riparian woodlands. Many have been recently burned and are scrubby. No shrub *Q. wislizeni* ssp. *frutescens* alliance (which would key in the pre-montane chaparral group) has been identified in the study area...

1114 = *Quercus wislizeni* (Interior live oak woodland) Alliance

27'. Broad-leaf trees other than oaks are dominant or codominant (two choices below)...

Juglans californica provides an open to intermittent tree overstory canopy (in some cases it may be a

large shrub). Stands are limited in the coastal drainages south of Cajon Pass and are associated with seeps and springs. Shrubs of chaparral (*Ceanothus* sp., *Heteromeles arbutifolia*, etc.) may occur in the understory...

1115 = *Juglans californica* (California walnut groves) Alliance

Aesculus californica is dominant (>60% relative cover) as a tree or tall shrub in the overstory. If buckeye is codominant with an oak species, see the *Quercus douglasii* and *Quercus wislizeni* Alliances. Only in the extreme western portion of the study area on north facing or concave slopes of the Liebre Mountains, adjacent to stands of chaparral or *Q. wislizeni*...

1116 = *Aesculus californica* (California buckeye groves) Alliance

26'. Canopy dominated by conifers, locally, either short shrubby trees of *Juniperus californica* or by taller *Pinus sabiniana*. Stands with other xeromorphic or mesomorphic trees may occur (*Yucca brevifolia*, *Quercus lobata*, *Q. douglasii*, *Q. chrysolepis*), if so, different relative cover rules apply, see comments within key to alliances in this group.

1120 = Californian evergreen coniferous forest and woodland Group (two choices below)...

Pinus sabiniana is the dominant tree in the overstory (> 60% relative cover), and it is generally >10%. Stands identifiable as the alliance occur largely over herbaceous or mixed shrub and herb understories. Stands limited in extent usually in proximity to *Quercus john-tuckeri* (shrub), *Quercus lobata* or *Quercus douglasii* stands. Most stands occur over grassy understory in northwest Liebre Mountains....

1121 = *Pinus sabiniana* (Foothill pine woodland) Alliance

Juniperus californica is evenly distributed and the dominant species in the tree or tall shrub layer. If *Yucca brevifolia* present and evenly distributed, it is no more than 30% relative cover (or if *J. californica* > 10% absolute cover, then *Y. brevifolia* no more than 4 % absolute cover) in the tree or tall shrub layer. In other words, when *Yucca brevifolia* is present, *Juniperus* must be at least 3X its cover. Most *Juniperus* stands range from 3 to 15% absolute *J. californica* cover. When sparse (e.g., < 5%), the general rule is that *J. californica* is evenly distributed and if present with large shrubs such as *Purshia tridentata*, *Adenostoma fasciculata*, *Quercus john-tuckeri*, or *Arctostaphylos glauca*, they must have strong dominance (e.g., *J. californica* is 4% and *Quercus john-tuckeri* is < 2% goes to *Juniperus*). This is because although *J. californica* is considered a "tree" it is ecologically a large shrub...

1122 = *Juniperus californica* (California juniper woodland) Alliance

25'. Stands are pure or mixed broad-leaved deciduous or needle-leaved evergreen tree growth forms, with a seasonal green understory of herbs. Winters are cool and summers may receive some montane thunderstorms. The tall-shrub layer is variable. Stands occur in higher elevation areas at the edge of the desert in the Transverse Ranges or Sierra Nevada, or in highest desert mountains. Snow may be on the ground between <1 to 6 months of the year...

1.B.2. Cool Temperate Forest Formation (F008)

27. Stands characterized by *Pseudotsuga macrocarpa* present and evenly distributed in canopy, usually with *Quercus chrysolepis* codominant, which may be up to 3 x the cover (e.g., *Q. chrysolepis* 30%, *Pseudotsuga macrocarpa* 10%). Restricted to sheltered sites (sheltered from canopy fire and relatively steep and shady lower canyons and slopes) in Liebre and San Gabriel Mountains ...

1200 = Californian-Vancouverian Montane and Foothill Forest Macrogroup MG023 (new name: M023 Southern Vancouverian Montane & Foothill Forest)

1210 = Californian montane conifer forest Group

1211 = *Pseudotsuga macrocarpa* (Bigcone Douglas-fir) Alliance

27'. *Pinus monophylla* > 1% absolute cover and evenly distributed throughout the stand. Stand may have equal or higher cover of *Juniperus californica*, *Yucca brevifolia* and/or shrubs such as *Quercus john-tuckeri*. Locally only represented by *Pinus monophylla* alliance. Difficulty in stand identification exists at ecoregional boundaries in stands with codominance of *Juniperus californica* or with scrub oaks such as *Quercus john-tuckeri* or other tall chaparral species such as *Fremontodendron californica*, *Cercocarpus montanus*, *Arctostaphylos glandulosa*, etc. If these species are present and < 10% absolute cover, *Pinus monophylla* must be > 1% absolute cover and evenly distributed in the stand (usually it is at least 5% cover and evenly distributed). If chaparral species are > 25% absolute cover throughout the stand, *P. monophylla* must be > 10% absolute cover (e.g., scattered low cover *P. monophylla* is overruled by dense chaparral stands). If *Yucca brevifolia* is present and > 1% cover, the stand must have > 3x as much *Pinus monophylla* to be *Pinus* (*Yucca brevifolia* takes precedence when codominant). If *Juniperus californica* is present, *Juniperus* must have > 3x as much absolute cover as *Pinus monophylla* (which takes precedence when codominant)...

1300 = Intermountain Basins Pinyon-Juniper Woodland Macrogroup MG026 (new name: M026 Intermountain Singleleaf Pinyon - Western Juniper Woodland)

1310 = Western Great Basin montane conifer woodland Group

1311 = *Pinus monophylla* (Singleleaf pinyon woodland) Alliance

13'. Vegetation dominated by shrubs or herbs; trees if present, generally < 10% absolute cover, or if greater cover, then characterized by trees with xeromorphic features such as succulence, spines, or drought-deciduousness...

28. Vegetation dominated by mesomorphic grasses and shrubs, with or without scattered trees (and trees typically <10% cover), ranging from temperate coastal and inland lowland and montane grasslands and shrublands, to bogs, fens, and marshes, with a strongly seasonal climate, with at least some frost to extended cold seasons.

Occasional desert border stands have a mixing of mesomorphic and xeromorphic (class 2000, mixing with desert classes 4000 and 5000) species. The woody species of the predominant class (e.g., chaparral shrubs with small proportion of emergent xerophyll Joshua Trees) would prevail in the key...

2000 = MESOMORPHIC SHRUB AND HERB CLASS (2 Temperate & Boreal Shrubland & Grassland)

29. Vegetation defined by plant growth strategies driven by a Mediterranean climate, characterized by dry summers and mild, humid, sometimes rainy winters. Sclerophyll-leaved shrub growth forms prevail, but drought-deciduous forms may also occur. Size and coverage of the shrubs range from arborescent (2 to 5 m tall) with a closed canopy, to < 1 m and open. Mediterranean grassland and meadow are included in this formation. In California, they are separated from Mediterranean scrub at the formation level. This formation occurs primarily in the western Mojave adjacent to the Transverse Ranges and the southern Sierra Nevada and Tehachapi Mountains where enough winter moisture affords persisting stands of non-desert chaparral, coastal scrub, and grasslands...

2.B Mediterranean Scrub & Grassland Subclass

30. Vegetation characterized by shrubs averaging $\geq 10\%$ in even distribution over the stand and herbaceous species not predominant.

2.B.1. Mediterranean Scrub & Grassland Formation (F038)

31. Shrubby sclerophylls (formation typically known as chaparral) dominant in the overstory; indicator genera include *Arctostaphylos*, *Adenostoma*, *Ceanothus*, *Fremontodendron*, *Quercus* (scrub oak species), etc. Note that at the margins of the desert, chaparral intermingles with desert scrub types where unusual stand composition can occur. Where diagnostic chaparral shrub species are present and make up > 50% relative cover of woody species in a stand and are evenly distributed, the stand would key to a member of this group...

2100 = California Chaparral Macrogroup MG043

32. Sclerophyll shrublands characterized by the dominance of one or more of the following species: *Adenostoma fasciculatum*, *Arctostaphylos glauca*, *Ceanothus crassifolius*, or *Fremontodendron*. Stands may have codominance of drought-deciduous *Salvia mellifera*. This shrubland group includes chaparral typically located inland from maritime chaparral from sea level up to 2000 m (6400 feet) elevation. It ranges from inland portions of northern Baja California, Mexico, southern, central and northern California through the northern end of the Great Valley and north into Oregon. Most stands occur on well-drained soils on exposures that are in full sun much of the growing season including upper slopes, spur ridges, and convexities. This group is made up of a mixture of obligate seeders, facultative seeders, and resprouters...

2110 = Californian xeric chaparral Group

33. *Adenostoma fasciculatum* is dominant or codominant (two choices below)...

Adenostoma fasciculatum occurs as a dominant, or as a codominant with *Eriogonum fasciculatum* or other shrubs such as *Eriodictyon crassifolium* or *E. trichocalyx*. If codominant with *Arctostaphylos glauca*, then key to *A. glauca* alliance. Occurs along the margins of the Transverse Range and the Tehachapi Mountains (probably mostly out of our area in the Tehachapis). Found on convex slopes above stands of *Quercus john-tuckeri*, *Artemisia tridentata*, *Salazaria mexicana*, and *Eriogonum fasciculatum* on slopes ramping up to Cajon Pass area...

2112 = *Adenostoma fasciculatum* (Chamise) Alliance

Salvia mellifera shares dominance with *Adenostoma fasciculatum* in the shrub canopy, with *A. fasciculatum* sometimes having twice as much cover as *S. mellifera*. Only a few stands occur in coastal drainages on steep south- or west-facing. Mostly sandstone slopes south of Cajon Pass, south and east of Mormon Rocks...

2115 = *Adenostoma fasciculatum* – *Salvia mellifera* (Chamise-black sage chaparral) Alliance

33'. Species other than *Adenostoma fasciculatum* are dominant or codominant (three choices below)...

Arctostaphylos glauca is the dominant or codominant overstory shrub (locally especially with *Adenostoma fasciculatum*); conifers (*Pinus*, *Juniperus*) absent or in very low cover. Occurs only on the edges of the study area in the Tehachapi Mountains or Transverse Ranges, usually adjacent to other chaparral stands such as *Adenostoma*, *Fremontodendron*, or *Quercus john-tuckeri*. May occur adjacent to *Juniperus* or to *Pinus monophylla* stands. Most stands are small, sometimes on steep slopes mixed with scattered *Yucca brevifolia*. Largest stands in map

area are near Cajon Pass and mixed with *A. fasciculatum*...

**2111 = *Arctostaphylos glauca*
(Bigberry manzanita chaparral)
Alliance**

Ceanothus crassifolius usually occurs as a dominant or as a codominant with other chaparral shrubs in lower numbers (e.g., *Adenostoma fasciculatum*, *Heteromeles arbutifolia* and *Cercocarpus montanus*). Only occurs in the southern portion of the mapping area, southwest of Cajon Pass, which drains to the Santa Ana River and Pacific Ocean...

**2113 = *Ceanothus crassifolius*
(Hoary leaf ceanothus chaparral)
Alliance**

Stands dominated by the tall shrub *Fremontodendron californicum*, with a mixture of desert chaparral and shorter shrubs including *Purshia tridentata*, *Hesperoyucca whipplei*, *Eriodictyon trichocalyx*, *Eriogonum fasciculatum*, *Cercocarpus montanus*, *Ceanothus leucodermis*, *Ericameria linearifolia*, *Salvia dorrii*, *Artemisia tridentata* ssp. *tridentata*, and scattered emergent *Yucca brevifolia*. Occurs on coarse alluvium including edges of arroyos and washes of upper valleys or lower steep slopes of mountains. Often adjacent to *Quercus john-tuckeri*, *Adenostoma fasciculatum*, *Pinus monophylla* and *Yucca brevifolia* stands, or recent burns with *Encelia actoni* and *Eriogonum fasciculatum* stands...

**2114 = *Fremontodendron californicum*
(flannelbush scrub)
Alliance**

32'. Chaparral stands of either cooler (winters with regular frost and snow) or moister (north-facing slopes and concavities) environments than previous group. Other shrub species predominant than those listed in couplet 32...

34. Stands are either codominated or dominated by *Arctostaphylos glandulosa* or *Ceanothus leucodermis*. This group consists of sclerophyllous shrublands that are more frost-tolerant and found at higher,

cooler and generally more mesic sites than the California Xeric Chaparral Group (G257) or the California Mesic Sclerophyll Scrub Group (G261). They are particularly well-developed in central and southern California mountains between 1000 and 2000 m, and tend to be composed of both shrubs that can resprout but also have obligate seeding indicator shrubs...

2120 = Californian pre-montane chaparral Group (two choices below)

Arctostaphylos glandulosa usually occurs as a dominant or codominant in the shrub overstory. Stands are found on north-facing slopes, outcrops, and ridges on shallow soils, only on the desert-facing slopes of the San Gabriel Mountains as near Pine Creek and south of Mormon Rocks. Common only on open ridges and convex slopes surrounded by other chaparral stands (e.g., *Adenostoma fasciculatum* or *Quercus berberidifolia* south of Mormon Rocks) or near Hwy 2 and Desert-Front Road. Difficult to tell from *A. glauca* in some cases without burl inspection...

2121 = *Arctostaphylos glandulosa* (Eastwood manzanita) Alliance

Ceanothus leucodermis characterizes the shrub canopy as a dominant or codominant. No consistent canopy tree overstory (top-killed stems of short resprouts of *Quercus wislizeni* may be present, though). Stands are found primarily on north-facing slopes only in recently burned areas of the Liebre Mountains. A common post-fire regeneration type following a 2000 fire in *Quercus wislizeni* and some resprouting chaparral stands. Occurs adjacent to *Quercus lobata* stands as well...

2122 = *Ceanothus leucodermis* (Chaparral whitethorn) Alliance

34'. This shrubland group occurs in mesic site conditions such as north-facing slopes, concavities and toeslopes with well-drained soils throughout Mediterranean California, mostly inland from the coastal fog belt. It occurs most often on north-facing slopes up

to 1500 m (4550 feet) in elevation and up to 1830 m (6000 feet) in southern California. This group tends to be dominated by a variety of mixed or single-species, evergreen, sclerophyllous shrubs that resprout from lignotubers following fire.
2130= Californian mesic chaparral Group

35. Stands characterized by *Quercus berberidifolia* with or without codominance of *Adenostoma fasciculatum* (two choices below)...

Quercus berberidifolia usually occurs as a dominant or, if it codominates, it is not with *Adenostoma fasciculatum* or *Cercocarpus montanus*. It is only in the southern part of the mapping area, southwest of Cajon Pass, and probably does not occur in the Liebre Mountains. Individuals of *Q. berberidifolia* occur within stands of *Q. john-tuckeri* near Cajon Pass and stands are difficult to discriminate from *Q. john-tuckeri* alliance. When in doubt go with surrounding associated species; if desert mix then *Q. john-tuckeri*, if mostly other California chaparral species, then *Quercus berberidifolia* ...

2132 = *Quercus berberidifolia* (Scrub oak chaparral) Alliance

Quercus berberidifolia codominates with *Adenostoma fasciculatum* (*Adenostoma fasciculatum* and *Quercus berberidifolia* each greater than 15% relative cover in the shrub layer), other shrubs in the stands are significantly less cover. Only in Cajon Pass and Pine Creek area...

2133 = *Quercus berberidifolia* – *Adenostoma fasciculatum*

(Scrub oak-chamise chaparral) Alliance

35'. Chaparral shrubs other than *Quercus berberidifolia* or *Adenostoma* dominant or codominant (two choices below)...

Cercocarpus montanus > 30% relative cover with no other shrub species exceeding it in cover, or *C. montanus* and *Arctostaphylos glauca* with equal relative cover (Keeler-Wolf et al. 1998b). Both *C. montanus* and *Eriogonum fasciculatum* between 30% and 60% relative cover in the shrub canopy. Only in the northwest portion of the Liebre Mountains in the extreme west of mapping area, or on north slopes of San Gabriel Mountains or Cajon Pass area, usually associated with *Quercus john-tuckeri*, *Arctostaphylos glauca*, *Eriogonum fasciculatum* or *Adenostoma fasciculatum* alliances. May have emergent *Pinus sabiniana* or *Pinus monophylla*...

2131 = *Cercocarpus montanus* (Birchleaf mountain mahogany) Alliance

Stands dominated or codominated by *Prunus ilicifolia*. *Heteromeles arbutifolia* may be codominant. If *Quercus john-tuckeri* codominates, the stand keys to *Q. john-tuckeri* alliance. Rare in the study area; usually on either steep south- or east-facing slopes near Canon Pass...

2134 = *Prunus ilicifolia* (Holly leaf cherry chaparral) Alliance

31'. Stands dominated by drought-deciduous shrubs, though at times can have characteristic (constant but not dominant) resprouting, deep-rooted sclerophyllous shrubs. Stands

include mixed coastal shrublands from central California south into Baja, Mexico. Stands generally occur below 1500 m (5000 feet) elevation and may extend inland from the maritime zone in hotter, drier conditions. Soils vary from coarse gravels to clays but typically only support plant-available moisture with winter and spring rains. Most predominant shrubs include *Artemisia californica*, *Salvia mellifera*, *Salvia apiana*, *Salvia leucophylla*, *Encelia californica*, *Eriogonum fasciculatum*, *Eriogonum cinereum*, and *Opuntia littoralis*. On recently disturbed sites, such as after fire, *Mimulus aurantiacus*, *Lotus scoparius*, and *Lupinus albifrons* can be dominant. Note: *Eriogonum fasciculatum* alliance stands occur in different expressions throughout the study area, including coastal scrub stands on the margins of the Transverse Ranges and the Tehachapi Mountains, often adjacent to chaparral, and desert stands where often mixed with *Ephedra nevadensis*, *Yucca* spp., *Viguiera parishii*, *Simmondsia chinensis*, *Grayia spinosa*, and other desert shrubs. At this point without detailed analysis it is unwise to further segregate desert and non-desert *Eriogonum fasciculatum* alliance stands, so *E. fasciculatum* alliances will be found in this macrogroup, as well as in the Inter-Mountain Dry Shrubland and Grassland Macrogroup MG098. Dominance in membership rules in the desert macrogroup will align with desert scrub cover and thus will be only $\geq 2\%$ cover as long as no other shrub exceeds *E. fasciculatum* in cover...

2200 = California Coastal Scrub Macrogroup MG044

36. Stands usually open and/or display recent evidence of fire or other disturbance. Stands are dominated or codominated by the following species: *Gutierrezia californica*, *Lotus scoparius*, *Lupinus albifrons*, *Ericameria linearifolia*, or a species of *Eriodictyon*.

2210 = Central and south coastal California seral scrub Group (nine choices below)...

The short shrub, *Gutierrezia californica* dominates an open shrub canopy, and other shrubs may occur at low cover. The herb layer is usually well-developed, including natives such as *Poa secunda* and non-natives such as *Bromus* and *Erodium* species. Appears in western Antelope Valley associated with grasslands...

2211 = *Gutierrezia californica* (California match weed patches) Provisional Alliance

Successional shrublands occurring in chaparral or coastal sage scrub in which short-lived subshrubs or shrubs of *Lotus scoparius* dominate following disturbance, particularly fire. The shrub canopy is sometimes over a higher cover of annual or perennial herbs such as *Bromus* spp.,

Corethrogyne filaginifolia, *Nassella*, *Erodium* spp., *Avena* spp., etc. Stands have been observed only on the portion of study area that drains to the Pacific Ocean in the vicinity of Cajon Pass...

2212 = *Lotus scoparius* (Deer weed scrub) Alliance

Lupinus albifrons dominates in the shrub canopy and grows on slopes that may be disturbed, steep, and unstable. A variety of coastal sage shrubs may be present, including *Ericameria linearifolia*, *Eriogonum fasciculatum* and others...

2213 = *Lupinus albifrons* (Silver bush lupine scrub) Alliance

Ericameria linearifolia is dominant to codominant in the shrub canopy with *Isomeris arborea* and/or *Gutierrezia californica*, *Eriophyllum confertiflorum*, *Eriogonum fasciculatum* and others. The herb layer can be well-developed, and *Poa secunda* is characteristically present...

2214 = *Ericameria linearifolia* (Narrowleaf goldenbush scrub) Provisional Alliance

An *Eriodictyon* species dominates. Locally stands are dominated by *E. trichocalyx* or *E. crassifolium*. Typically, stands are *E. crassifolium* on the north side of the Liebre Mountains and *E. trichocalyx* on the north side of the San Gabriel and San Bernardino Mountains and in the Cajon Pass region...

2215 = *Eriodictyon (crassifolium, trichocalyx)* (Thick leaf and hairy yerba santa scrub) Provisional Alliance

Malacothamnus fasciculata dominates recently burned chaparral. No mappable stands have been inventoried, but small patches occur in the vicinity of the Liebre Mountains and Cajon Pass...

2216 = *Malacothamnus fasciculatus* (Bush mallow scrub) Alliance (not mapped and not inventoried in study area)

Eriogonum elongatum or *E. nudum* occur within herbaceous stands often codominated by native and non-native grasses and annual forbs. These mid-size perennial herbs are sometimes abundant colonizers of recent burns or areas released from repeated regular grazing in the foothill regions of the west Mojave Desert...

2217 = *Eriogonum elongatum, nudum* (Longstem buckwheat) Provisional

Alliance (not mapped and not inventoried in study area)

Corethrogyne filaginifolia dominates recently burned margins of chaparral or coastal sage scrub in the foothill regions of the Western Mojave Desert. Small sub-MMU stands occur in the vicinity of Cajon Pass and the northern Liebre Mountains foothills...

2218 = *Corethrogyne filaginifolia* (Common sandaster scrub) Alliance (not mapped and not inventoried in study area)

Dendromecon rigida, a short-lived shrub characteristic of recent post fire chaparral stands at the border of the study area, is dominant. Only one stand is mapped near Cajon Pass. Post fire stands are to be expected in similar areas of chaparral. Usually within a few years, stands are replaced by longer lived shrubs such as *Adenostoma fasciculatum* or *Quercus berberidifolia*...

5216 = *Dendromecon rigida* (Bush poppy scrub) Alliance

36'. Stands are characterized by the presence of *Eriogonum fasciculatum*, *Salvia mellifera*, or *Eriogonum wrightii* without significant cover (subdominant) of previous group of seral scrubs.

2220 = Central and South Coastal Californian coastal sage scrub Group (three choices below)...

Eriogonum fasciculatum \geq 2% absolute cover or >50% relative cover in the shrub canopy; other shrubs if present < half its cover, but *Hyptis emoryi* or *Salvia dorrii* may be higher (Thomas et al. 2004). Most pure stands occur along the east face of the Tehachapi and Scodie Mountains. These stands and those in the Cajon Pass area surrounded by chaparral tend to have substantially higher shrub cover and usually do not codominate with many species. Instead they are often single dominant stands. In the desert hills and mountains > 1000 m (3000 ft) elevation, *Eriogonum fasciculatum* co-occurs with many other semi-desert shrubs; if *Encelia actoni*, *Ericameria teretifolia*, *Purshia tridentata*, or *Ericameria linearifolia* are equal or higher cover, stands key to those alliances. Mixed stands with *Ephedra nevadensis*, *Ambrosia salsola*, *Ericameria cooperi*, *Grayia spinosa*, and other mid elevation shrubs only require *Eriogonum fasciculatum* to be higher cover

and more evenly distributed than any of the other shrubs...

**2221 = *Eriogonum fasciculatum*
(California buckwheat scrub) Alliance**

Eriogonum wrightii is dominant. Associated with *Eriogonum fasciculatum*, *Ericameria cooperi*, and *Prunus fasciculata* stands in the vicinity of Hesperia and Cajon Pass. May have emergent *Yucca brevifolia* or *Juniperus californica*...

2222 = *Eriogonum wrightii* (Wright's buckwheat patches)

Salvia mellifera usually > 60% or combined with a coastal scrub species > 30% relative cover in the shrub canopy (Klein and Evens 2005, Keeler-Wolf and Evens 2006). Only occurs southwest of Palmdale on Sierra Pelona and Signal Ridge adjacent to the California Aqueduct...

2223 = *Salvia mellifera* (Black sage scrub) Alliance

30'. Vegetation characterized by grasses and herbs adapted to Mediterranean climates. Shrubs, if present, not > 10% and/or not evenly distributed in the stand...

2.B.2 Mediterranean Grassland & Forb Meadow Formation

D021 California Grassland & Meadow Division

2300 = California Annual and Perennial Grassland Macrogroup MG045

Note: stands that are assumed to contain native species especially without wildflower color signature are placed into this mapping unit...

2305 = California annual and perennial grassland Mapping Unit (Native component)

37. Stands dominated or characterized by mostly annual grasses and forbs. Native herbs are characteristic and evenly distributed across the herbaceous layer, though non-native forbs and grasses may be dominant. Cover and composition vary year to year, but indicators usually present in sufficient amounts to differentiate from non-native stands. Diagnostic species include *Amsinckia* spp., *Eschscholzia* spp., *Lasthenia* spp., *Plantago erecta* and *Vulpia microstachys*...

2310 = California annual forb/grass vegetation Group

38. *Eschscholzia californica* is seasonally dominant on upland slopes or flats with well-drained sandy to loamy soils. *Amsinckia*, *Avena*, *Bromus*, *Castilleja exserta*, *Erodium cicutarium*, *Lupinus bicolor*, *Lupinus microcarpus*, *Uropappus lindleyi* and a variety of other native and non-native forbs and grasses may be present. Known from famous wildflower fields in Antelope Valley on non-alkaline soils from west of Lancaster and Palmdale to Gorman; associated with *Poa secunda*,

Achnatherum speciosum, *Gutierrezia*, *Eriogonum fasciculatum*, *Ericameria linearifolia*, and *E. nauseosa* alliance stands. Tolerates regular spring grazing and some agricultural tilling history...

2311 = *Eschscholzia (californica)* (California poppy fields) Alliance

38'. *Eschscholzia californica* is not conspicuous in the spring flowering season. Other wildflower species characteristic of Mediterranean California are present...

39. *Amsinckia menziesii*, *A. tessellata*, *A. vernicosa*, and/or *Phacelia* spp. are seasonally characteristic in the herbaceous layer with greater than or equal to 10% relative cover. Soils are often well-drained and loamy and may have high levels of bioturbation (e.g., rodent burrows) and/or high levels of (past/current) grazing. Stands may also occur well into the central Mojave Desert in good rainfall years in recently disturbed (mostly burned) desert scrub...

2312 = *Amsinckia (menziesii, tessellata)* (Fiddleneck fields) Alliance

39'. Other species besides *Amsinckia* and *Phacelia* are characteristic and/or dominant in proper phenology (three choices below)...

Native annual species *Vulpia microstachys*, *Plantago erecta* and/or *Lasthenia californica* (or *L. gracilis*) are characteristically present in stands and usually at least 10% relative in cover to other herbs. Other native species such as *Castilleja exserta*, *Crassula connata*, *Lepidium nitidum*, *Lupinus*, and *Trifolium* species are often well-represented (and sometimes codominant to dominant) as well as a variety of herbs. Soils may be clayey, wet to moist in spring and dry by summer. Stands occur in Antelope Valley west of Lancaster and Palmdale, but may occur in clearings in *Larrea tridentata* or related alliances all the way to the Granite and Sidewinder Mountains. Stands with an even distribution of *Larrea* or other desert shrubs >2% would key to the shrub type. Must have properly timed aerial imagery for detection...

2313 = *Lasthenia californica* - *Plantago erecta* - *Vulpia*

***microstachys* (California
goldfields - Dwarf plantain - Six-
weeks fescue flower fields)
Alliance**

Monolopia lanceolata is seasonally dominant or codominant on fine-textured, moderate to steep slopes, or *Coreopsis calliopsidea* is the dominant annual herb with < 2% absolute shrub cover (shrubs not evenly distributed and < 2%), forming bright golden-yellow seasonal patches on fine textured soils in the West Mojave (especially the area around Four Corners)... *Coreopsis calliopsidea* is actually likely to be a better indicator of mud hills and clay beds (6113) in the Lithomorphic class than of California annual forb/grass vegetation Group (2310)...

**2314 = *Monolopia (lanceolata)-
Coreopsis (calliopsidea)*
(*Monolopia* and Tickseed)
Alliance** (not mapped and not inventoried in study area)

Plagiobothrys nothofulvus or other similar *Plagiobothrys* are dominant and characteristic in springtimes of good rainfall. Probably does not form mappable stands in the study area; most stands with

Plagiobothrys or other ecologically similar species (e.g., *Cryptantha* or *Pectocarya* sp.) are part of other alliances such as *Eschscholzia*, *Bromus-Schismus*, *Lasthenia*, etc...

**2315 = *Plagiobothrys nothofulvus*
(Popcorn flower fields) Alliance**
(not mapped and not inventoried in study area)

37'. Stands characterized by native perennial bunch grasses such as *Nassella pulchra* or completely dominated by non-native annual grasses (e.g., *Bromus rubens*, *Schismus* spp.) or forbs (e.g., *Brassica* sp., *Salsola* sp.) with little or no native component...

40. Stands dominated or characterized by perennial native bunch grasses with usually a number of native and non native annuals present...

2320 = California perennial grassland Group

**2321 = *Nassella cernua* (Nodding
needle grass grassland)**

Provisional Alliance (not mapped and not inventoried in study area)

2322 = *Nassella pulchra* (Purple needle grass grassland) Alliance (not mapped and not inventoried in study area)

40'. Stands strongly dominated by non-natives, lacking evenly distributed diagnostic native plants (usually < 5% relative cover). Annual *Bromus*, *Schismus*, *Avena*, *Brassica* and other non-native herbaceous species are strongly dominant, with little regular cover of native herb species.

2330 = Mediterranean California naturalized annual and perennial grassland Group (four choices below)...

Locally either *Hirschfeldia incana* or *B. tournefortii* are the most prevalent dominant species (where shrubs < 2% absolute cover and not evenly distributed). Usually on sandy substrates, often near road cuts or in clearings, fallow fields, large washes, riverbeds, etc. Stands can include *Sisymbrium irio* and other species as the local dominants. *Brassica tournefortii* shows up as a straw colored signature over sandy substrates in areas such as Dunn, East of Dale Lake, etc. Can model for exotic class 2 in these settings where *Larrea tridentata* or other alliances still prevail....

2331 = *Brassica nigra* and other mustards (Upland mustards) Semi-natural Stands

Bromus rubens (*B. madritensis* ssp. *r.*) and/or *Schismus arabicus* or *S. barbatus* with the highest percent cover of non-native grasses present, generally strongly dominant; without even shrub layer or without even distribution of native desert annual herbs or grasses. In the main part of study area *B. rubens* stands are usually the result of multiple short interval fires in desert scrub such as *Larrea tridentata* – *Ambrosia dumosa*; *Schismus* tends to dominate on sandier or siltier substrates than *Bromus rubens* and tends to not take up as much area as *Bromus rubens* stands (not usually mappable)...

2332 = *Bromus rubens* - *Schismus* (*arabicus*, *barbatus*) (Red brome or Mediterranean grass grasslands) Semi-natural Stands

2333 = *Lolium perenne* (Perennial rye grass fields) Semi-natural Stands (not mapped and not inventoried in study area)

2334 = *Pennisetum setaceum* (Fountain grass swards) Semi-natural Stands (not mapped and not inventoried in study area)

29'. Vegetation not adapted to Mediterranean climate. Either typically cooler winters (more continental climate), moister summers (ameliorated locally by cool, shady slope exposures, or surface and subterranean runoff), or both. Stands are higher in the mountains or more strictly associated with cooler and moist to wet microsites...

3000 = TEMPERATE AND BOREAL SHRUBLAND AND GRASSLAND SUBCLASS (3000)

41. Stands are upland grasslands, herblands, or shrublands that are not associated with areas of moisture accumulation that could be considered marshes, wet meadows, or swales...

3100 = Western North American Temperate Grassland and Meadow Macrogroup MG048

42. Stands dominated by non native annual cool season cheat grass (*Bromus tectorum*).

3110 = Vancouverian and Rocky Mountain naturalized annual grassland Group (now considered part of G600 Great Basin & Intermountain Ruderal Dry Shrubland & Grassland Group)

No stands seen in any reconns of the area, but some mixes with *B. trinii* and *B. rubens* were seen in the Antelope Valley. *Note: B. tectorum* is very common as an understory of chaparral and *Juniperus* stands in the higher and cooler margins of the study area and would be likely to form stands if subjected to repeat short interval fires...

3111 = *Bromus tectorum* (Cheatgrass grassland) Semi-natural Stands

42'. Stands dominated by evergreen or deciduous shrubs or by native bunchgrasses...

43. Native perennial grasses such as *Poa secunda* (the only certain member of this group locally) are diagnostic and evenly distributed although may be codominant with non native annuals.

3120 = Western dry upland perennial grassland Group

Poa secunda is dominant or codominant with *Bromus rubens* on clayey soils on both flats and north-facing hillslopes along with *Allium* spp., *Claytonia parviflora*, *Erodium cicutarium*, *Dichelostemma capitata*, *Trifolium willdenovii* and other herbs. Sole member of this group in the mapping area is *Poa secunda* alliance, restricted to the area west and south of Lancaster. Occurs as stands on Tejon Ranch just across fence-line in west

Antelope Valley, and probably also large stands nearby (need field verification). If *Eschscholzia* seen in air photos, always label as *Eschscholzia*, but in other areas that have high natives and no obvious wildflower fields, might expect this alliance following fire and removal of shrub cover on north-facing slopes as far east as Ord Mountain area...

3122 = *Poa secunda* (Curly or one-sided blue grass grassland) Alliance

3121 = *Elymus multisetus* (Big squirreltail patches) Provisional Alliance (not mapped and not inventoried in study area)

3220 = Western Cordilleran montane moist graminoid meadow Group

3221 = *Muhlenbergia richardsonis* (Mat muhly meadows) Provisional Alliance (not mapped and not inventoried in study area)

43'. Shrubs dominant...

44. Winter deciduous shrubs dominant and characteristic...Sole member locally is *Ribes quercetorum* alliance

3200 = Western Cordilleran Montane Shrubland and Grassland Macrogoup MG049 (M049 Southern Rocky Mountain Montane Grassland & Shrubland)

3210= Western Cordilleran montane deciduous scrub Group (G275 Northern Rocky Mountain Montane-Foothill Mesic Deciduous Shrubland Group)

Ribes quercetorum is the dominant shrub in the canopy, often growing clonally in stands that have resprouted recently after fire on steep or concave north facing slopes. Seen only in small stands in the foothills of Liebre Mountains adjacent to *Ericameria nauseosa*, *Quercus lobata* and *Q. douglasii* stands. Most stands are <1 acre, so few can be mapped...

3211 = *Ribes quercetorum* (Oak gooseberry thickets) Alliance

44'. Stands characterized by two shrub layers, a moderately open to intermittent cover of sclerophyll shrubs and a shorter layer of drought deciduous subshrubs with at least some presence of xerophylls including *Platyopuntia*, *Cylindropuntia*, *Yucca* spp., etc and presence of many drought deciduous species such as *Ericameria* spp, *Encelia actoni*, etc., which have desert affinities. True Mediterranean California chaparral species such as *Adenostoma*, *Arctostaphylos*, and

Ceanothus (other than *C. greggii*), and others are lower in cover or absent, although stands of California chaparral Macrogroup may occur adjacent to these stands...

D061 Western North American Interior Sclerophyllous Chaparral Shrubland
3300 = Warm Interior Chaparral Macrogroup MG051
3310 = Western Mojave and Western Sonoran Desert borderland chaparral Group

45. Stands are characterized by *Quercus cornelius-mulleri* and/or *Q. john-tuckeri*. *Quercus* spp. are $\geq 2\%$ cover and not exceeded by any other tree cover; however *Juniperus californica* can be almost as high in cover. Note: These two desert scrub oak species, along with the common California chaparral species *Q. berberidifolia* (map code 2132) are extremely difficult to tell apart in the field and serve similar ecological roles. If two or more scrub oaks are both present (most likely in the Cajon Pass area for *Q. berberidifolia* and *Q. john-tuckeri*) use the sum of their absolute covers and key to the predominant species...

46. *Quercus john-tuckeri* is the dominant oak or it intermixes as a low tree with similar or higher cover than *Juniperus californica*. A variety of shrubs, such as *Arctostaphylos glauca*, *Garrya flavescens*, *Ericameria linearifolia*, *E. cooperi*, *Adenostoma fasciculatum*, *Eriogonum fasciculatum*, etc., may be present at low cover in the shrub layer...

3312 = *Quercus john-tuckeri* (Tucker oak chaparral) Alliance

46'. *Quercus cornelius-mulleri* is dominant, or codominant and evenly distributed across the stand. It occurs with a mixture of shrubs including *Eriogonum fasciculatum*, *Ericameria*

linearifolia, *E. teretifolia*, and *Yucca schidigera*. Stands are limited the vicinity of Morongo Valley and the town of Joshua Tree.

These stands are on north-facing slopes and often mix with stands of *Juniperus californica* or *Coleogyne ramosissima*. If *J. californica* and *Quercus cornelius-mulleri* are codominant, and stands contain some *Coleogyne ramosissima* and *Achnatherum speciosum*, then key to *Juniperus californica* alliance...

3314 = *Quercus cornelius-mulleri* (Muller oak chaparral) Alliance

3313 = *Quercus palmeri* (Palmer oak) Alliance (not mapped and not inventoried in study area)

45'. *Ceanothus greggii* is dominant or codominant in the stand. Although the species has been identified from the study area in the margins of the Transverse Ranges, stands have not been defined...

3311 = *Ceanothus greggii* (Cup leaf ceanothus chaparral) Alliance (not mapped and not inventoried in study area)

42'. Stands dominated by tall to short herbs and graminoids in wet to moist meadows, seasonal ponds, vernal pools, or in regularly to episodically flooded bottomlands or depressions (including saline and alkaline depressions such as playas). Due to the proximity of fresh and saline or alkaline water sources in the study area it is possible for freshwater and salt or alkaline stands to be immediately adjacent to one another. This makes mapping of wetland complexes such as near large playas, rivers, or artificial sources such as water treatment plants, sometimes problematic...

47. Stands restricted to freshwater seeps, marshes, and wet meadows. Stands are of three main types; 1) dominated by tall emergent perennial herbs such as reeds, tules, and cattails found in permanently wet soil or standing water; 2) stands dominated by smaller mostly annual herbs of ephemeral ponds and swales, drying by mid-growing season, and 3) stands dominated by mid-sized perennial wetland graminoids such as rushes (*Juncus* sp.)...

[2.C.5 Temperate & Boreal Freshwater Wet Meadow & Marsh

D031 Western North American Freshwater Wet Meadow & Marsh]

Type 1 falls within the following macrogroup and group:

3400 = Western North American Freshwater Marsh Macrogroup MG073 (now called Warm Desert Freshwater Shrubland, Meadow & Marsh MG 076)

3410 = Arid West freshwater emergent marsh Group

This type has three main mappable categories, listed below:

Phragmites australis, the tall stoloniferous wetland grass, dominates the stand. Most stands are small and occur adjacent to permanent water sources such as springs, flowing streams and rivers. Most are below mappable size for this project, with the exception of the larger stands along the Colorado River. Hybrids between the native races of the American Southwest and non-native Eurasian races are likely in some areas and make conservation prioritization difficult without detailed taxonomic study. Most stands in isolated wetlands appear native...

3411 = *Phragmites australis* (Common reed marshes) Alliance

Schoenoplectus spp., tall bulrushes, dominate the stand. Small, but often mappable stands occur in all areas of the study where ponds and sluggish permanently flowing water exist. Note: two or more alliances are treated within this mapping unit. It is not possible to map them individually. They have similar ecologies. *S. acutus* occurs in fresh or brackish water; *S. californicus* appears more regularly at edges of open water; and *Bolboschoenus maritimus* (formerly *Scirpus m.*, and not represented by a specific map code in this project), the alkali bulrush, occurs in brackish to salty or alkaline water near or on playas such as Coyote Lake...

3412 = *Schoenoplectus* spp. (Bulrush) Mapping Unit

3413 = *Schoenoplectus acutus* (Hardstem bulrush marsh) Alliance

3414 = *Schoenoplectus californicus* (California bulrush marsh) Alliance

Typha spp. dominate the stands. Most stands growing within water with slightly alkaline or saline chemistry are *T. domingensis*. Stands of *T. latifolia* have only been inventoried in fresh water at Lost Lake (Cajon Pass region)...

3415 = *Typha (angustifolia, domingensis, latifolia)* (Cattail marshes) Alliance

Type 2 is probably restricted to the Antelope Valley region and is likely present but due to inaccessibility, no certain stands inventoried. Three polygons (all privately owned and not accessible) are mapped as vernal pools or swales in this area. If accurate, these would likely be similar to those of the southern San Joaquin Valley or the inner south Coast Ranges, farther west and outside of the study area. Landscapes with pools and swales near Rosamond and north of Lancaster, which contain an even distribution of low cover *Atriplex confertifolia* over *Lasthenia* and other native herbs are keyed in the *A. confertifolia* alliance (couplet 67)...

3500 = Western North America Vernal Pool Macrogroup MG074

3510 = Californian mixed annual/perennial freshwater vernal pool/swale/plain bottomland Group

One particular alliance that may possibly occur in the area is characterized by the summer or late spring flowering *Deinandra fasciculata*. No verified occurrences of this type have been seen in the area...

3511 = *Deinandra fasciculata* (Clustered tarweed fields) alliance (not mapped and not inventoried in study area)

Type 3 is composed of mid-sized grasses and graminoids, characteristic of moist to wet meadows and marsh edges with fresh to slightly alkaline or saline water chemistry. Most stands are small, although some meet MMU requirements and have been mapped.

3600 = Western North America Wet Meadow and Low Shrub Carr Macrogroup MG075 (now called Western North American Montane & Subalpine Wet Shrubland & Wet Meadow)

3610 = Californian warm temperate marsh/seep Group (three choices below)...

Juncus arcticus, the dark brownish-green rhizomatous rush, is dominant and characteristic. Stands may include similar to lower cover of other native and non-native herbs, but *J. arcticus* is prevalent throughout. Largest mappable stands occur

in the Silverwood Lake area and around seeps and springs such as on the west side of Coyote Lake and Paradise Springs...

3611 = *Juncus arcticus* (var. *balticus*, *mexicanus*) (Baltic and Mexican rush marshes) Alliance

Leymus triticoides, the pale green creeping grass, characterizes the stands. Stands are usually too small to map and often occur adjacent to edges of permanent wetlands (when the group 3610 or Macrogroup 3600 categories are acceptable) or riparian woodland as near Victorville. May be saline, alkaline, or fresh water. Usually in slightly drier conditions than *Juncus arcticus*...

3612 = *Leymus triticoides* (Creeping rye grass turfs) Alliance

Muhlenbergia rigens, the mid size tufted perennial grass, characterizes the stands. No mappable stands have been seen within the study area, but very small stands exist in the upper Mojave River drainage upstream from Silverwood Lake...

3613 = *Muhlenbergia rigens* (Deer grass beds) Alliance (not mapped and not inventoried in study area)

47'. Stands restricted to marshes or episodic wetlands that concentrate salts or alkaline minerals. Many stands are too small to effectively be mapped...

2.C.6 Salt Marsh.

48. Stands are wet, flooded, or at least moist throughout the growing season and support dense herbaceous growth...

3700 = Warm Semi-Desert/Mediterranean Alkali-Saline Wetland Macrogroup MG083

3710 = Southwestern North American alkali marsh/seep vegetation Group

49. *Juncus* spp. (rushes) or *Schoenoplectus* or *Bolboschoenus* (bulrushes) dominate the stands (two choices below)...

Vegetation of saline or alkaline marsh vegetation dominated by densely tufted *Juncus cooperi* (similar to *J. acutus* of s coastal California). Most stands occur to the east of our study area in the Death Valley-Soda Lake Trough in the eastern Mojave Desert...

3714 = *Juncus cooperi* (Cooper's rush marsh) Alliance (not mapped and not inventoried in study area)

Schoenoplectus americanus, a vivid green triangular stemmed bulrush, forms open to dense stands in moist to flooded borders of saline or alkaline marshes in basins or near playas, such as China Lake. Large stands also mix with *Bolboschoenus maritimus* in seeps below the water treatment storage ponds north of Lancaster (edge of Edwards Air Force Base). Local stands of alkali bulrush (*Bolboschoenus maritimus* or *B. robustus*) also occur in a few areas of the mapping area, including Harper Lake Wildlife Observation area, the center of Coyote Dry Lake, and the tail below the Lancaster waste water ponds on the border of Edwards Air Force Base. Currently these are being treated as members of the bulrush mapping unit ...

3715 = *Schoenoplectus americanus* (American bulrush marsh) Alliance

49'. Stands may contain *Juncus* or *Schoenoplectus* species, but are not dominated by them (four choices below)...

Stands in moist meadows and flats often associated with alkaline water and stands of *Distichlis spicata*, *Schoenoplectus americanus*, or *Juncus arcticus*. Stands are small and uncommon, with none known to be large enough to be mapped in this study. Individual stands occur in the vicinity of Victorville near the Mojave River narrows some in actively grazed irrigated meadows...

3713 = *Anemopsis californica* (Yerba mansa meadows) Alliance
(not mapped and not inventoried in study area)

3711 = *Spartina gracilis* (Alkali cordgrass marsh) Alliance (not mapped and not inventoried in study area)

Sporobolus airoides may occur as small stands in *Atriplex confertifolia* stands near Hwy 14 and west side of Edwards Air Force Base...

3712 = *Sporobolus airoides* (Alkali sacaton grassland)

Alliance (not mapped and not inventoried in study area)

Although formally considered part of the next group (3720 = Southwestern North American salt basin and high marsh Group), *Distichlis spicata*, the short rhizomatous salt grass, is likely to be seen adjacent to episodically flooded basins, springs, playas, and salt marshes, sometimes without visible salt deposits on the surface. Mappable stands occur at Harper Lake and several other sites in the study area...

3726 = *Distichlis spicata* (Salt grass flats) Alliance

48'. Stands usually restricted to alkali or salt basins, margins of springs, or river terraces where salt deposits or alkali deposits are often present...

3720 = Southwestern North American salt basin and high marsh Group

50. Stands are dominated by succulent stemmed herbs or shrubs in the Chenopod family. Leaves are usually scale like and inconspicuous. Restricted to salty basins that may be seasonally inundated or saturated (two choices below)...

Allenrolfea occidentalis > 2% absolute cover in the shrub canopy and no other species with greater or equal cover (Keeler-Wolf et al. 1998b, Thomas et al. 2004). Saline playas and margins of salt pannes. Can occur in hummocks that are widely spaced on relatively flat playas like China Lake. Stands may also form borders around edges of playas on edges of stabilized dunes. In general, stands in the Mojave and Colorado deserts have small, low, and widely to intermittently spaced shrubs, but stands at China Lake may be denser and have *Distichlis* understory...

3721 = *Allenrolfea occidentalis* (Iodine bush scrub) Alliance

The annual pickleweed *Salicornia depressa* is the dominant herb. No other species approaching cover of *S. depressa*. In winter, skeletons of the annual plants are usually visible

to confirm identification. The only stands known are at Harper Lake, probably only 2-3 acres on average and adjacent to *Suaeda moquinii* and enhanced (trenched and watered) *Bolboschoenus* stands. Based on the imagery used for this project, stands were not mappable...

**3727 = *Salicornia depressa*
(Pickweed flats) Herbaceous
Alliance**

50'. Stands are dominated by shrubby members of the Chenopod family without succulent stems or by other species of shrubs or herbs without succulent stems. Stands are rarely inundated or if so briefly, compared to the above couplet 50...

51. Stands dominated or characterized by shrubby members of the genus *Atriplex* (three choices below)...

Stands strongly dominated (> 60% relative cover) by *A. lentiformis*. Stands are uncommon and are of two kinds. The tall, broad, bushy form of *A. lentiformis* ssp. *lentiformis* occurs on river terraces adjacent to *Populus fremontii* stands near Victorville below the Mojave River narrows mixed with *A. polycarpa*, and *A. canescens*. Rare small stands of *A. lentiformis* ssp. *torreyi* (New Mexico saltbush) occur on the beds of dry lakes. Mapping sites of *A. lentiformis* ssp. *torreyi* might be in Edwards, Lucerne, Koehn, or perhaps China lakes, locally. Charlton (in Lichvar et al. 2004) states: New Mexico saltbush is strongly associated with specific environments and occurs as pure stands in clay washes and on the playa edge where drainages empty out into the playa. New Mexico saltbush is commonly associated with

shadscale or spinescale in communities adjacent to washes in or near playas.

A. lentiformis ssp. *torreyi* is usually associated with *Suaeda moquinii* and/or *Atriplex canescens*. Based on codominance to dominance by *Suaeda moquinii*, key to *Suaeda*. Strong dominance by *A. confertifolia* keys to *Atriplex confertifolia*. For example, Charlton (at Edwards AFB) has a description of a *Suaeda–Atriplex lentiformis* ssp. *torreyi* Shadscale Association...

3722 = *Atriplex lentiformis* (Quailbush scrub) Alliance

Atriplex spinifera (spinescale saltbush) dominates or codominates with *A. polycarpa*, *Larrea tridentata*, or *Ambrosia dumosa* in the shrub canopy. When *A. confertifolia* is codominant with *A. spinifera*, key to *A. confertifolia*. The herb layer has open to intermittent cover including *Bromus rubens*, *Erodium cicutarium* and *Lasthenia*. *A. spinifera* prefers fine textured silty or clay soils that aren't strongly alkaline or salty; stands tend to have evenly spaced moderate-sized shrubs with small gray clay lenses scattered throughout, and not highly reflective whitish pannes.

Spinescale occurs with *Artemisia spinescens* and *Lepidium fremontii* near playa lakebeds, where either of these additional species may codominate. Occasionally, stands may occur in less silty and sandier soils as near Hinkley and Helendale

(west of Barstow) where one would normally predict *A. polycarpa*. Stands may be extensive or can occur in a fine matrix with *Ambrosia dumosa* (present or codominant), *Atriplex polycarpa*, and *Krascheninnikovia lanata* (e.g., in Four Corners area north to Cuddyback Lake). *Lasthenia gracilis* or *Coreopsis calliopsidea* are often the dominant annual species associated with *Atriplex spinifera*. *Atriplex spinifera* often occurs on hydrophobic soils that saturate to only a few centimeters during the rainy season. The soil surface remains moist throughout spring. These conditions favor the development of the blacktop form of cryptobiotic crust. Sensitive species observed in this community include Mojave spineflower (*Chorizanthe spinosa*) and crowned onion (*Muilla coronata*)...

**3723 = *Atriplex spinifera*
(Spinescale scrub)
Alliance**

Stands are strongly dominated by the mid-sized shrub *Atriplex parryi*. Stands are only mappable at China Lake where they occur in alkaline basins on fine textured soils just upslope from *Allenrolfea occidentalis* stands or down slope from the *Atriplex confertifolia* stands. Small stands (non mappable) have been noted within larger *A. confertifolia* stands at Coyote Lake, and with *A. spinifera* near California City...

**3729 = *Atriplex parryi*
(Parry's saltbush)
Provisional Alliance**

51'. Stands dominated by either the low subshrub *Frankenia salina*, the taller succulent leaved *Suaeda moquinii* or the low grass *Distichlis spicata* (four choices below)...

Frankenia salina is dominant or codominant (>30% relative cover) in playas, alkaline depressions and alkali sinks that have seasonally moist, poorly drained soils. *Atriplex* spp., *Cressa truxillensis* and other species may be present. Stands are part of the high marsh vegetation in southern California salt marshes; inland they occur on alkaline flats. In our study area, they are often intermixed with *Bassia*, *Salsola*, or other alkali-tolerant weeds. Stands of mappable size are unusual in the study area; they are relatively uncommon at Harper and China Lakes, and usually below 1 acre in size...

**3724 = *Frankenia salina*
(Alkali heath marsh)
Alliance**

Suaeda moquinii is $\geq 2\%$ cover and evenly distributed. No other shrub (or tree) species have greater or equal cover. Stands typically occupy strongly alkaline playas, usually with distinct salt deposits on the soil surface, but may occur in upland areas adjacent to playas (e.g., Lucerne Lake). Stands are often in fine scale drainage patterns in cracks in lake beds. They are mapped as low cover (1-5% shrub) over broad areas in such situations (as at Coyote Lake). Where wind-blown salts are deposited, *Suaeda moquinii* and *Kochia* may co-occur (Rosemond and China

Lakes), and in these cases should be mapped as the *Suaeda moquinii* alliance. There is no *Kochia* alliance defined yet. If *Suaeda moquinii* and either *Atriplex confertifolia* or *A. lentiformis* ssp. *torreyi* codominate, the alliance is *Suaeda*. If *Suaeda moquinii* and *Allenrolfea occidentalis* codominate, the alliance is the latter...

**3725 = *Suaeda moquinii*
(Bush seepweed scrub)
Alliance**

Distichlis spicata ≥ 2%, dominant or codominant with >30% relative cover in the herb layer. Soils are often deep, alkaline or saline, and poorly drained. A variety of native and non-native forbs and grasses may be present. *Distichlis spicata* is restricted to moderate to strongly alkaline and saline soils. Large (> 5 ha, 12 ac) stands do occur at Harper Lake (the south shore margin has a stand > 100 m (300 ft) wide for > 4 ha, 10 acres). Stands are associated with alkali springs, playa and panne margins...

**3726 = *Distichlis spicata*
(Salt grass flats) Alliance**

Isocoma acradenia is dominant on flat to gentle slopes near salty margins of dry lakes and playas or on episodic alkaline outwash deposits from springs and seeps. Larger stands occur on western shore of Coyote Lake and some at China and Lucerne Lakes (may not always be mappable due to small size and intermixing with *Atriplex confertifolia*, *A. parryi*, and *Suaeda moquinii* stands)...

**3728 = *Isocoma acradenia*
(Alkali goldenbush)
Alliance**

28'. Vegetation is defined by scrubs and occasional grasslands within the warm and cold desert of California. Diagnostics usually exhibit characteristics of xeromorphy and are naturally distributed in scrubs and grasslands with lower overall cover than mesomorphic (class 1000 or 2000) vegetation...

SEMI-DESERT CLASS

52. Principal indicator species are adapted to very hot dry summers and mild winters, characteristic of the majority of the lower to mid elevations of the Mojave, Sonoran and Colorado deserts. The main indicator genera include: *Ambrosia*, *Acacia*, *Chilopsis*, *Encelia*, *Larrea*, *Hyptis*, *Senna*, *Parkinsonia*, *Olneya*, *Ferocactus*, *Psoralea*, and *Krameria*. Stands with emergent *Yucca brevifolia* over higher, more uniform cover of sclerophylls (e.g., *Fremontodendron*, *Adenostoma*, or shrubby *Quercus*) key to chaparral. If stands do not meet these criteria, please go to couplet 52'.

4000 = WARM SEMI-DESERT SCRUB AND GRASSLAND SUBCLASS

53. Vegetation of the lower slopes, fans and small sheet flow areas of the warm desert parts of the state, but not of well defined washes or arroyos with clear banks and channels. Dominated by small to moderate sized shrubs (or perennial grasses) of the genera *Ambrosia*, *Encelia*, *Larrea*, *Senna*, *Parkinsonia*, *Olneya*, *Ferocactus*, *Psoralea* and *Krameria*. If *Yucca*, *Salazaria*, *Grayia*, or *Ephedra nevadensis* present, they are equal or lower cover and are overshadowed by members of the aforementioned genera. Winters may experience short frosts, but generally are not subject to persistent freezes and snow accumulation...

4100 = Mojavean–Sonoran Desert Scrub Macrogroup MG088

4110 = Lower Bajada and Fan Mojavean–Sonoran desert scrub Group

54. *Ambrosia dumosa* covers > 2% of a single stand and exceeds any other shrub in cover with the exception of *Grayia spinosa*. Stands lack significant cover of *Larrea tridentata* or *Larrea* cover is patchy and not uniformly distributed and less than 2% absolute cover. No other shrub has similar or greater cover. Stands are either on uplands with relatively fine textured soil or in active washes or terraces adjacent to medium to large washes. They also may occur on steep slopes with neutral or southerly exposures (but not too bouldery). In the West Mojave (especially the northwestern portion of the mapped area) stands often result from fire or clearing of *Larrea* in formerly mixed *Larrea tridentata* – *Ambrosia dumosa*. Many stands commonly occur mixed with *Krascheninnikovia lanata*, *Grayia spinosa*, *Ambrosia salsola*, *Ericameria cooperi*, and other species of "bathtub ring" shrubs above *Atriplex spinifera* or *A. polycarpa* and below *Larrea tridentata* – *Ambrosia dumosa* of mid- or upper fans. When stands are codominated with *Krascheninnikovia lanata*, *Ericameria cooperi*, *Tetradymia* spp., or *Eriogonum fasciculatum* (but not *Grayia spinosa*, or *Atriplex spinifera*), stands are *Ambrosia dumosa* alliance...

4111 = *Ambrosia dumosa* (White bursage scrub) Alliance

54'. *Ambrosia dumosa* may be present, either codominant with, or have less cover than *Atriplex*, *Larrea tridentata*, *Encelia*, *Pleuraphis*, or *Cylindropuntia* spp...

55. *Atriplex polycarpa* has the highest shrub cover and covers > 3% of a single stand and makes up > 50% relative shrub cover. Scattered along broader washes and on

adjacent terraces. May occur on playa edges, in washes through alkaline areas, or occasionally uplands with alkaline substrate. Stands can occur on broad flats, in washes, or on steep volcanic ravines and slopes. In this alliance, *Atriplex polycarpa* is always dominant in the shrub canopy if these shrubs are present: *Ambrosia dumosa*, *A. salsola*, *Atriplex canescens*, *Chamaesyce polycarpa*, *Cleome isomeris*, *Isocoma acradenia*, and *Larrea tridentata*. Emergent *Prosopis glandulosa* trees may be present at low cover. Stands codominant with *A. spinifera* key to *A. spinifera*...
4113 = *Atriplex polycarpa* (Allscale scrub) Alliance

55'. *Atriplex polycarpa* is not dominant; either *Encelia farinosa*, *Larrea tridentata* dominant or conspicuous and evenly spaced...

56. *Encelia farinosa* is the pronounced dominant (at least >1%), no other species have equal or higher cover. *Larrea tridentata* is largely absent. *Ambrosia dumosa* and *Fagonia laevis* have been noted as associated species. Occurs on mid to upper (most exposed) hot and dark rocky substrate on south-facing slopes, of the southern or lowest elevation parts of the study area (as far north as Trona and Spangler Hills, south side of Alvord Mountains, Paradise Range, south side of Sidewinder Mountains near Lucerne). Usually bordered by *Larrea tridentata* – *Encelia farinosa* alliance on slightly less exposed slopes (lower or less steep adjacent slopes) and giving way to *Larrea tridentata* – *Ambrosia dumosa* alliance on more neutral slopes...

4114 = *Encelia farinosa* (Brittle bush scrub) Alliance

56'. *Encelia farinosa* is not the sole dominant. If present, it is codominant with *Larrea tridentata* or subdominant to absent with other shrubs...

57. *Larrea tridentata* and *Encelia farinosa* are both present and in similar cover (broadly codominant), although sometimes *E. farinosa* < 1% cover, but if so, then *A. dumosa* also very low cover or absent. *Larrea tridentata*, *Encelia farinosa*, and sometimes *Ambrosia dumosa* codominate, sometimes along with these other conspicuous or codominant warm desert rupicolous shrubs: *Pleurocoronis pluriseta*, *Viguiera parishii*, *Trixis californica*, and *Simmondsia chinensis*. Stands sometimes have *Ambrosia dumosa*, *Hyptis emoryi*, or *Acacia greggii* as a third codominant. Usually in rocky/bouldery uplands or on well drained bajadas...

4118 = *Larrea tridentata* – *Encelia farinosa* (Creosote bush - brittle bush scrub) Alliance

57'. *Encelia farinosa* does not have significant cover with *Larrea tridentata* or other shrubs...

58. *Larrea tridentata* is either codominant with *Ambrosia dumosa*, or is the sole indicator, either mixing with other shrubs (not *Ambrosia dumosa* or *Encelia farinosa*) or as the sole dominant shrub. *Yucca* species are generally less than 1% and not evenly distributed...

59. *Larrea tridentata* is broadly codominant with *Ambrosia dumosa* and both species are evenly distributed across the stand and in combination the two species (if covers added) clearly dominate. However, mapping aggregations sometimes take into account mixes of *Larrea tridentata* without *Ambrosia dumosa* or *Ambrosia dumosa* without *Larrea tridentata* if they occur in fine scale patches within broader *Larrea tridentata* – *Ambrosia dumosa* stands. *Ambrosia dumosa* conspicuously present ($\geq 1\%$ cover) as short shrub between evenly spaced *Larrea*; may have higher cover than *Larrea tridentata*. If *Encelia farinosa* is present it is $\leq 1\%$ cover. *Yucca schidigera* if present is $< 1\%$ cover or is unevenly distributed (see *Yucca schidigera* alliance if higher cover and evenly distributed). *Atriplex polycarpa* can be codominant. Widespread on all but the hottest, rockiest, sandy, or alkaline areas of the middle and lower elevations. Do not expect *Larrea tridentata* – *Ambrosia dumosa* alliance

on old alluvial surfaces where *A. dumosa* tends not to grow. Map older alluvial fans with interfluves as *Larrea tridentata* alliance rather than *Larrea tridentata* – *Ambrosia dumosa*. Tends to occur on north-facing slopes at the lowest and hottest exposures, where south-facing would be *Larrea tridentata* – *Encelia farinosa* or *Encelia farinosa*. Only occurs on shadier north-facing exposures in the Colorado Desert, where extremely dry...

**4115 = *Larrea tridentata* – *Ambrosia dumosa*
(Creosote bush - white burr sage scrub) Alliance**

59'. *Larrea tridentata* is the dominant shrub with at least 2% cover and is evenly distributed in the stand. Associated shrubs other than *Ambrosia dumosa* or *Encelia farinosa* may be present or absent (e.g., *Krameria* spp. *Bebbia juncea*, *Ericameria teretifolia*, *Acamptopappus sphaerocephalus*, *Ephedra nevadensis* or *Opuntia acanthocarpa*)...

**4119 = *Larrea tridentata*
(Creosote bush scrub) Alliance**

58'. *Larrea tridentata* may be present but if so, at least one of the following shrubs is present in significant cover: *Tidestromia oblongifolia*, *Pleuraphis rigida* (actually a shrubby grass), *Brickellia desertorum*, *Tetracoccus hallii*, *Peucephyllum*, *Viguiera*, *Salazaria*, or *Yucca*...

60. The small stellate-pubescent subshrub, *Tidestromia oblongifolia* is unlikely to form major stands in the study area. Stands inventoried east of this area occur on sand

ramps and dune aprons in hot and very dry locations tending to have < 5% perennial cover. No mapped polygons in our area...

121 = *Tidestromia oblongifolia* (Arizona honey sweet sparse scrub) Provisional Alliance

60'. Species other than *Tidestromia* are characteristic of the stands...

61. The shrubby grass *Pleuraphis rigida*, is the dominant perennial species although may occur with scattered shrubs of *Larrea*, *Ambrosia dumosa*, *Ephedra trifurca*, or other shrub species. Stands are on sand ramps, dune aprons, stabilized dunes near playas, or wide washes adjacent to *Larrea tridentata* – *Ambrosia dumosa* stands. Stands are often mixed with lower cover of *Larrea*, *Ambrosia*, or other shrubs. Sandy stands adjacent to freeways and disturbance often have a significant non-native component of *Brassica tournefortii*, *Schismus* spp, etc.

Pleuraphis rigida is dominant and can be considered in the same layer as commonly occurring

shrubs (e.g.,
Larrea, Atriplex).
Stands not often
separable from
*Larrea – Ambrosia
dumosa* (really
*Larrea-Ambrosia
dumosa* /
Pleuraphis rigida
association), but
may occasionally
be large enough to
map in the far
eastern portion of
map area; western-
most occurrence is
west of Barstow
near Hinkley...

**4122 = *Pleuraphis
rigida* (Big galleta
shrub-steppe)**

61'. Stands of rocky
or bouldery hills and
lower mountains.
Dominated or
codominated by
either *Viguiera
parishii* or by
*Brickellia
desertorum* (two
choices below)...

Viguiera parishii
with ≥ 1 percent
cover. No other
species with greater
or equal cover,
except for *Acacia
greggii, Ambrosia
dumosa,
Simmondsia
chinensis,
Pleuraphis rigida,
Lotus rigidus, or
Encelia actoni* .
Usually on rocky
slopes in areas with
cobbles, boulders,
and/or outcrops at
low to mid
elevations or,
rarely, in washes.
No other species
with greater or
equal cover. On
northerly slopes of

the Mojave and Sonoran Desert in California. Found just above *Larrea tridentata* – *Ambrosia dumosa* alliance or in washes in east Mojave Desert. Generally at higher edge of *Larrea tridentata* – *Ambrosia dumosa* zone on bouldery, often granitic slopes in the southeast portion of study, not found elsewhere in our study area. Note: this shrub is usually non-descript (to photointerpreters) and tends to codominate with several of the above listed shrub species...

4151 = *Viguiera parishii* (Parish's goldeneye scrub)

Brickellia desertorum occurs on steep colluvial or boulder-strewn slopes of granitic or volcanic rocks; usually with relatively large, but scattered shrubs amongst rocks and boulders. Often associates with *Ericameria teretifolia*, *Atriplex polycarpa*, and may be adjacent to *Larrea tridentata* or *Larrea tridentata* – *Ambrosia dumosa*. Appears to resprout or recolonize after fires. Technically a sparsely vegetated rock outcrop type. Map when there are

not enough other definitive shrubs of other alliances and on boulders, often recently burned slopes (grass carries the fires up into the rocks)...
4123 = *Brickellia desertorum* (Desert brickellbush scrub) Alliance

53' Vegetation of well defined desert washes that shows distinct changes in plant composition and/or cover from adjacent upland vegetation stands. These washes may be broad and many-channeled or narrow and contain one or a few channels. They may occur in hills, flow across moderate sloping fans, or have a barely discernable slope and meander across lower toeslopes or basins. Diagnostic species include *Ephedra (californica or trifurca)*, *Lepidospartum squamatum*, *Ericameria paniculata*, *Ambrosia salsola*, *Prunus fasciculata*, *Brickellia incana*, *Artemisia tridentata* ssp. *parishii*, *Acacia greggii*, *Hyptis emoryi*, *Prosopis glandulosa* P. *pubescens*, *Chilopsis linearis*, *Psoralea spinosus*, *Parkinsonia florida*, and *Oleña tesota*...

4200 = Madrean Warm Semi-Desert Wash Woodland/Scrub Macrogroup MG092

Note: there are many warm and cool desert alliance indicators that may occur in washes, including *Larrea tridentata*, *Ambrosia salsola*, *Salazaria mexicana*, *Lycium cooperi*, *Atriplex polycarpa*, *A. canescens*, etc. This macrogroup is defined based on a series of indicators which are largely diagnostic of washes. Although it does exclude other alliances with broad ecological tolerances, this does not by any means exclude other alliances from occurring in washes on occasion.

62. Vegetation of washes and dominated, codominated by, or containing evenly distributed shrubs of the following species: *Ephedra (californica or trifurca)*, *Lepidospartum squamatum*, *Ericameria paniculata*, *Ambrosia salsola*, *Prunus fasciculata*, *Brickellia incana*, or *Artemisia tridentata* ssp. *parishii*...

4210 = Mojavean semi-desert wash scrub Group

63. Vegetation either dominated or codominated by *Ephedra californica* *Ephedra trifurca*, *Senna armata*, or *Brickellia incana*.. *Ephedra californica* is typically of broad, active washes of mid to upper bajadas and fans, and this species may be confused with the similar *Ephedra trifurca* of washes and sand dunes from Barstow eastward. *E. trifurca* is characteristic of low dunes and sand-sheets in the Colorado Desert but generally attains higher cover than vegetation types in the lithomorphic class that includes sparsely vegetated dunes. Due to similar ecology, both species are treated together in this *Ephedra californica* alliance...

4211 = *Ephedra californica* (California joint fir scrub) Alliance

63' Vegetation not dominated or-codominated by *Ephedra californica* or *E. trifurca*...

64. Vegetation characterized by *Lepidospartum squamatum*. Stands are concentrated along washes on the eastern base of the San Bernardino, San Gabriel and southern Sierra

Nevada Mountains. Usually in larger washes with regular flooding, the substrate texture is coarse sand to small cobbles to gravel. Stands rarely occur well out into the desert (except along the Mojave River) and are usually at the bases of mountains in arroyos or on upper fans. (Compare with *Ericameria paniculata*, *Ambrosia salsola*, and *Chilopsis linearis* alliances)...

4212 = *Lepidospartum squamatum* (Scale broom scrub) Alliance

64' Vegetation not dominated by *Lepidospartum squamatum* but instead is dominated or characterized by *Ericameria paniculata*, *Prunus fasciculata*, *Brickellia incana*, *Ambrosia salsola*, or *Artemisia tridentata* ssp. *parishii* (five choices below)...

Ambrosia salsola is strongly dominant (> 60% relative cover in dominant shrub layer). Stands of washes or disturbed uplands. Upland stands are usually associated with fire, clearing, or other disturbance in former *Larrea-Ambrosia dumosa*, *Juniperus californica*, *Yucca schidigera*, *Coleogyne ramosissima* or other upland vegetation. Most non-fire related stands of *A. salsola* are associated with washes in mid and lower elevations. *A. salsola* may mix with equal or somewhat higher amounts of *Senna armata* in washes and still be considered *A. salsola* alliance. (However, new evidence, following the conclusion of the field work for this project, suggests *Senna armata* may be a separate type in the northwest Mojave, found in the Spangler Hills and the vicinity of Red Rock Canyon State Park). Stands in washes must be > 5 acres in order to map. If they are smaller, map with adjacent best wash indicator (e.g., *Ericameria paniculata*, *Chilopsis linearis*, *Ephedra californica*, *Psoralea glandulosa*, etc)...

4216 = *Ambrosia salsola* (Cheesebush scrub) Alliance

Ericameria paniculata ≥ 2% absolute cover and ≥ 25% relative cover. Widespread throughout a broad elevation range in much of the mapping area in relatively large, recently active washes. Usually in lower energy portions of washes than *Lepidospartum squamatum* and if mixed with it, then must be > 2 x the cover of it to make this alliance. If adjacent to *Chilopsis linearis* stands and with lower extent, subsume these stands into *Chilopsis* map polygon...

4213 = *Ericameria paniculata* (Blackstem rabbitbrush) Alliance

Prunus fasciculata ≥ 2 percent cover and ≥ 25 percent of total relative cover. *Gutierrezia sarothrae* and *Lycium cooperi* may have higher cover (up to 2x). If *Prunus fasciculata* co-occurs with other tall

shrubs such as *Acacia greggii*, it must have twice the cover of other species to make the alliance definition. Usually of upper elevations (above 1000m, 3000 ft) and usually in mountains in well defined canyons or valley bottoms. Typically of washes and arroyos but may occur on wash terraces or on concave rocky slopes. Cover may be high following resprouting from fire. The following species are common associates: *Salazaria mexicana*, *Ericameria teretifolia*, *Lycium cooperi*, *Yucca schidigera*, *Rhus trilobata*, and *Purshia tridentata*. Occurs adjacent to *Eriogonum fasciculatum*, *Grayia spinosa*, or *Salazaria mexicana* stands, and also occurs adjacent to *Artemisia tridentata* stands near base of San Gabriel Mountains...

4214 = *Prunus fasciculata* (Desert almond) Alliance

Rarely strongly dominant, *Brickellia incana* usually occurs with *Ephedra californica*, and may be a part of *Ephedra californica* alliance. It occurs in sandy washes in the Central Mojave, usually at mid to lower elevations as around Coyote Lake, and south of Barstow. Some large stands strongly dominated by *B. incana* occur in areas without *E. californica* as in hills northwest of Hodge Road exit off of Interstate 15 between Victorville and Barstow.

4215 = *Brickellia incana* (Woolly brickellia) wash scrub Provisional Alliance

Artemisia tridentata ssp. *parishii* is the dominant shrub. Stands may have *Atriplex polycarpa*, *Ericameria nauseosa*, or *Atriplex confertifolia*, and may have emergent *Forestiera* or *Prosopis* at low cover or widely scattered. Small, usually linear stands associated with low gradient channels and washes in around Lancaster-Palmdale. Associated with *Prosopis glandulosa* on Edwards AFB, and with *Forestiera* and *Atriplex polycarpa* on finer soils west and north of Lancaster. This vegetation is often habitat for the rare species, *Calochortus striatus*. Mapped if ≥ 1 acre in size...

4217 = *Artemisia tridentata* spp. *parishii* (Parish's sagebrush) Provisional Alliance

62'. Wash or wetland margin vegetation of the warmer desert parts of the study area. Shrubby "trees" of mesquite (*Prosopis glandulosa* or *P. pubescens*), desert willow (*Chilopsis linearis*), smoke tree (*Psoralea spinosus*), paloverde (*Parkinsonia florida*), ironwood (*Olneya*) or tall wash or wetland shrubs such as *Pluchea sericea* and *Hyptis emoryi* characterize these stands. Usually found in areas that concentrate water such as edges of springs, river terraces, washes, etc. It is important to note that all of these taller desert woody plants are considered "trees" and thus require only a minimum of 2% absolute cover (evenly distributed) to be considered the dominant layer. This is true even if the lower shrub or herb layer is noticeably higher in cover...

4220 = Sonoran-Coloradan semi-desert wash woodland/scrub

65. Stands dominated by generally large woody plants that tend to be taller or broader than the typical *Larrea tridentata*. These include mesquite (*Prosopis glandulosa* or *P. pubescens*), desert willow (*Chilopsis linearis*), smoke tree (*Psoralea spinosus*), paloverde (*Parkinsonia florida*), or ironwood (*Olneya*) (five choices below)...

Prosopis glandulosa > 3% absolute cover as the dominant plant (including shrub and trees together), not exceeded in cover by any other species of microphyllous tall shrub or tree. Usually associated with stabilized dunes or sand sheets adjacent to playas or basins. Mapped even if very low cover, especially if evidence of recent die-off due to dewatering via ground pumping, etc. Stands along Mojave River near Daggett-Yermo are almost completely dead, but are mapped if possible...

4222 = *Prosopis glandulosa* (Mesquite bosque, mesquite thicket) Alliance

Prosopis pubescens stands are sporadic and associated with stands of the usually more abundant *P. glandulosa*. Small stands have been noted for Barstow near the Mojave River, Paradise Springs, and a few other areas. Not expected to be mapped unless field verification available. Only large stands seen have been along Colorado River in Colorado River Indian Reservation...

4223 = *Prosopis pubescens* (Screwbean mesquite bosques) Alliance

Stands characterized (1% or higher cover) by *Chilopsis linearis* no other tree-size or tall shrub species equals or exceeds *Chilopsis linearis* cover. *Chilopsis* is usually higher cover than any other large shrub/small tree, although stands may contain similar cover of *Acacia greggii* and/or *Prunus fasciculata*. Occurs in washes, intermittent channels, arroyos, or lower canyons that are intermittently flooded. Stands rarely at permanent springs or seeps and not usually associated with *Populus fremontii*, *Salix* spp., or other true riparian species; adjacent to *Ericameria paniculata*, *Ephedra californica*, *Ambrosia salsola*, and *Atriplex polycarpa* or *A. canescens* stands in washes as far west as Daggett along the Mojave River. Stands tend to occupy sandy or gravelly washes where wash energy is dissipated across a relatively wide flood path. It does not range up into mountain valleys and narrow arroyos as much as the *Acacia greggii* or *Prunus fasciculata* shrublands, and does not tend to occupy the most active wash centers such as do *Psoralea spinosus*, *Ericameria paniculata*, or *Ambrosia salsola* shrublands...

4224 = *Chilopsis linearis* (Desert willow woodland)

Psoralea spinosus is consistently distributed in low energy washes (normally at >1%, but occasionally lower). No other tall shrub or short tree species with greater cover. *Chilopsis linearis* may occur in some stands at equal cover. *Larrea tridentata* or *Ambrosia salsola* may be similar in cover. Only in eastern part of study area on lower or mid fan

wash systems out of Newberry Mountains or Twentynine Palms area. Usually associated with *Ericameria paniculata* or *Ambrosia salsola* washes, occasionally with *Ephedra californica* stands.

4225 = *Psorothamnus spinosus* (Smoke tree woodland)

Olneya tesota and *Parkinsonia florida* occur together or on their own at usually greater than 2% cover (including both shrub and tree layers). Associated species may include *Larrea tridentata* and *Ambrosia salsola*, and they may be similar in cover to *Olneya tesota* and/or *Parkinsonia florida*. Stands occur east and south of Joshua Tree National Park. They are usually tied to small to large washes and occasionally are spread out over the middle portions of large alluvial fan systems...

4227 = *Parkinsonia florida* – *Olneya tesota* (Blue palo verde-ironwood woodland) Alliance

65'. Stands with the major woody species including *Acacia greggii*, *Pluchea sericea*, or *Hyptis emoryi* visually prominent (may be as low as 2% cover) and of even distribution. These species are smaller in stature than those in couplet 65. Although the stands may include scattered individuals of those species characteristic of the previous couplet, these are in insufficient cover or dispersion to be diagnostic (three choices below)...

Pluchea sericea is present in the canopy with >2% absolute cover and no other shrub species have equal or greater cover. Occurs around springs, seeps, irrigation ditches, canyon bottoms, stream sides, and seasonally flooded washes. May include *Baccharis salicifolia*, *Atriplex*, *Ericameria nauseosa*, and others. Stands occur abundantly adjacent to the Colorado River Valley in alkaline terraces adjacent to *Prosopis glandulosa*, *Suaeda moquinii*, and occasionally *Phragmites australis* stands...

4221 = *Pluchea sericea* (Arrow weed thickets) Alliance

Acacia greggii > 2% cover. No other single tall shrub species had greater cover, but *Prunus fasciculata* or *Hyptis emoryi* may be equal or slightly greater cover than *Acacia*. Smaller shrubs such as *Ericameria paniculata* or *Ambrosia salsola* can have higher cover but no more than twice the cover of *Acacia greggii*. Occurs in washes and arroyos, as well as upland valleys and bouldery slopes. Proliferates after disturbance including flood and fire. Found in Ord Mountains and as far west as the north slope of Sidewinder Mountain into area of our study and also in eastern portion near Twentynine Palms...

4226 = *Acacia greggii* (Catclaw acacia thorn scrub)

Vegetation characterized ($\geq 2\%$) by *Hyptis emoryi*. Other shrub species are not in high cover, but may include *Acacia greggii*, *Larrea tridentata*, and *Sarcostemma hirtellum*. In rocky washes of upper bajadas and low-elevation canyons only in the eastern portion of the area north of Joshua Tree NP...

4228 = *Hyptis emoryi* (Desert lavender scrub) Alliance

52'. Stands occur in dry interior portions of the cool temperate zones of continents where precipitation is low but winters are also relatively cold. In this study, stands occur primarily in the Mojave Desert and adjacent desert mountain borderlands and not in the southerly lower elevation portions of the study area such as the Sonoran or Colorado Deserts. Locally it is difficult to differentiate some alliances in this subclass from those in the warm semi desert subclass because the Mojave Desert is transitional between cool and warm deserts and has great topographic variety. Therefore it is common to encounter stands of the warm and cool deserts (52 and 52') adjacent or even intermixed. Thus, this seemingly large dichotomy in the key is often locally less significant than it appears. Mid- and upper-elevation vegetation characteristic of the Mojave Desert ecoregion can be placed into cool semi-desert subclass based on overarching ecological conditions. This includes such iconic vegetation as *Yucca brevifolia* and *Yucca schidigera* alliances.

Stands include long persisting woody species but also commonly include species that colonize readily and rapidly when conditions are met. *Ericameria* is a particularly diagnostic genus (except *E. paniculata*). *Atriplex confertifolia* and *A. canescens* are both placed in cool desert subclass even though at the subspecific level the local expression of this vegetation may occur in warmer parts of the desert...

5000 = COOL SEMI-DESERT SCRUB AND GRASS SUBCLASS

66. Stands with the following generally short-lived and fast-colonizing shrubs common and diagnostic: *Atriplex canescens*, *A. confertifolia*, *Artemisia tridentata* (various subspecies), *Ericameria cooperi*, *E. nauseosa*, *Encelia actoni*, *Ambrosia salsola*, and *Gutierrezia microcephala*...

67. Stands are dominated by *Atriplex canescens*, *A. confertifolia* or *Sarcobatus vermiculatus*. Occur in many settings including in dry lakebeds, low dunes adjacent to them, and in rocky uplands or in sandy washes...

5100 = Cool Semi-Desert Alkali-Saline Flats Macrogroup MG093

5110 = Shadscale-saltbush cool semi-desert scrub Group

(three choices below)...

Atriplex canescens has the highest shrub cover. Stands may have emergent *Yucca brevifolia*. Prefers sandy substrates, usually stabilized dunes or sand ridges, and also sandy washes surrounded by *Larrea tridentata* – *Ambrosia dumosa*, *Yucca brevifolia* or *Yucca schidigera* alliances. May occur above 1000m (3000 ft) elevation in sandy washes in granitic mountains (such as the Sidewinder Mountains). The subspecies *linearis* prefers saltier or more alkaline sand at edges of Coyote Lake, adjacent to *Suaeda moquinii* (downslope) or *Atriplex polycarpa* (upslope). A different but ecologically similar subspecies, *A. c. var. laciniata*, occurs around the low dunes and playa margin at Palen Dry Lake. This taxon appears to be more salt-tolerant and can occur in low numbers adjacent to *Allenrolfea occidentalis* on the playa...

5111 = *Atriplex canescens* (Fourwing saltbush scrub) Alliance

Atriplex confertifolia has the highest shrub cover. Stands may occur in alkaline valleys or playas and in the upper mid-elevation Mojave Desert on rolling hills and slopes. Stands are particularly common in the northern portion of the mapping area on rhyolite, upland alkaline soils or in silty badlands. According to Charlton (in Lichvar et al. 2004), at Edwards Air Force Base, *Atriplex confertifolia* tolerates more saline and finer soils than *A. spinifera* (in areas that have high salt and clay concentrations from hydrological activity at lower elevations). When codominant with *Suaeda* on playas, stands key to

Suaeda. When mixed with *Stanleya pinnata*, *Lepidium fremontii*, and *Atriplex lentiformis* var. *parryi*, stands key to the *Atriplex confertifolia* alliance. When associated with pool and swale topography and *Lasthenia* spp. in Antelope Valley key to *A. confertifolia*. When *Atriplex spinifera* and/or *Artemisia spinescens* are codominant with *A. confertifolia* on playa edges (as at Edwards Air Force Base) stands key to *A. confertifolia* alliance...

5112 = *Atriplex confertifolia* (Shadscale scrub) Alliance

Sarcobatus vermiculatus \geq 2%. *Sarcobatus* is the relative dominant and may have *Suaeda moquinii* and *Atriplex confertifolia* or *A. canescens* associated in lesser cover. Only known in study area from small (< 1ha) stands in the alkali dunes and flats above the southeast shore of Rodgers Lake, the southwest shore of Rosamond Lake and the southeast margin of Buckhorn Lake. *Sarcobatus* seems to prefer sandy and salty soil just above the more abrupt transition to more alkaline/saline and fine textured lake bed. Locally, stands occur adjacent to *Suaeda* or *A. confertifolia* alliance stands. In one case, the edge of the stand was marked by a sandy ridge covered by an *A. canescens* – *Yucca brevifolia* stand.

Note: formally this alliance is part of the **5500 = Cool Semi-Desert Alkali-Saline Wetlands MG082**, nested within

5510 = Great Basin cool semi-desert alkali basin Group.

However, ecologically it is associated with *Atriplex confertifolia* and *A. canescens* in the few stands known from this study area...

5511 = *Sarcobatus vermiculatus* (Greasewood scrub) Alliance

67'. Stands not characterized by *Atriplex canescens*, *A. confertifolia* or *Sarcobatus*. Stands have received recent disturbance from fluvial action, fire, or clearing and are usually in rocky uplands, cobbly washes, or other areas that are not alkaline, playa-like, or particularly sandy...

5200 = Cool Semi-desert wash and disturbance scrub Macrogroup MG095

68. Dominant plants are relatively small, short-lived plants that colonize uplands following natural or unnatural disturbance such as clearing or fire. The main species are members of the genera *Ericameria*. Other common diagnostics in our area include *Encelia actoni* or *Ambrosia salsola*. *Eriogonum fasciculatum*, especially ssp. *polifolium* also occurs in such settings, especially in codominance with other species in this group, as well as *Ephedra nevadensis*, *Thamnosma montana*, and *Tetradymia* sp.

5210 = Intermontane seral shrubland Group

(four choices below)...

Encelia actoni \geq 2% cover. No other shrub species with greater or equal cover. Typically of washes or other disturbed areas (such as recently burned mid elevation desert slopes) throughout the Mojave Desert. In the West Mojave borders of the Transverse and Tehachapi ranges, stands often occur on steep south-facing slopes associated with *Hesperoyucca whipplei* or *Eriogonum fasciculatum*. The biggest stands are upland steep south- or southeast-facing slopes near Valyermo or Gramercy Avenue off of Hwy 138 south of Phelan. Small stands occur in washes and on recently burned slopes in the Stoddard Wells and Whitehorse

Mountain area and as far east as Ord Mountain. Stands may also have relatively high cover of *Achnatherum speciosum* and *Salazaria mexicana*. (Note: *Encelia actoni* was previously considered a subspecies of *Encelia virginensis*. The two share very similar ecological traits. *E. actoni* is the taxon most common in the study area)...

5211 = *Encelia (actoni, virginensis)* (Acton's encelia and Virgin River brittlebush scrub) Alliance

Ericameria nauseosa has $\geq 2\%$ absolute and $\geq 25\%$ relative cover. Found in mid and upper elevations, usually in areas with fire, flood or agricultural or grazing history. Most mappable stands are in the Antelope Valley to Hesperia. Particularly abundant on the margins of the Liebre, Tehachapi, and Sierra Nevada mountains and also abundant off alkali soils in Antelope Valley. Several subspecies are included...

5212 = *Ericameria nauseosa* (Rubber rabbitbrush scrub) Alliance

Gutierrezia microcephala or *G. sarothrae* are locally dominant on loose substrates such as gravelly washes, and on steep rocky slopes with unstable substrate. Cover of *Gutierrezia microcephala* or *G. sarothrae* is higher than any other shrub. Other shrubs may include *Grayia spinosa*, *Salvia mohavense*, *Ericameria teretifolia*, or *E. cooperi*. Most stands in our area are composed of *G. microcephala*, not *G. sarothrae*...

5214 = *Gutierrezia sarothrae* (Broom snake weed scrub) Provisional Alliance

Ericameria cooperi is evenly distributed and dominant across the stand. Stand shows evidence of recent disturbance (typically fire) and is usually adjacent to stands with larger and longer-lived shrubs more easily keyed to *Grayia spinosa*, *Ericameria teretifolia*, *Coleogyne ramosissima*, or *Larrea tridentata* – *Ambrosia dumosa*.

This is highly provisional based on strong dominance (generally $> 60\%$ relative cover) of *E. cooperi*. This is unusual and most stands with codominant *E. cooperi* can be better placed in *Ambrosia dumosa*, *Grayia spinosa*, or *Ambrosia salsola* alliances. Stands codominated by *E. nauseosa* or *E. teretifolia* key to those alliances respectively. The species occurs commonly in the West Mojave in all sub regions. It is spring flowering and shorter lived and more directly a disturbance responder than *Ericameria teretifolia*...

5215 = *Ericameria cooperi* (Cooper's goldenbush) Provisional Alliance

68'. A subspecies of *Artemisia tridentata* is dominant or codominant in the shrub canopy. *A. tridentata* (*sensu lato*) ≥ 2% absolute cover in the shrub canopy. Classification of vegetation using *Artemisia tridentata* has proceeded using different subspecies to indicate alliances due to the ecological stereotypy of many of the races. However, identifying subspecies is difficult because genotypic and phenotypic variation is common. Two different ecologically distinct races are segregated in the classification within the study area:

5300 = Western North America Tall Sage Shrubland and Steppe Macrogroup MG096

5310 = Inter-Mountain West mesic tall sagebrush shrubland and steppe Group...

69. *Artemisia tridentata* ssp. *tridentata* is dominant or codominant. No other single shrub species has greater cover except *Ericameria nauseosa*, *Eriogonum fasciculatum*, or *Eriodictyon trichocalyx*. These stands occur in coarse alluvium (granitic sands and gravels) of valleys on the north side of the San Gabriel and Sierra Pelona ranges. Stands with codominance of *Prunus fasciculata* key to *Prunus*. Stands with > 2% cover of *Juniperus californica* or *Yucca brevifolia* (regardless of height) key to *Juniperus* or *Yucca*, respectively...

5311 = *Artemisia tridentata* (Big sagebrush) Alliance

69'. *Artemisia tridentata* ssp. *parishii* is dominant or codominant. No other single shrub species with greater cover except *Ericameria nauseosa*, *Eriodictyon trichocalyx*, *Atriplex polycarpa*, *A. canescens*, *A. confertifolia*, or *A. spinifera*. Those species may be no more than 60% relative cover as long as *A. t.* ssp. *parishii* is at least 30% relative cover and evenly distributed. *A. t.* ssp. *parishii* occurs in finer textured soils more immediately adjacent to swales or intermittent channels than typical stands of *A. tridentata* ssp. *tridentata*. It is largely restricted to silty alluvial sediments of the west Mojave and adjacent Transverse Ranges and appears relatively tolerant of alkalinity. Stands occur adjacent to stands of *Atriplex spinifera*, *A. polycarpa* and *A. confertifolia*, *Prosopis*, *Larrea tridentata* – *Ambrosia dumosa* and other core Mojave Desert Alliances. They may also occur immediately adjacent to wetlands (*Juncus arcticus*, *Distichlis spicata*) and riparian (*Populus fremontii*, *Forestiera pubescens*) stands. Note: this can also be keyed in desert xero-riparian group of this key (at couplet 64')...

***Artemisia tridentata* ssp. *parishii* Provisional Alliance**

66'. Vegetation is usually characterized by scrubs of the cooler (higher elevation) desert. Most diagnostic species are long-lived. Some resprout following fire, but

some are extremely sensitive to fire (especially *Coleogyne*). Although widespread in the higher elevations of the Mojave Desert, in many areas of western and central Mojave and Sonoran desert fires and clearing have resulting in a confusion of types that intergrade between seral scrub alliance stands and more stable persistent stands. *Larrea tridentata*, *Encelia farinosa*, and for the most part *Ambrosia dumosa* are not present as major alliance indicators. Indicator species include: *Ericameria teretifolia*, *Coleogyne ramosissima*, *Grayia spinosa*, *Purshia tridentata*, *Salazaria mexicana*, *Ephedra nevadensis*, *Yucca schidigera* and *Yucca brevifolia*...

5400 = Inter-Mountain Dry Shrubland and Grassland Macrogroup MG098

70. Stands with the following species as diagnostics: *Grayia spinosa*, *Krascheninnikovia lanata*, *Ephedra nevadensis*, *E. viridis*, *Eriogonum fasciculatum*, *Lycium andersonii*, *L. cooperi*, and *Salazaria mexicana*. This vegetation merges with the upper edge of the *Larrea tridentata* – *Ambrosia dumosa* belt and is usually seen on north-facing slopes at lower elevations but will occupy basins and slopes at all elevations above 1050 m (3500 ft). May occur on slopes, but also in medium textured soils of basin margins and lower fans, especially in cool air drainages. This group includes many similar vegetation types with subtle distinction between them based on soil texture, chemistry and disturbance regime. Most types recover rapidly following fire compared to the Mojave and Great Basin Upper Bajada and Toeslope Group. Note: *Eriogonum fasciculatum* (esp. ssp. *polifolium*) is a common associate in this group especially on rocky uplands. If *Eriogonum fasciculatum* is a dominant with any of the following alliance level indicators present, key to *Eriogonum fasciculatum* (desert version), if lower in cover than any other shrub, key to appropriate other alliance based on rules in categories in the following alliances...

5410 = Intermontane deep or well-drained soil scrub Group

71. Stands are dominated or characterized by resprouting members of the genus *Ephedra* (two choices below)...

Ephedra nevadensis > 2% cover. No other species with greater cover with the exceptions of *Acamptopappus sphaerocephalus* or *Chrysothamnus viscidiflorus*. Stands in this study occur in two basic situations: (1) cooler mid- or upper slopes of mountains where codominant with *Salazaria*, *Encelia actoni* and *Ericameria cooperi*, often replacing *Grayia spinosa* or *Coleogyne ramosissima* following repeated fire; or (2) broad terraces adjacent to large washes (e.g., Little Rock Wash, Rock Creek Wash) between Pearblossom and Palmdale, often codominant with *Encelia actoni* and containing emergent *Yucca brevifolia*. Both expressions are often adjacent to *Larrea tridentata* – *Ambrosia dumosa* stands (upper elevation associations of *Larrea tridentata* – *Ambrosia dumosa* often contain *E. nevadensis*). Fire stimulates resprouting in *E. nevadensis* as does occasional fluvial disturbance. *E. nevadensis* stands are difficult to predict because of their dependence upon fire or fluvial disturbance. Rocky substrate, either cobble-alluvium or shallow broken colluvium on slopes are usually important. Stands are not found on extensive sandy or fine textured soils.

Stands often mix with other mid-elevation scrub species such as *Grayia*, *Salazaria mexicana*, *Tetradymia* spp., *Ericameria cooperi*, *Eriogonum fasciculatum*, or (near

Twentynine Palms), *Viguiera parishii* and *Simmondsia chinensis*. *Achnatherum speciosum* is common in many stands. *Coleogyne ramosissima* and *Ephedra nevadensis* often occur in similar situations and exposures, but *Coleogyne* is killed outright by fire, while *E. nevadensis* is stimulated by it. Thus, *E. nevadensis* may in some cases be a type conversion from *Coleogyne* in many burned areas of the desert mountains. If *Eriogonum fasciculatum* is codominant with *E. nevadensis*, the stand would key to *Eriogonum fasciculatum* (desert version)...

5413 = *Ephedra nevadensis* (Nevada jointfir) Alliance

Ephedra viridis ≥ 2% cover, dominant or codominant with *Ericameria teretifolia* or *Grayia spinosa*. In our area *Ephedra viridis* is dominant or codominant with *Grayia spinosa*, *Salazaria mexicana*, *Krascheninnikovia lanata*, *Ericameria cuneatus*, or *Eriogonum fasciculatum*. On steep boulder-covered slopes of mid-elevation to higher mountains in the study area from the Scodie Mountains west of the California Aqueduct to the highest points of the Ord Mountains (> 1800 m, 5900 ft). Associated with steep talus or rock outcrops except at highest elevations, when it can occur on more moderate slopes. Tends to mix with *Grayia spinosa*, *Salazaria mexicana* or with *Ericameria teretifolia* at slightly lower and warmer rocky settings. Also may mix with *Brickellia desertorum* on slopes of the Sidewinder or Granite mountains (near Apple Valley)...

5417 = *Ephedra viridis* (Mormon tea scrub) Alliance

71'. Stands not characterized by *Ephedra* (though it may be present in lesser numbers than other shrub species)...

72. Stands characterized by the dominance of a species of *Lycium* (two choices below)...

Lycium andersonii is strongly dominant without high cover of other alliance indicators such as *Salazaria*, *Grayia*, *Ephedra nevadensis*, or *Eriogonum fasciculatum*. Rarely mappable and poorly defined in the western Mojave, although a common widespread shrub of many mid- and upper-elevation scrubs of the Mojave and southern Great Basin. Stands are rare in our study area and usually small in extent. Small stands have been noted in several situations from low granitic rock outcrops, to rocky uplands above *Grayia* stands in cold air drainages and basins. This type is closely ecologically related to several other alliances, (*Grayia spinosa*, *Larrea tridentata* – *Ambrosia dumosa*, *Yucca schidigera*). (Type not described in Thomas et al. (2004), but discussed in Barbour et al. (2007). Beatley (1976) discusses its relationship to *Atriplex confertifolia* and *Grayia spinosa* in the Nevada test site...

5414 = *Lycium andersonii* (Anderson's boxthorn scrub) Alliance

Lycium cooperi is dominant and evenly distributed across the stand. Stands may include codominance of

Ambrosia salsola. If *Grayia spinosa*, *Krascheninnikovia lanata* or shrubby *Atriplex* species are codominant, stands key to those alliances, respectively. Occasional stands exist at margins of alkaline or saline basins, or on terraces above washes where soils are moderately fine textured. *L. cooperi* increases following fires relative to more weakly sprouting or non sprouting species such as *Larrea tridentata*, *Ambrosia dumosa*, and *Atriplex* spp. When codominant with *Salazaria*, *Atriplex canescens*, or *Prunus fasciculata*, as in washes and arroyos, key to those alliances, respectively...

**5418 = *Lycium cooperi* (Cooper's boxthorn scrub)
Provisional Alliance**

72'. Stands not characterized by dominance by *Lycium*...

73. Stands with *Grayia spinosa* or *Krascheninnikovia lanata* characteristic, either codominant with other shrubs, or dominant (two choices below)...

Grayia spinosa \geq 2% cover, evenly distributed, and no other species with substantially greater cover. Codominance is the rule with *G spinosa* stands; they rarely are strongly monospecifically dominant. Thus, careful assessment of shrub dispersion is important for proper identification. In many cases stands have been affected by fire, clearing, grazing, or other disturbances and seral shrubs or increasers like *Tetradymia stenolepis*, *Ericameria cooperi*, *Lycium cooperi*, *Lepidium fremontii*, *Senna armata*, or *Lycium andersonii* can have similar cover. *G. spinosa* alliance stands occupy the transition between warm desert and cool desert vegetation in much of the study area. *G. spinosa* does resprout after fire, and along with *Salazaria*, *Encelia actoni*, and *Lycium andersonii* may replace *Coleogyne ramosissima* stands as a result. There are many post-fire seral stands that have strong mixtures of multiple species, but if *G. spinosa* is evenly distributed in such stands they key to *G. spinosa*. If *Ephedra viridis* is present at \geq 2% cover and evenly distributed, please see *E. viridis* alliance.

At lower elevations *Grayia* stands usually occur on north-facing slopes in regions dominated by *Larrea tridentata* – *Ambrosia dumosa*, in lower basins and cold-air drainages on relatively well-drained medium-textured soils. Larger stands occur on moderate to gentle mid- and upper-slopes above approximately 1000 m (3000 ft) or on basin margins in a "bathtub ring" above *Atriplex polycarpa* or *A. spinifera* stands. At similar elevation to *Coleogyne ramosissima*, *Grayia* stands are on relatively less rocky and less exposed sites (mid-slope, not convex upper-slope) and often have evidence or more recent fire. Stands transition to *Ambrosia dumosa* adjacent to *Larrea tridentata* – *Ambrosia dumosa* on lower slopes and

transition to *Ericameria teretifolia* or *Ephedra nevadensis* on convex rocky slopes or *Salazaria mexicana* on concave (often burned) rocky slopes, or *Ephedra viridis* on higher elevation rocky crags or slopes. Abrupt shifts in soil texture in flats and basins give way to *Atriplex spinifera* or *A. polycarpa* on fine textured soils, or *Krascheninnikovia lanata* on calcareous soils; transitioning to *Ambrosia dumosa* or *Larrea tridentata* – *Ambrosia dumosa* on well-drained slopes above cold air pockets...

5411 = *Grayia spinosa* (Spiny hop sage scrub) Alliance

Vegetation usually of mid- to upper-elevation flats and small basins dominated strongly by the low shrub *Krascheninnikovia lanata*, without any other species in higher cover. Stands occur in small basins with silty, but not strongly alkaline soil, southeast of California City, where *Atriplex spinifera* is not dominant. These stands give way to *Grayia*, then *Ambrosia dumosa*, and then *Larrea tridentata* – *Ambrosia dumosa* in sequence up-slope. Largest stands occur in Superior Valley, in matrix adjacent to *Atriplex spinifera* on calcium rich soils (whitish caliche layer at surface), or *Atriplex confertifolia* on saltier basin soils, or *Ambrosia dumosa* on slightly higher slopes with better drained soils.

Krascheninnikovia lanata also occurs on altered calcareous soils adjacent to volcanics on northwest slopes and eastern slopes of the El Paso Mountains on shallow caliche or dolomite (?) where stands tend to codominate with *Grayia* and *Tetradymia fasciculatum*. Stands were assumed rare throughout the California deserts, but some extensive stands have been mapped in this study. These include the largest ones now known from California...

5412 = *Krascheninnikovia lanata* (Winterfat scrubland) Alliance

73'. Stands neither with *Grayia* or *Krascheninnikovia lanata* as conspicuous or dominant...

74. Stands with *Ericameria teretifolia* \geq 2% cover. No other species with greater cover, but can share dominance with *Eriogonum fasciculatum*, *Gutierrezia sarothrae*, or *Opuntia chlorotica*. Found in disturbed uplands in the mid-elevation Mojave or Sonoran Desert, but also occurs in longer-persistent stands on shallow granitic pediments and rock outcrops. *Ericameria teretifolia* is the dominant or codominant shrub and may occupy shallow rocky post-fire stands associated with *Juniperus californica* or other upland alliances. In our area it is usually found as low cover

shrubland in granitic or other rocky uplands on south- or north-facing steep, bouldery slopes, and is more warm-tolerant than *Ephedra viridis* and thus usually at lower elevations. When codominant with *Grayia*, *Ephedra viridis*, *Coleogyne*, or *Salazaria*, key to those alliances...

5416 = *Ericameria teretifolia* (Needleleaf rabbitbrush scrub) Alliance

74'. Stands with *E. teretifolia* absent or in low relative cover. Instead, stands are dominated or codominated by *Eriogonum fasciculatum* or *Salazaria mexicana* (two choices below)...

Eriogonum fasciculatum \geq 2% absolute cover or $>$ 50% relative cover in the shrub canopy; other shrubs if present $<$ half its cover, but *Hyptis emoryi* or *Salvia dorrii* may be higher (Thomas et al. 2004) and *Ephedra nevadensis* can be broadly codominant with *Eriogonum fasciculatum*. Most pure stands occur along the east face of the Tehachapi and Scodie Mountains. These stands and those in the Cajon Pass area surrounded by chaparral tend to have substantially higher shrub cover and usually do not codominate with many species, instead they are often single dominant stands. In the desert hills and mountains $>$ 1000 m (3000 ft) elevation, *Eriogonum fasciculatum* co-occurs with many other semi-desert shrubs; if *Encelia actoni*, *Ericameria teretifolia*, *Purshia tridentata* or *Ericameria linearifolia* are equal or higher cover, then go with those alliances. Mixed stands with *Ephedra nevadensis*, *Ambrosia salsola*, *Ericameria cooperi*, *Grayia spinosa* and other mid-elevation shrubs only require *Eriogonum fasciculatum* to be higher cover and more evenly distributed than any of the other shrubs...

2221 = *Eriogonum fasciculatum* (California buckwheat scrub) Alliance

Salazaria mexicana $>$ 2% cover. Other shrubs, if present, are each less than half of the cover with the exceptions of *Hyptis emoryi*, *Senna armata* or *Salvia dorrii*, which may have higher or equal cover. *Salazaria* stands are mostly restricted to sandy or gravelly washes in terrain where fire has been minimal, but may occur on burns or in other disturbed uplands throughout steep and rocky uplands. Substrate for all expressions of this alliance is frequently granitic or crystalline non-calcareous

metamorphic (gneiss, schist, phyllite). In washes, often commonly occurs with *Ambrosia salsola*, *Bebbia juncea*, *Eriogonum fasciculatum* or *Senna armata*. On rocky slopes tends to occupy bases of larger outcrops or narrow concave defiles, or ravines where water is channeled during run-off. Depending upon the site topography, many upland *Salazaria* polygons may actually contain a fine scale matrix of several vegetation alliances including *Encelia actoni*, *Ephedra nevadensis*, *Eriogonum fasciculatum*, *Ambrosia salsola* or *Ericameria teretifolia*. The overall expression of burned rocky uplands with a number of concavities or concentrations of vegetation at the base of rock outcrops tends to emphasize the balance of overall cover toward *Salazaria*, even though other small stands may be present...

5415 = *Salazaria mexicana* (Bladder sage scrub)

70'. Stands with other shrub or herbaceous species as diagnostics...

75. Shrubs form the dominant layer (at least 2% and evenly distributed). If present, *Grayia spinosa*, *Krascheninnikovia lanata*, *Ephedra nevadensis*, *E. viridis*, *Eriogonum fasciculatum*, *Lycium andersonii*, *L. cooperi*, *Salazaria mexicana* usually less conspicuous or subordinate to species in the genera *Coleogyne*, *Purshia*, *Menodora*, *Cercocarpus*, or *Yucca*...

5420 = Mojave and Great Basin upper bajada and toeslope Group

76. *Coleogyne ramosissima* is the dominant or codominant shrub, typically with no species of *Yucca*, *Juniperus*, or taller shrub greater than 33% of the total relative cover of *Coleogyne*, though other smaller shrubs such as *Ephedra nevadensis* may equal it in cover. If present with *Yucca schidigera*, that species needs only to be >1% absolute cover and evenly distributed to shift into a *Yucca schidigera* alliance if *Coleogyne* is as much as 3 x the cover of *Yucca schidigera*. If *Coleogyne* >3 x *Yucca schidigera* then it is a *Coleogyne* stand. *Coleogyne* typically dominates stands, but may be exceeded by species of disturbance (*Ambrosia salsola*, *Salazaria mexicana*, *Ericameria* spp. or *Eriogonum fasciculatum*). A widespread type on shallow rocky soils on upper bajadas, pediments and hill slopes, generally upslope from *Larrea tridentata* – *Ambrosia dumosa* on shallower soil of old alluvium or shallow rocky pediments. Does not prefer steep colluvial deposits with larger rocks and boulders. It is extremely susceptible to even low-intensity fire and many thousands of acres of *Coleogyne ramosissima* are now converted to *Grayia*, *Salazaria*, *Ericameria*, and *Ambrosia* types throughout the mapping area. Particularly devastated

are the Stoddard Wells, Fairview Valley, and Ord Mountain areas...

5421 = *Coleogyne ramosissima* (Black brush scrub) Alliance

76'. *Coleogyne* not important. Instead *Yucca*, *Purshia*, *Cercocarpus* or *Menodora* conspicuous and or dominant...

77. A *Yucca* species conspicuous and evenly distributed (two choices below)...

Yucca brevifolia present and evenly distributed throughout stand, though usually only between 1 and 5% cover (highest cover may be 10% in clonal stands in the western part of the mapping area), often with substantially higher cover of shorter shrubs or perennial grasses beneath the well-spaced emergent trees. *Y. brevifolia* must be evenly distributed, not scattered and clumped, and must be 1% or greater absolute cover to map alliance stands. It is usually difficult to discern juvenile *Yucca brevifolia* < 3 m tall, so these are not always accounted for in mapping. If *Juniperus californica* is present, *Yucca brevifolia* must be > 2 x the cover of *Juniperus*. *Pinus monophylla* must be lower than 1% absolute cover and not evenly distributed. If sclerophyll shrubs such as *Fremontodendron* or *Quercus john-tuckeri* are present, these are less than 10% absolute cover...

5423 = *Yucca brevifolia* (Joshua tree woodland) Alliance

Yucca schidigera conspicuous, evenly distributed and generally $\geq 2\%$ absolute cover. However, since sub-meter imagery has been used and signatures are generally recognizable at low covers, *Y. schidigera* has been pulled out as low as closer to 1% absolute cover as long as evenly distributed. At lower elevations may have *Larrea tridentata*, *Ambrosia dumosa*, and other shrubs at equal or even higher cover. *Yucca brevifolia* is often scattered in tree layer (<1%). At upper elevations *Juniperus californica* may be present (< 2% cover). If *Juniperus* is $\geq 2x$ the cover of *Yucca schidigera*, then it is *Juniperus californica* Woodland Alliance. If *Coleogyne ramosissima* is conspicuous, *Yucca schidigera* $\geq 2\%$ cover. On pediments and hillslopes near the upper-elevation range of the alliance, and common on bajadas and moderate to gentle hillslopes at mid-elevations. Predominantly in southeast portion of the study area including Ord Mountains, Stoddard Wells, and Fairview Valley especially...

5424 = *Yucca schidigera* (Mojave yucca scrub)

77'. *Yucca* not important and evenly distributed. Instead, either *Cercocarpus*, *Purshia*, or *Menodora*

are conspicuous, dominant, or codominant (three choices below)...

Purshia tridentata \geq 2% absolute cover and comprises higher relative cover than any other single shrub (but see below). If *Artemisia tridentata* or *Ephedra viridis* are present they have less than 1% cover. Stands are locally represented adjacent to San Bernardino, San Gabriel and southern Sierra Nevada Mountains. Confusion between this alliance and *Eriogonum fasciculatum*, *Encelia actoni*, and *Ericameria linearifolia* alliance stands occurs on the north slopes of the San Bernardino Mountains south of Lucerne Valley; all stands with evenly distributed codominant to dominant *P. tridentata* are keyed to *P. tridentata*. Stands in the current study area are often the result of burned *Juniperus californica* stands and resprouting *Purshia*. Exceptions are steep rocky slopes near Cushionberry Grade and steep incised faces of old alluvial surfaces at the base of the San Gabriel Mountains near Valyermo, where stands are codominant with *Eriogonum fasciculatum*, *Encelia actoni*, and *Hesperoyucca whipplei*...

5422 = *Purshia tridentata* (Bitter brush scrub) Alliance

Menodora spinescens \geq 2% cover, no other single species with greater cover although many other species may be present. Represented by a few localized stands in well-defined, shallow rocky soils characteristically just above *Larrea tridentata* – *Ambrosia dumosa*. Locally restricted to pale limestone (or dolomite?) outcrops associated with pediments north of Ord Mountain. Upon further reconnaissance, may be restricted to a single stand in study area...

5425 = *Menodora spinescens* (Greenfire scrub) Alliance

Vegetation characterized by the relative dominance of the shrubby tree *Cercocarpus ledifolius*, which must be at least 2% cover and evenly distributed. No other large shrub or tall tree equals or exceeds cover of *C. ledifolius*. Stands occur in dry, rocky, and usually very well-drained exposures in the highest portions of the Inyo, Panamint, and other tall ranges of the northern Mojave Desert, but only two places in the study area: adjacent to *Coleogyne* on limestone on the north slope of the San Bernardino Mountains. These low elevation stands (1400 m, 4600 ft) are likely to be the only stands in the study area...

5441 = *Cercocarpus ledifolius* (Curl leaf mountain mahogany scrub) Alliance (formally within the 5440 = Intermountain shallow/calcareous soil scrub Group)

75'. Shrubs not evenly distributed and perennial grasses comprise the dominant layer (three choices below)...

5430 = Southern Great Basin semi-desert grassland Group

Achnatherum speciosum is dominant, with no shrubs comprising more than 10% relative cover and none evenly distributed across the stand. Stands are the result of fire eliminating desert shrub cover from stands formerly *Coleogyne ramosissima*, *Larrea tridentata* – *Ambrosia dumosa*, *Atriplex canescens*, etc. The largest stands noted occur in the western Antelope Valley and Superior Valley. Persistence is unknown, but probably gives way to shrub dominance in < 50 years without fire or other disturbance...

5431 = *Achnatherum speciosum* (Desert needlegrass grassland) Alliance

Pleuraphis jamesii comprises > 2% cover with no other perennial grasses or shrub species in greater cover. Stands are restricted to below minimum map unit size except perhaps on parts of the upper slopes of Ord Mountain, Stoddard Mountain, Sidewinder Mountain and other peaks ≥ 1500 m (4500 ft) in the central or northern portions of our study area that receive some reliable summer precipitation. They occur on rocky gentle to moderately steep slopes, in some cases adjacent to unstable scree or talus. The small < 1 acre stands are often associated with *Grayia*, *Ephedra viridis*, *Gutierrezia microcephala* and *Opuntia eriantha*. Fire on the top of Ord Mountain has probably enhanced the species extent, but most individuals are associated with sufficient shrub cover to be considered as an understory rather than the dominant species in a single herb layer...

5432 = *Pleuraphis jamesii* (James' galleta shrub-steppe) Alliance

Achnatherum hymenoides is dominant; cover is usually < 10% and stands transition to *Larrea tridentata* or *Larrea tridentata* – *Ambrosia dumosa* when shrubs become evenly distributed at > 2% cover. Small stands are widely scattered throughout the mapping area on dune aprons and other sandy soils. The most extensive stands seen were in the area east of Lucerne Lake north of Old Woman Springs, but other smaller stands exist well westward. Stands that have < 2% *Achnatherum hymenoides* and > 2% *Abronia* or *Dicoria* would key to the *Dicoria canescens* – *Abronia villosa* Alliance (6121)...

5433 = *Achnatherum hymenoides* (Indian rice grass grassland) Alliance

Appendix D

Land Use Mapping Criteria

Criteria For Mapping Land Use In the 2012 DRECP Mapping Area

General Rules

Although land use is not considered a “special” vegetation type with a minimum mapping unit (MMU) of 1 acre, the MMU is lowered from the standard 10 acres for upland vegetation to an MMU of ~2.5 acres for all land use types (codes 9000’s through 9300 and 9805). Water types 9800, 9801, 9803 and 9804 have a 1 acre MMU and are assigned a land use code in the Land Use layer (land use code = 9800 for these water types).

Note that the study area includes small portions or fringes of the Tehachapi Mountains, Sierra Pelona, San Gabriel Mountains, and San Bernardino Mountains, which technically are not a part of the desert ecoregion, but do transition into it. For the sake of future mapping continuity these fringe “Foothills” areas are mapped at 2.5 acre MMU for vegetation, using Statewide vegetation mapping criteria. The land use in these areas, however, uses the Desert criteria.

VEGETATION CODES (Vegetation Layer ONLY):

9000 = MISCELLANEOUS CLASSES

9200 = Agriculture (within the current 5-year cycle)

A two-tiered coding system (Vegetation type and Land Use) is used. A given polygon can be coded as both a natural vegetation type and a land use type indicating agriculture if the situation warrants a dual code. 2005 through 2010 image sources are used to interpret and code an area as Agriculture or as a natural vegetation type.

Agriculture in the desert may be difficult to map, especially in areas that have historically been farmed. When mapping an area of agriculture, the question becomes, “when should an area no longer be considered agriculture?” Old plow and irrigation marks on land that has not been cultivated in the past 10 to 20 years or longer may still be visible on the current imagery sources, giving the impression that the area may be agriculture. Based on the successional trends of vegetation in a given area, if vegetated with any identifiable vegetation type, other than herbaceous, any area fallow for < 5 years is considered a candidate for vegetation/land use dual coding.

The Agriculture (Ag) polygon boundary is drawn to the largest actively farmed area seen on any of the imagery in the 2005-2010 time span. Using the multiple image sets as a guide to code the agriculture areas may result in polygons coded as Ag that are not photomorphic to the photo signature on the 2010 imagery. For example, if an area was last actively farmed in 2005, but since then was not active, then the Ag polygon is coded based on the 2005 image. However, the Ag delineation is still drawn to the 2010 image base (e.g. following roads, fence lines, etc. that usually appear on both sets of imagery).

In order to be consistent, the following criteria have been set:

- a. Land that has been actively farmed within ~5 years is considered agricultural.
 - If the area in question shows signs of **active** agricultural use (e.g., crop irrigation patterns, or other signs of actively managed crops) on any of the image sources, then the area is called Agriculture in the Land Use layer (use either land use code 2100 or 2200).
 - If an area is shown on the 2010 imagery as fallow or weedy but earlier sources (2005-2010) show active agriculture (crops, plowed dirt, etc.), then the area is still mapped as Agriculture (either 9210 or 9220) in the Vegetation layer.
 - Grassy or vacant areas between and at the outer corners of crop circles are left in Ag regardless of the 10 acre mmu. The crop circle areas are mapped as a box, using fence lines or roads as the boundary where possible.

Appendix D – Land Use Mapping Criteria

- b. If the area appears to be **inactive** agriculture (based on the 2005-2010 imagery) and remains unchanged from image source to image source, then the mapper made the assumption the area is no longer being used for agricultural purposes even though old plow and/or irrigation marks are still visible. Usually the imagery will show a mottled grass or herbaceous signature. Shrubs may be present in varying amounts and distribution.

Sometimes the signature may appear as a smooth, tan grassy signature, which may be due to previous *Hordeum* plantings or other single grass/grain crop (prior to 2005) that has since been inactive. In this area, the *Hordeum* is currently growing from old seed and not actively maintained. These areas are mapped as grass unless the field data notes otherwise, such as metal irrigation structures present.

- c. An **exception** to the 5 year active Ag rule is if the area in question appears to have enough naturally colonizing shrubs to be coded as a vegetated shrub type (on the 2010 imagery), then the Veg layer is coded the appropriate shrub type (e.g., 5212 = *Ericameria nauseosa*) and the Land Use layer is coded with the appropriate Land Use Agriculture code (e.g. 2100 = Non-woody row & field crops).
- d. In rural or agricultural regions, if there is a structure within a mapped unit that meets the 2.5 acre MMU, the polygon is urban (9300 or 9310) even if the structure is Ag related. This includes structures that are associated with dairies, horse ranches or other Ag related uses. If it doesn't meet the 2.5 acre MMU, it is ignored.
- e. Agricultural areas within an Urban Window are mapped if they are >10 acres.

One reason the vegetation code-dependent attribute “anthropogenic disturbance” was created is to account for visually obvious, but > 5 years pre-imagery agricultural disturbance (see definition in “mapping attributes with examples” document).

NOTE: Polygons should not be coded with the generic 9200 code but are assigned a more specific 9210 or 9220 code as described below:

9210 = Woody Agriculture (orchards, vineyards)

Corresponding Land Use layer code is 2200.

Defined as showing orchards, vineyards, jojoba farms, etc. within the past 5 year set of imagery (usually from NAIP early summer 2005-early summer 2010). Abandoned orchards remain as 9210 until the trees have actually been removed. If an orchard/vineyard is completely removed based on the 2010 imagery, then the area is mapped as its current (2010) vegetation and/or land use type.

9220 = Non-woody Row and Field Agriculture

Corresponding Land Use layer code is 2100.

Defined as showing as active agriculture at least once within the past 5 year set of imagery (usually from NAIP early summer 2005-early summer 2010).

9300 = Built-up & Urban Disturbance

Corresponding Land Use layer code is 1000.

Settlements – Suburban Areas

Areas of urban development that are below the MMU for Urban Window: MMU for code 9300 is 2.5 acres; MMU for Urban Window (9310) is 1 square mile.

- a. Settlements or other developed areas (9300) are not mapped next to an Urban Window (9310). The 9300 is incorporated into the 9310.

- b. Vacant areas within a settlement that are disturbed, scraped, or grassy are cut off from the natural vegetation around the settlement and included with the Urban. There is usually a road at the urban break that is used as a polygon boundary.
- c. Vacant areas within a settlement or rural residential area that has “natural” vegetation are mapped using the following criteria:
 - A vegetation polygons is pulled out if it is at least 10 acres of contiguous vegetation not split or disrupted by roads or other man-made features.
 - If the vegetated area meets the 10 acre unsplit (by roads) MMU then other vegetated areas adjacent to the “main” unit but split by roads are added to the vegetation polygon.
 - “Snaky” vegetation polygons are not created just to meet the 10 acre MMU.
 - This rule applies to areas that are “more built-up” (settlements) and does not apply to “more natural” areas that are just split by multiple roads.
- d. Non built-up “holes” within the settlement that are scraped or otherwise disturbed are left as part of the Urban built up (9300) polygon.
- e. Natural vegetation that comes into the settlement from the outside is continued into the Urban area as a natural vegetation type if the natural area meets the 10 acre MMU. This linework can either be straight or more natural looking. When the natural vegetation is riparian, the MMU is lowered to 1 acre to maintain continuity.
- f. If all the vegetation within the urban settlement is >10% natural then it is coded as a vegetation type in the Vegetation layer, with the Land Use code of Urban (1000) in the Land Use layer.

Scraped lots and any urban built up areas <1ac adjacent to urban are included in the urban polygon. Context is used for this guideline - scraped areas may not always get mapped into the land use, especially if the scrapings are linear along a roadway or fence.

- g. If an area is under construction (including buildings or cleared land with an urban development footprint), it is coded as 9300. This includes under construction areas that are adjacent to existing land uses such as residential developments as well as areas that are isolated.
- h. Cleared areas on horse-related property that may be dirt or grass (looks green in most cases) are coded as 9300 (no dual coding for veg and land use).
- i. Flood Control Basins – are included in the 9300 polygon but are mapped separately as a 9805 if >10 acres.

Military Target Areas

Cleared areas that are on the DRG as “target” are coded as 9300 in the Veg. These have been mapped on Edwards Air Force Base. Additionally, areas that are maintained as cleared areas are also coded as 9300.

Isolated Built-up Areas (9300) vs. Natural Vegetation PI Code

Isolated built-up areas not associated with an urban window or settlement/ suburban area are typically more rural in character, and can range from one isolated homestead to a group of houses on large lots mixed with vacant lots, small agricultural plots, and pods of natural vegetation.

- a. If Natural vegetation is <10% the entire area is coded as 9300. This polygon would get a Land Use layer value of 1000.

- b. If Natural vegetation is >10% the entire area is coded as a natural vegetation type in the Veg code and a Land Use layer value of 1000.

Miscellaneous Considerations

- a. Photointerpreters were instructed to try to keep the polygon delineation tight to the land use and associated land use disturbance by pulling out as much natural vegetation without land use as possible.
- b. If there is a large area (at least 2.5 acres) of disturbance (scraped land) with very little to no development, it is mapped as a vegetation code of 9320 with Anthropogenically Altered Disturbance = 3.
- c. Major four-lane divided highways and freeways, such as State Highways 14 and 58, and Interstates 15 and 40 are pulled out to the fenced right-of-way as Urban. Vegetation within it is not pulled out separately.
 - i. If the right-of-way (ROW) falls below the 2.5 acre mmu width (1/2 the width of a 2.5 acre box), that portion of the highway is not mapped unless it is a very short segment, thus keeping the roadway connectivity intact.
 - ii. Typically the type of vegetation within the ROW is a disturbance type of vegetation and is different from the natural type of vegetation outside of the ROW. In this case, the ROW boundary is used and the disturbance vegetation in the road polygon is included.
 - iii. The exception to this is when the ROW extends beyond 90' past the pavement edge, when the disturbance corridor is reevaluated and the ROW boundary may not be used as the road boundary.
 - iv. Lines are not drawn between the same type of vegetation based on a ROW fence line. The vegetation is kept together in one poly and the road in a different poly where possible.
- d. Active quarries are mapped as 9300 (the Land Use layer attribute will be coded as Urban = 1000). Vegetated areas within an active mine operation are included.
- e. Underground mines are mapped as natural vegetation with no land use coding. The mapper can code for Anthropogenically Altered Disturbance level 1, 2 or 3 depending on surficial disturbance visible on images.
- f. Inactive quarries, usually where vegetation has been reestablished, are dual coded with a natural vegetation type, an Anthropogenically Altered Disturbance code of 1, 2, or 3, and a Land Use layer value of 1000.

9310 = Urban Window

- a. Fully developed contiguous area of built up and disturbed lands greater than 1 square mile in size are considered the Urban Window.
- b. Urban/disturbed polygon (9300) are not mapped next to an Urban Window, but are incorporated into the 9310.
- c. If an urban area is smaller than 1 square mile, it is mapped as a 9300.
- d. If an area is within the Urban Window and under construction (including buildings or cleared land with an urban development footprint), it is coded as part of the 9310. This includes under construction areas that are adjacent to existing land uses such as residential developments and areas that are isolated.

Appendix D – Land Use Mapping Criteria

- e. Agricultural areas are only mapped within an Urban Window if they are >10 acres. However, agricultural areas along the edge of an Urban Window adhere to the 2.5 acre MMU rule.
- f. Vacant areas within an Urban Window that are “natural” vegetation are mapped as vegetation polygons if they are at least 10 acres of contiguous vegetation not split or disrupted by roads or other man-made features.
- g. If the vegetated area meets the 10 acre (unsplit by roads) MMU then other vegetated areas adjacent to the “main” vegetation unit but split by roads are added to the vegetation polygon. The minimum mapping width (MMW) for a 10 acre MMU polygon is 333 feet, approximately half the width of a 10 acre MMU square.
- h. Natural vegetation that comes into the settlement from the outside is continued into the Urban area as a natural vegetation type if the natural area meets the 10 acre MMU and MMW. This linework can either be straight or more natural looking.
- i. Snaky vegetation polygons are not created just to meet the 10 acre MMU.
- j. Cleared areas on horse-related property that may be dirt or grass (looks green in most cases) will be coded as 9300 (no dual coding for veg and land use).
- k. Flood Control Basins – are included in the 9310 polygon but are mapped separately as a 9805 if >10 acres.
- l. An exception to the Urban Window criteria of not dual coding an urban window polygon is in the Piñon Hills region of the study area. In this situation, the natural vegetation is composed of a fairly large dense and continuous (non disjunct) cover of California Juniper, Joshua Trees, and Pinyon Pine overlapping the urban window polygon. In order not to lose this continuous overstory of natural vegetation, this area has dual coding of the vegetation and land use within the urban window polygon.

9320 = Anthropogenic areas of little or no vegetation

Isolated scrapes with no apparent built up uses associated with them that are >2.5 acres are mapped as 9320 with an Anthropogenic Disturbance of 3.

Intensely used OHV areas where the vegetation may have been removed because of high vehicle traffic are not considered a 9320. However, cleared or scraped OHV staging areas used for camping or rendezvous are considered 9320. For these situations, a Roadedness Disturbance a value of 3 and the Anthropogenically Altered Disturbance a value of 0 are assigned.

- a. Isolated scraped land and urban built up areas <2.5 acres are ignored. These visible patterns, when < 2.5 acres, are treated within the vegetation polygon by using the anthropogenic disturbance rating or the development rating (both 0 to 3 scales).
- b. When scraped land abuts an urban polygon (9300 or 9310) and is >2.5 acres, it is mapped as part of the 9300/ 9310 polygon.
- c. When scraped land abuts an urban polygon (9300 or 9310) and is <2.5 acres, discretion about how it is mapped is left up to the mapper.

9500 = Exotic Trees

A variety of non-native tree plantings occur in the study area. They are associated with human habitation but not mapped as part of a 9300 or 9310 polygon. These are considered “hortomorphic” as opposed to “agromorphic” classes in the National Vegetation Classification. The exotics that are not mapped as this code are:

1431- *Arundo donax*

1432- *Tamarix spp* (not *T. aphylla*)

9501- Eucalyptus

9501 = Eucalyptus

As a genus, Eucalyptus is widely planted and has some unique habitat values. It can usually be distinguished from other exotic plantings, hence the separate coding. No escaped, self-sustaining stands occur in the mapping area, so Eucalyptus is always considered as a sub-category of exotic trees.

9502 = Tamarix aphylla (athel)

Planted *Tamarix aphylla* (athel) should be considered in this category.

This type has a MMU = 1 acre.

9800 = Water

Artificially created water is mapped in the Vegetation layer as Water (9800) and is coded as 9800 in the Land Use field.

The boundary of a water polygon is determined considering the following items:

- a. A review all the available imagery and topographic layers
 - b. The high water line is used
 - c. The water signature is not cut
 - d. The best delineation that represents all the imagery is used
- 1ac MMU; 10ac in Urban Window
 - Perennial water
 - Usually water body shown on topo map
 - Water in playa mapped as part of the playa (6116)
 - Examples include:
 - Park ponds
 - Residential development with recreational lake
 - Reservoirs
 - Curvilinear shaped duck ponds with water
 - Bermed agricultural ponds with water

9801 = Perennial Stream Channel

This type is restricted to a few locations such as near Mojave River Narrows, Afton Canyon, etc. Water is expected to flow throughout most average years. These polygons are coded as 9800 in the Land Use layer and have a MMU=1 acre.

9803 = Small Ponds & Natural Lakes

This type is coded as 9800 in the Land Use layer. These polygons have a MMU=1 acre.

- Small dammed ponds on creeks contain ephemeral water from natural seasonal flow
- May have naturally ponded water in the Mojave River and Colorado River floodplain
- Bermed ponds in agricultural fields are not included

9804 = Aqueduct (California and Colorado River Aqueducts only)

Aqueducts are coded separately because of their unique attributes as a cement-lined or covered water conveyance system. Only open water aqueducts are mapped as a water type in vegetation (code 9804). The aqueduct is also coded in the Land Use layer attribute as 1436 (Water Transfer). The MMU=1 acre for this type.

9805 = Water Impoundment Feature:

These are typically utility or other straight-edged water bodies impounded by berms and may or may not contain water at time of imagery exposure. The MMU is 2.5 acres. Examples are

Appendix D – Land Use Mapping Criteria

settling ponds, sewage treatment ponds, salt evaporators, non-curvilinear duck ponds (with and without water), curvilinear duck ponds (without water) and bermed agricultural ponds (no water). The Land Use code for this type is 9810 (Water Impoundment Feature).

Water impoundment features are coded with Development Disturbance of 3 and Anthropogenically Altered Disturbance of 3.

The following criteria are provided to give additional clarification for specific situations regarding water impoundment features:

Flood Control Basins – are not mapped as a 9805 unless they are >10 acres in size. Flood control basins <10 acres in size are mapped as part of a 9300 or 9310 polygon.

Duck Ponds – In situations where there is a mix of duck ponds (curvilinear, non-curvilinear, with and/or without water) that are determined to be in current use, these are, as a whole, mapped as 9805 - not separated into individual 9800 and 9805 polygons.

Inactive duck pond areas with shrubs growing in them are mapped as natural vegetation. Water impoundment features are coded with a Development Disturbance of 3 and Anthropogenic Disturbance of 3.

LAND USE CODES (Land Use layer ONLY):

This code is primarily used for polygons coded in the Vegetation layer as some type of Agriculture, water type, Urban or Urban Window. It is also used for polygons that contain mappable Land Use with >10% natural vegetation. These polygons are coded as vegetation types in the Vegetation layer.

If a polygon is coded in the Vegetation layer as an active agriculture type (code 9210 or 9220), then there is a Land Use code assigned in the Land Use layer (either 2100 or 2200). To determine if a polygon is coded as active agriculture, see the description under vegetation code “9200 = Agriculture” above.

0000 = Not Assigned/Not assessed

1000 = Urban - Areas of built-up land characterized by intensive land use, where most of the land is covered by man-made structures because of human activity. See discussion under vegetation code 9300 and 9310.

1436 = Water Transfer (California and Colorado River Aqueducts only) - see discussion under vegetation code 9804 above.

2100 = Non-woody row & field crops – see discussion under vegetation code 9200 and 9220.

2200 = Orchards & Vineyards – see discussion under vegetation code 9200 and 9210.

9800 = Undifferentiated water – see discussion under vegetation code 9800, 9801, 9803, and 9804.

9810 = Water Impoundment Feature – see discussion under vegetation code 9805.

Appendix E

Reconnaissance Field Form

RECON FIELD FORM (May 17, 2011)

Date:	Surveyors (circle recorder):								
Waypoint ID:	GPSname: Projected? Yes / No / Base If yes, enter base Waypoint ID: Bearing: _____(degrees) Distance: _____(meters)								
UID:	Base UTM's / projected UTM's (circle one) UTME _____ UTMN _____ PDOP: +/- Elev.(m)								
Size of stand (acres): <1 1-10 >10 Camera/Photos:									
Field alliance name:									
Comments:									
<table border="0" style="width:100%;"> <tr> <td align="center">%</td> <td align="center">Cover -</td> <td align="center">Conifer Tree:</td> <td align="center">Hardwood tree:</td> <td align="center">Joshua Tree:</td> <td align="center">Tree:</td> <td align="center">Shrub:</td> <td align="center">Herb:</td> </tr> </table>		%	Cover -	Conifer Tree:	Hardwood tree:	Joshua Tree:	Tree:	Shrub:	Herb:
%	Cover -	Conifer Tree:	Hardwood tree:	Joshua Tree:	Tree:	Shrub:	Herb:		
Strata	Species	% cover	Strata	Species	% cover	Strata	Species	% cover	

Date:	Surveyors (circle recorder):								
Waypoint ID:	GPSname: Projected? Yes / No / Base If yes, enter base Waypoint ID: Bearing: _____(degrees) Distance: _____(meters)								
UID:	Base UTM's / projected UTM's (circle one) UTME _____ UTMN _____ PDOP: +/- Elev.(m)								
Size of stand (acres): <1 1-10 >10 Camera/Photos:									
Field alliance name:									
Comments:									
<table border="0" style="width:100%;"> <tr> <td align="center">%</td> <td align="center">Cover -</td> <td align="center">Conifer Tree:</td> <td align="center">Hardwood tree:</td> <td align="center">Joshua Tree:</td> <td align="center">Tree:</td> <td align="center">Shrub:</td> <td align="center">Herb:</td> </tr> </table>		%	Cover -	Conifer Tree:	Hardwood tree:	Joshua Tree:	Tree:	Shrub:	Herb:
%	Cover -	Conifer Tree:	Hardwood tree:	Joshua Tree:	Tree:	Shrub:	Herb:		
Strata	Species	% cover	Strata	Species	% cover	Strata	Species	% cover	

Date:	Surveyors (circle recorder):								
Waypoint ID:	GPSname: Projected? Yes / No / Base If yes, enter base Waypoint ID: Bearing: _____(degrees) Distance: _____(meters)								
UID:	Base UTM's / projected UTM's (circle one) UTME _____ UTMN _____ PDOP: +/- Elev.(m)								
Size of stand (acres): <1 1-10 >10 Camera/Photos:									
Field alliance name:									
Comments:									
<table border="0" style="width:100%;"> <tr> <td align="center">%</td> <td align="center">Cover -</td> <td align="center">Conifer Tree:</td> <td align="center">Hardwood tree:</td> <td align="center">Joshua Tree:</td> <td align="center">Tree:</td> <td align="center">Shrub:</td> <td align="center">Herb:</td> </tr> </table>		%	Cover -	Conifer Tree:	Hardwood tree:	Joshua Tree:	Tree:	Shrub:	Herb:
%	Cover -	Conifer Tree:	Hardwood tree:	Joshua Tree:	Tree:	Shrub:	Herb:		
Strata	Species	% cover	Strata	Species	% cover	Strata	Species	% cover	

Appendix F

Field Verification Form

Field Verification Form: DRECP Vegetation Mapping

(5/2/2012)

Surveyors (circle recorder):						Date:				<input type="checkbox"/>			
Waypoint ID:		GPSname: _____ Projected? Yes / No / Base / Digitized If projected or digitized, enter base Waypoint ID: Bearing: _____ (degrees) Distance: _____ (meters)									<input type="checkbox"/>		
Polygon UID:		Base UTM's / projected UTM's (circle one) UTME _____ UTMN _____ PDOP: +/-									<input type="checkbox"/>		
Strata	Species	%	cover	Strata	Species	%	cover	Strata	Species	%	cover	<input type="checkbox"/>	

Notes: (including recommendations for line-work revision, state of veg. "discernability" based on season and topography, classification interpretation, homogeneity and unusual sightings of plants or animals)		<input type="checkbox"/>
USGS Quad Name		<input type="checkbox"/>
Map Unit (code and name)		<input type="checkbox"/>
Camera/Photos		<input type="checkbox"/>
Conifer Cover	None >0-1% >1-5% >5-15% >15-25% >25-50% >50-75% >75-100% NA	<input type="checkbox"/>
Hardwood Cover	None >0-1% >1-5% >5-15% >15-25% >25-50% >50-75% >75-100% NA	<input type="checkbox"/>
Joshua Tree Cover	None >0-1% >1-5% >5-15% >15-25% >25-50% >50-75% >75-100% NA	<input type="checkbox"/>
Total Tree Cover	None >0-1% >1-5% >5-15% >15-25% >25-50% >50-75% >75-100% NA	<input type="checkbox"/>
Shrub Cover	None >0-1% >1-5% >5-15% >15-25% >25-50% >50-75% >75-100% NA	<input type="checkbox"/>
Herb Cover	0-2% >2-15% >15-40% >40% NA	<input type="checkbox"/>
Exotics	None or not visible 1 2 3 Not Applicable	<input type="checkbox"/>
Development	None or not visible 1 2 3 Not Applicable	<input type="checkbox"/>
Anthropogenic Alteration	None or not visible 1 2 3 Not Applicable	<input type="checkbox"/>
Hydrologic Modification	NO YES Not Applicable	<input type="checkbox"/>
Estimated area of identifiable vegetation viewed	Radius (m) _____ or rough % of polygon viewed from point _____	<input type="checkbox"/>
Is this a "multiple" point assessment?	NO YES if yes: ____ of ____ points for this polygon	<input type="checkbox"/>