

**California Wildlife Habitat Relationships System**  
**California Department of Fish and Wildlife**  
**California Interagency Wildlife Task Group**

---

SIERRA NIGHT LIZARD

*Xantusia sierrae*

Family: XANTUSIIDAE  
R099

Order: SQUAMATA

Class: REPTILIA

Written by: R. Marlow

Reviewed by: T. Papenfuss

Edited by: S. Granholm

Updated by: CWHR Program Staff, May 2013

#### DISTRIBUTION, ABUNDANCE, AND SEASONALITY

The Sierra night lizard is restricted to the Greenhorn mountains in the southwest Sierra Nevada. It is found in association with yucca, foothill pine, chamise, pinyon pine and juniper. This is a secretive lizard, spending most of its time in and under yucca logs and other cover. This species is found at elevations of 300 to 2070 m (990 to 6800 ft) (Macey and Papenfuss 1991). Activity may begin in early April at low elevations and last until early fall, while emergence may be retarded until late spring at higher elevations.

#### SPECIFIC HABITAT REQUIREMENTS

**Feeding:** This lizard eats termites, orthopterans, small beetles, homopterans, hemipterans, collembolans, moths, caterpillars, flies, ants, ticks, and spiders. It waits for prey items that wander into the cover area and seldom searches actively (Brattstrom 1952, Stebbins 1954)

**Cover:** Most commonly this species is found under and in logs of the several species of yucca, and much less frequently under logs and debris of foothill pine, root channels of creosote, and other natural or unnatural cover objects (Brattstrom 1952, Stebbins 1954).

**Reproduction:** Courtship and parturition take place in the shelter of yucca logs or whatever cover is utilized.

**Water:** Probably does not require water but will drink in captivity.

**Pattern:** A closely related species, the desert night lizard is secretive and reclusive. It feeds, reproduces, and lives most of its life under cover (usually yucca logs). It is difficult to overestimate the importance of cover to this species.

#### SPECIES LIFE HISTORY

**Activity Patterns:** Populations of the closely related desert night lizards at low elevations become active in early April and remain so until late summer, while higher-elevation populations may not become active until May. Daily activity patterns are difficult to determine, due to secretive behavior. Laboratory studies have demonstrated that the desert night lizard is diurnal (LaPointe 1966, Mautz and Case 1974).

**Seasonal Movements/Migration:** None.

Home Range: The home range is restricted to the cover site and the area immediately adjacent to it (Miller 1951).

Territory: Both males and females are territorial during the breeding season (Miller 1951).

Reproduction: Copulation occurs in late spring and egg development takes about 3 months. The young are born in September and October with an average of 2 young in a brood (range 1-3). The mother attends the young for a short time after birth (Cowles 1944, Miller 1948, 1951, Stebbins 1954).

Niche: Predators probably include most larger lizards (*Crotaphytus*, *Gambelia*, *Sceloporus*, etc.), many snakes, and predatory birds. This species rarely occurs away from cover, and the most commonly-used cover objects are yucca logs. Replacement of yucca logs is slow and their destruction by collectors can have a deleterious impact on local populations.

#### REFERENCES

- Brattstrom, B. H. 1952. The food of the night lizards, genus *Xantusia*. *Copeia* 1952:168-172.
- Cowles, R. B. 1944. Parturition in the yucca night lizard. *Copeia* 1944:98-100.
- LaPointe, J. L. 1966. Investigation of the function of the parietal eye in relation to locomotor activity cycles in the lizard *Xantusia vigilis*. Ph.D. Thesis, Univ. California, Berkeley. 95pp.
- Macey, J. R. and T. J. Papenfuss. 1991. Reptiles. Pages 291-360 in C.A. Hall, Jr., editor. *Natural History of the White-Inyo Range eastern California*. Univ. Calif. Press, Berkeley, California. 536 pp.
- Mautz, W. J., and T. J. Case. 1974. A diurnal activity cycle in the granite night lizard, *Xantusia henshawi*. *Copeia* 1974:243-251.
- Miller, M. R. 1948. The seasonal histological changes occurring in the ovary, corpus luteum and testis of the viviparous lizard, *Xantusia vigilis*. *Univ. Calif. Publ. Zool.* 47:197-224.
- Miller, M. R. 1951. Some aspects of the life history of the yucca night lizard *Xantusia vigilis*. *Copeia* 1951:114-120.
- Stebbins, R. C. 1954. *Amphibians and reptiles of western North America*. McGraw-Hill, New York. 536pp.

R099

---

Life history accounts for species in the California Wildlife Habitat Relationships (CWHHR) System were originally published in: Zeiner, D.C., W.F.Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. *California's Wildlife*. Vol. I-III. California Department of Fish and Game, Sacramento, California. Updates are noted in accounts that have been added or edited since original publication.