Species Notes for Sandhill Crane (*Grus canadensis*):

California Wildlife Habitat Relationships (CWHR) System Level II Model Prototype



William F. Laudenslayer, Jr. (author)¹ and Monica D. Parisi (editor)²

California Department of Fish and Game California Interagency Wildlife Task Group

October, 2007

¹ United States Forest Service, Pacific Southwest Research Station (ret.)

² California Department of Fish and Game, Biogeographic Data Branch

PREFACE

This document is part of the California Wildlife Habitat Relationships (CWHR) System, operated and maintained by the California Department of Fish and Game (CDFG) in cooperation with the California Interagency Wildlife Task Group (CIWTG). The information will be useful for environmental assessments and wildlife habitat management. For more information on the CWHR System and all of its components, please see http://www.dfg.ca.gov/biogeodata/cwhr/.

Notes such as these were prepared for 32 species by the US Forest Service Pacific Southwest Research Station as part of a 2000/2001 contract with CDFG. Each is part of a prototypical "Level II" model for a species. As compared with the "Level I" or matrix models initially available in the CWHR System, "Level II" models incorporate spatial issues such as size of a habitat patch and distance between suitable habitat patches.

The notes are divided into three major sections. First, "Distribution, Seasonality and Habitats" represents information in the existing Geographic Information System (GIS) range data and in the Level I matrix model for a species. There is a vector-based GIS layer of geographic range and seasonality for each species in CWHR as well as a matrix containing all suitability ratings – High (H), Medium (M), Low (L) or Unsuitable (-) – by habitat (e.g. BOW or Blue Oak Woodland), stage (e.g. 4P or small tree, open canopy) and life requisite (reproduction, cover, or feeding.). Tools such as "Bioview" within the CWHR software will return these suitability ratings for a species to a user-supplied data set containing habitats and either stages (e.g. 4P) or stage values (e.g. trees of 16.0 average diameter at breast height in a stand of 30% canopy closure).

Second, "Required Attributes of Suitable Habitat Patches" represents spatially-explicit requirements of a species. The information here builds upon what is known about habitat patch size and the most critical attributes of a habitat patch needed by an individual of the species. Applications such as "GRABS", which stands for "Grouping Resources Algorithm for Biological Data Sets", will "clump" pixels of a user-supplied raster-based GIS data set representing patches of a suitable habitat and stage for a species. It will calculate area, perimeter, and complexity within each patch and analyze its outside edge for juxtaposition with other habitats and stages of interest. Many of the attributes are what were once called "elements" in the CWHR model.

Third, "Spatial Habitat Requirements for Persistence of Population" represents estimates of the amount of habitat needed to maintain a population of a species. This may be considered the starting point for a "Level III" CWHR model, which would take into account spatial issues as well as a number of population parameters not yet incorporated into CWHR. Such information is included for most, but not all, Level II-modeled species.

B150 Sandhill Crane Grus canadensis

Distribution, Seasonality and Habitats

Model Parameter	Threshold Value(s) for Species
Biogeographic Range and Seasonality range of the species, by season, in the state	Species breeds in or near montane meadow- emergent wetland areas in northeastern California and winters in lower elevation grasslands or wetlands primarily in the Sacramento and San Joaquin valleys.
Suitable Habitats and Stages habitats rated in the California Wildlife Habitat Relationships (CWHR) System as high (H), medium (M), or low (L) suitability for reproduction, cover, or feeding	Species finds suitability (H>L) for reproduction, cover, and/or feeding in some or all stages of: Annual Grassland, Dryland Grain Crops, Fresh Emergent Wetland, Irrigated Grain Crops, Irrigated Hayfield, Irrigated Row and Field Crops, Lacustrine, Rice, Valley Foothill Riparian, and Wet Meadow.
Water whether water is required, enhances, or is irrelevant for habitat suitability	Water is required. Shallow water is needed for feeding, cover, and sometimes reproduction.

Required Attributes of Suitable Habitat Patches

Model Parameter	Threshold Value(s) for Species
Patch Size	minimum nesting/roosting patch
L = low suitability. This is the	0.5 acre (L)
minimum patch size for	5 acres (H)
persistence of an individual.	
	minimum foraging patch (shallow water of ponds or lakes,
H = high suitability. Above this	wet meadows, grasslands, open wetlands) for persistence of
patch size, area alone does not	an animal
increase habitat suitability for	5 acre (L)
an individual.	50 acres (H)
	distance between - nesting/roosting and foraging patches
	5 miles (L)
	< 1 mile (H)

Edges requirements for a transition between two life form types – tree/shrub, tree/grass, tree/water, tree/agricultural, shrub/grass, shrub/water, shrub/agricultural, grass/water, grass/agricultural, or water/agricultural	A grass/water edge or an agriculture/water edge is preferred but not essential for reproduction, cover and feeding.
Structural Habitat Attributes requirements for live vegetation, dead or decadent vegetation, vegetation residues, physical features, or human- made features	Emergent aquatic vegetation is essential for reproduction, cover and feeding. Gravel and organic soils are preferred but not essential for feeding.
Food vegetative or animal diet requirements	Graminoids, forbs, grains, roots, or invertebrates are essential. Species also eats seeds, berries, fruits, acorns, amphibians, reptiles, small birds and small mammals.