BALTON SRA NATIONAL WILDLIFE REFUGE

AND

WATERFOWL DEVELOPMENT AREA

HARRATIVE REPORT

JANUARY, FEBRUARY, MARCH, APRIL

1957

XXXXXXXXXXXXXX

UNITED STATES DEPARTMENT OF INTERIOR FISH AND WILDLIFE SERVICE BRAWLEY, CALIFORNIA



CONTENTS

PERS	omic	L	1
I	C M	ERAL CONDITIONS	
	A.		2
	В.		2
	Ċ.	Fires	3
II	AIL	DLIFE.	
	A.		_
		1. Populations and Behavior	5
		2. Geose and Swans	3
		5. Ducks	5
		4. Water, Marsh, and Wading Birds	5
		5. Shorebirds, Gulls and Torns	5
		6. Discaso	7
		Upland Game Birds	5
	C.	Other Birds	3
	D.	Fur Animals, Predators, otc	9
		Predaceous Birds	ううりりりりりちんる??
		Mah	7
	G.	Waterfowl Depredations	7
111	nsf	UGE DEVELOPMENT AND MAINTENANCE	
	A,	Physical Developments	7
	4-4	1. Cultivated Crops	7
	₿.	Receipt of Seed and Stock	8
IA	BCC	NOMIC USE OF REFUGE	8
V.	FUE	PLIC RELATIONS	
	A.	Recreational Vacs	8
	В.	Refuge Visitors	8
	Ç.	Refuge Participation	9
	D.	Bunting	9
	E.	Publicity	999
	F.	Violations	9
VI	APF	LIED RESEARCH	10
VTT	OTT	TER ITHMS	12

REFIGE PERSONNEL

Regular Personnel

roward J	. 0	N	ZI:	LL	•	•	•	•	٠	•	•	٠	•	•	•		•	٠	٠	•	•	. Refuge Manager . Refuge Manager, Asst
WILLIAM	NUE	88.	•	•	•	•	•	•	•	•	•	•	•	٠	•	٠	•		•	•	٠	 Refuge Manager, Asst
														(1	'n.	a P e	i.	M.	lni	de	k	Refuge, Idahe)
Joseph L	. C	UDI	ĐΥ	•	•		•	٠	•	•	•	•	٠	•	•	•		•	•	•	•	. Clerk-typist . Mechanie, Hwy. Duty
ekiry st	EER	•	•	•	•	•	•	٠	•	•	٠	•	•	•	•	•	•	•	•	•	•	. Mechanic, Hwy. Duty
Jose' Ba	RRO	S	•	•	•	•		•		•	•			•	•		•	•	٠	•	•	• Tractor Operator • Tractor Operator • Tractor Operator • Tractor Operator
CARL W.	FOR	D					•		•			•	•		•	•		•	•			. Tractor Operator
ALFRED W	¥	C 1	ΓĀJ	RÌJ	NI)	_								•			•	•			. Tractor Operator
CI YDE W.	ST	Her.	A TO	*		_	Ĭ	_	_	_	_	_	_	_		_			_	_	_	. Tractor Operator
CHEST PT	wit t	1.7	W	* 2	•		•		•	Ī	•	•	-	-	-	-	•	•	•	•	•	• Tractor Operator
lko e. o																						
																						. Irrigator
SYLVESTE	R B	ARI	RO	3		•	•	•	•	•	٠	•	•	•				•	•	•	•	. Irrigator
MANUEL C	ARD	OP7	05			•																. Irrigator
JULIO RI	BER	IO	•	•	•	•	•	•	•	•	•		•	•	•							. Irrigator

Temporary Personnel

Mone during period.

BARRATIVE REPORT

I GENERAL CONDITIONS

A. Weather & Conditions

imperial Valley experienced cooler days in January, 1957 than it did during the corresponding menth of 1956. Mean average temperature was 54 degrees or about 3.5 degrees lower than January 1956. Mean maximum for January of this year was 66.5 degrees, compared to 72.9 degrees for January 1956. Mean minimum was 42.7 degrees, just slightly lower than the same period of 1956. There were 11 clear days, 6 partly cloudy days and 14 cloudy days during January. On the 25th a heavy fog capped the Valley from about 7:00 to 9:00 AM.

February 1957 was warmer than the corresponding weeks of 1956. Ween temperature this year was 63.7 degrees or 9.4 higher than February, 1956. Mean maximum this year was 77.6 degrees. The mean minimum was 49.9 degrees ----- 11.1 above the minimum for February, 1956.

Weather conditions during March and April were rather mild compared with previous seasons on record. Maximum temperature during March was 92 degrees, during April, 94 degrees. Minimum temperature in March was 49 degrees, in April, lowest temperature recorded was 49 degrees. Winds which generally rip through Emperial Valley in March and April were lacking this year.

Tabulated below is the weather data as compiled by the El Centro Naval Air Station, Seeley, California.

MONTH	MAX.	MIH.	PRECIPITATION	DAYS OF #25MPH WINDS
January	73°	35°	0.61*	
February	90 ⁶	36°	0.10"	
March	9 2º	430	0.09*	6
April	84°0	149°	T.	17
	Totals		0.80"	25

B. Precipitation & Water Conditions

Rainfall for January totaled 0.61 inches. Storms ecoured on January 3, 7, 8, 9, 29, and 30. The total was about .13 higher than that measured one year ago. The general area storm

of the 28th and 29th put considerable variety in the winter weather for local residents. Howling winds, spiked with rain, sleet and snow left the mountains west of Salton Sea covered with 4 feet of snow. For the first time in the memory of most citizens the Chocolate mountains, east of Imperial Valley, were whitened with a light snow blanket.

Precipitation for February totaled 0.10 inches. Traces of rain fell on the 21 and 22nd but main moisture was recorded on the 23rd when .08 inch fell and the 28th when .02 inch was received.

A light earthquake joited the general Imperial Valley area about 12:15 PM February 23rd. No damages were noted but the tembler was strong enough to rattle dishes and utensils in most everyones home.

For a period of about two weeks light earthquake tremors were felt daily at the refuge in early April. On April 25 four separate heavy shocks jarred the entire Emperial Valley area. Worst damage done was within a narrow 2-mile wide strip about 1 mile south of refuge Unit II extending about 4 miles in a southwesterly direction. The hard ground was left broken and cracked. In the fields, canals, and roads, water and fine sand flowed to the surface through "sand-boil" fissures. The boils flowed for about 3 hours leaving large acreages of wet fields. The Vail 4 road (4 miles) was about three-fourths under water. About 2,000 acres of farm land south of refuge sub-headquarters was affected by the boils. In the wake was mounds of sand in cotton rows spread over a dozen adjacent rows leaving high spots and cracked ground. Undoubtedly a lot of land will require releveling. Tiled lines too, appeared to have slipped and mudded in here and there.

C. Fires

No fires during period.

II WILDLIFE

A. Migratory Birds

1. Populations and Behavior

Geese and Swans

Hunting pressures both on and around refuge Unit II caused unprecedented shifting of populations and feeding sites among the geese.

Although Canada goose hunting after December 15th was illegal, nevertheless some of the hunters persisted both through ignorance (other counties were not elesed) and nafarious attitudes.

A number of Canada goose cases were made by wordens. At least 5 Canadas were shot on the refuge during the balance of the season.

Another contraband species, the Ress' goose, owing to it's habit of mingling and flying with canada goose on refuge Unit II, received very heavy gun pressure for such a tiny flock. Of the 31 Ross' goese here during the previous period only 4 remained at the close of the season.

Snow goese, although showing the effects in numbers of a flock reduction over last year, seemed to fair well. Most of the Snows moved into refuge Unit II and shuttled between cattail fields and the Alamo river delta or open sea during the day.

A single Blue goose was observed January 29th.

Returns from banded Canada geose which were killed on Salton Sea Refuge originated at Bear River Refuge, Utah, Neponset Reservoir, Utah, and Pathfinder Reservoir, Wyoming.

On March 20, when the last general serial census was made some 500 Snow geese, 1 White-front, and 5 Canada geese were still here.

In previous reports the figures taken from annual census or winter inventory totals have been pointed out to show the definite downward trend of geese in this part of the flyway. Although total number in the Salton Sea area appear insignificant to some they nevertheless are real and represent the general attitude or negligence to keep closer tab on certain species locally. (Canada geese have shown a drop from 4150 in 1952 to 300 in 1956. Snow geese have dropped from 20,830 in 1952 te 9,000). It is interesting to note how the trend is reflected in fugures for the Colorado River. Population figures for Arizona counties along the river, taken from 1950 through 1957 are as follows:

Canada and Snow Goose Populations-Lower Colorado River

Year				Ar	Lzona Cou	nties		
(Jan.)	G1	la	Mar	icopa	Moj	AVE	Yum	Maria
•	Snow	Can.	Snow	Can,	Snow	Can.	Snow	Can.
1950		2304		170	1200	2500	14.3	1853
1951		1826		215	1150	9 5 5	430	1280
1951 1952 1953	7	1150		282	885	985	500	1612
1953	•	1135		26 0	11,00	946		473
1954	5	1907		437	1817	830	143	315
	Ž	1486		527	900	401		270
1956	17	2110	1	527 604	232	152	63	611
1955 195 6 195 7	10	2388		350	50tt 535	152 254	2	510

Daoks

Duck species continued to use the refuge similar to population numbers and pattern mentioned in the previous report.

Shovelers, which had shown lower numbers this year, showed local population increases through March and early April.

Widgeon remained stable population-wise until March 15th when they showed signs of dwindling population and migration. Most duck species appeared to have moved from Salton Sca slightly earlier than previously.

Fulvous-tree ducks were first noted when 9 were seen March: 50th.

Discase

No known diseases occured during the period.

Shorabirds, Gulls and Torns

We change in status noted.

Water, Marsh and Wading Birds

No change in status noted.

An estimated 10 Common Egrets were found incubating eggs in small colony at the north end of the sea March 20th.

Food and Conditions

No change noted.

B. Upland Came Birds

He change in status noted.

C. Other Birds

An estimated 500 Mountain Bluebirds were present at Unit II from January 20 to March 5th.

A Mourning Dove banded at this station March 13, 1953 as an adult bird was re-trapped here March 20th.

A light influx in Mourning Doves occured about January 20th at which time a definite increase was noted. By January 24 the increase was quite pronounced. By February Lith the species was again common.

One ground dove was seen near trifolium 15 canal at Unit I February 12.

Robins were present from January 8 to April 15. About 500 were at Vendel's Corner February 7 where they were feeding on dates. February 20th about 500 were observed at Brawley feeding on pyracantha berries.

Some 50 Ecuntain Plover used Tract C, Unit I from February 4 to about February 20th.

A flock of about 1500 Horned larks remained at tracts 18-19 Unit I from February 5 to April 15th, and Rough-winged smallows were observed here in mid-March.

A pair of Verding were observed feeding a young verdin at Unit

Pine sisking were observed east of Unit II February 9.

Mr. William Anderson of California Fish and Game Department observed 1 Ladder-backed woodpooler near Finney-Ramer February 9.

Mostern Kingbirds were first seen April 5th.

Readminiers were observed constructing nests as early as February 19th.

Night Hawks arrived about March 28 when 4 were observed at Unit I.

Shrikes, which nested at headquarters, had young on the wing by March joth.

Red-winged Blackbirds were actively constructing nests in mid-April.

Several Kinglishers were seen throughout the period.

A single Phainopepla showed up at Unit I April 25.

D. Fur Animals, Predators, etc.

No change in status noted.

A Coyote was observed a few times at Unit I in mid-February.

B. Predaceous Birds

One Caprey was seen March 5th near Calipatria.

One Golden eagle was observed often in early February.

On February 7th a lone Swainson's Hawk was seen at Unit II.

Cooper's and Sharp-shinned hawks occured in the general area as uncommon visitors.

F. Pish

University of California biologist continue to predict a tremendous sportsfishing resource at Salton Sea from the ocean fish transplanted here in recent years. Latest prediction is that good fishing will be enjoyed within the year 1957.

Corvina were found abundant and believed to have spawned in February. Since 1948, when 1000 adults were introduced, it is estimated 65,000 now inhabit the sea.

G. Materfowl Depredations

Complaints of crop damage by waterfowl and particularity Widgeon reached a new low. This does not infer that losses were not received. It does indicate, we believe, that without spokesmen and general rabel-rousing the problem, although present, didn't receive the publicity of other years.

Alfalfa damages were noted on the Seibert Farms 3 miles cast of refuse headquarters at Gate 155 Trifolium 8 canal March 2;, Mimore ranch, John Williams place one-half mile south of headquarters, C.T. Dearborn and others along Trifolium 12 canal also reported some widgeon damages.

The widgeon showed an increased population over last year. Alfalfa prices dropped slightly during the winter months. Rains, which rendered alfalfa fields too wet for grazing, caused a surplus of livestock feed and alfalfa for dehydration meal.

III REFUGE DEVELOPMENTS AND MAINTENANCE

A. Physical Developments

Development work was confined to land clearings at Tract 9 Unit I and Unit A for contours and eventual leashing.

Cultivated Crops

Cropping program in previous report continued. Lands in Unit A are being prepared and about 200 additional acres will be brought under leach this summer. Approximately 2; acres of mature barley, 30 acres of Sudangrass, 1300 acres of cattails and wild millet and 900 acres of green barley are planned for fall-winter use.

B. Receipt of Seed and Stock

During the period 6,000 pounds wild millet seed was purchased at Secremento, California.

Approximately 18 tons of outs were shipped from Tule Lake Refuge.

IV ECONOMIC USES OF REFUGE

Mone in effect during the period.

V PUBLIC RELATIONS

A. Recreational Uses

Eighty members of the Falomar Club toured the refuge January 26.

Sixty-two visitors checked in at headquarters during the period. A total of 1450 people visited the refuge including hunters who used the Lea Act lands.

B. Refuge Visitors

Name	Date	Identification	Purpose
Mr. Graham Elmore	1/15	County Read Supt.	Gravel haul-Red hill
Mr. Paul Quick		Asst. Rog. Dir.	Inspection tour
Mr. K.F. Mao Donald		Reg. Refuge Supv.	Inspection tour
Dr. H. Lee	1/26-29		Photography
Mr. AMrs. Deveroux Butcher	2/5	Nat'l Park Assn.	Photo & observation
Mr. L. Me Kibben	2/11	Reg. truck driver	Sood haul
Mr. Phil. Douglas	3/1	Cal. Fed Dept.	Boat launching site
Mr. Wendell Miller	3/15	Cal. PAG Dopt.	Experiment plantings
Mr. Bob Reynolds	3/22	I.I.D.	Water order
Mr. Wes. Fleming	4/57	Aris. Biologist	Contact
Mr. Chas. Copley	4/57	Aris.Area Supv.	Contact
Mr. AMrs. fom Brown	14/0	San Diego Audubon	Observation
Mr. Lop L. Laythe	1/8	Regional Director	Inspection tour
Mr. dire. Bugene Cardiff	4/10		Bird collecting
Mr. E.R. Quartrup	4/30	San Diego Cty. Vet.	Tour
Mr. Paul Williams	Several		Contect
Mr. C. Lostetter	Several	Biologist	Contact
Mr. A.W. Elder	Several	Game Mgmt. Agent	Contact
Mr. Jim Johnson		Came Egat. Agent	Contact
Mr. Raymond Gallen	Severa l	Pilot-biologist	Acrial census
Mr. Frank Baldan		Oul. Fed Dept.	Menting & patrol
Mr. John Parrish	Mumerous Visits	g Cal. Pag Dopt.	Contact

C. Refuge Participation

法的证券者

Mr. William Anderson and Mr. Gale Horn of California Fish and Game Department continued waterfowl trapping activities on the refuge throughout the period. Traps were set up at Tracts 9, and 11, Unit I and 7-11, Unit II was well as Finney-Ramor, Hazard area, and Wister unit.

April 6 a joint meeting of Service and California Fish and Game personnel w/the So. California Duck Hunters Assoc. and Federation So. California Sportsmen was attended at the Laffayette Hotel, San Diego, California.

Local Scout troops participating in conservation activities visited and toured the refuge. Refuge personnel served as wildlife conservation merit badge counselor for troops in Brawley and Calipatria.

D. Bunting

Restricting the acreage of Lea Act lands opened to hunting to 50% of what it was one year ago appears to have been a step in the right direction - now that the canada goose population has been scatterguined down to a more remnant of former numbers.

B. Publicity

During the period another 6500 words were added to the weekly column "Wildlife Commente".

The Miland Tomato Festival brochure featured an article entitled "Desert Ducks" which covered refuge operations and objectives.

F. Violations

Court proceedings of apprehensions made by refuge personnel and not reported in the previous narrative are listed below:

XAMO	Violation	Date	Amount
Jerry Lee Young-	Munting on refuge	11/21/56	35.00 -
Spring Valley, Calif. William A. Richards-	w # W	12/7/56	\$100.00 -
Redondo Beach, Calif. Bon Aland-	Pos. of 1 Snow goose	12/7/56	#100.00 /
El Segundo, Calif. Thomas R. Rrank-	Hunting on refuge	11/17/56	\$ 35.00 -
Garden Grove, Calif. P.A. Luke-	Shooting waterfowl	12/15/56	♦ 75.00 1
Redlands, Calif. Robert H. MeKinney- San Yeidro, Calif.	after legal shooting ! Bunting on refuge Three Snow geese seis	1/4/57	\$100.00 -

Hame (Cont'd)	Vio	lation	Date	Asount	
Pierce Sherman-		fter legal	12/8/56	\$ 35.00	r
Malibu, Calif. Robert K. Derigo-	shooting Bunting o		11/17/56	\$ 35.00	*****
Anaheim, Calif. Bugene J. Sneed-	# N	**	11/17/56	\$ 35.00	
Seal Beach, Calif. Paul Landers-	# W	# ^			_
Garden Grove, Calif			11/17/56	\$ 35.00	
Albert H. Saunders- Los Angeles, Calif.	,	· · · · · · · · · · · · · · · · · · ·	10/20/56	\$ 35.00	-
Donald R. Helland- Los Angeles, Calif.	*	#	10/11/56	\$ 35.00	maus.,
Robert L. Gebert- San Diege, Calif.	*	#	12/7/56	\$ 35.00	WATER

VI APPLIED RESEARCH

Supplementary Feeding

During the period another supplemental green feed experiment was attempted similar to work done once by Bert Wardwell and twice by present refuge personnel.

The following shows dates and amounts of chopped alfalfa put out at refuge Tract 9, Unit I.

Date	Pounds	Delivery No.	Operator
1/21	2620	3-6826	Jose' Barros
1/25	3550	6884	Jose' Barros
2/4	1640	7040	Jose' Barros
2/5	1630	7062	Jose Barros
2/4 2/5 2/6	1190	7078	Joseph L. Gaddy
2/7 2/8	1600	7093	William Muses
2/8	1890	7204	Edward J. O'Neill
2/11	4130	7137	Jose Barros
2/12	al ₁ 30	7152	William Moss
2/13	3140	7166	Jose Barros
5/11	1820	7179	Joseph L. Ouddy
2/15	5530	7195	Jose Barres
2/16	3940	7201	Clinton Lostetter

Some sixty tons of alfalfa were contracted for by biologist Clinton Lostetter. It was necessary to travel to the East Brawley, Fudge Milling Company, for each load. All leading was done by the truck operator by pitch fork. The stage bruck was weighed and reweighted each trip.

It was predetermined that alfalfa would be spread on the water at fract 9 in four locations where widgeon loafed daily. It was

decided that as rapidly as the feed was taken by widgeon more would be obtained and set out.

Er. Lostetter, accompanied by Agent Jim Johnson, hauled and put out one load of feed. All other feeding and observations was by refuge personnel as indicated on the foregoing tabulation.

Below is listed the sailient points of the feeding observations:

Date	Observation
2/4	time, 4 PM; estimated 500 widgeon at feeding station. When feed spread 5,000 widgeon present.
2/5	time, 9 AM; estimated 450 widgeon at feeding station. Alfalfa about 30% cleaned up; less than 25% of widgeon
2/6	around feeding stations; 1500 widgeon present. time, 9 AM; estimated 300 widgeon & 30 coots at Tract 9; alfalfa 50% cleaned up.
2/7	time, 9:30 AM; estimated 3,000 widgeon, 200 coots, 200 pintails; about 500 using alfalfa feeding stations.
5/8	time, 10:30 AMI estimated 1,000 widgeon, 350 coots in general area, about 200 at stations.
2/9-10 2/11 2/12	No supplemental feed put out.
2/11	time, 9 AM; estimated 5,000 widgeon in general area.
	time, 9:30 AM; estimated 3,000 widgeon in general area, about 150 at feeding stations.
2/13	time, 845 AM; estimated 2,000 widgeon in general area, about 500 at feeding stations.
ड/ग्रो	time, 9 AM; estimated 7,000 widgeon in general area, about 0,000 at feeding stations.
2/15	time, 12:30 PM; estimated 300 widgeon in general area, about 50 at feeding stations.

February 21 through March 1, 1957 (except February 23 & 24), eight sacks of oats were distributed each day at Tract 9. An estimated 2,000 birds, mostly pintails, green-winged teal and widgeon were held during the time feed was being distributed.

A total of about 15 tons chopped alfalfa and 6,00 pounds of oats were put out as supplemental feed to hold widgeon.

It is our spinion that the supplemental feeding efforts hold insufficient widgesn numbers to warrant its use as a depredations proventative. Each feeding required 6 man hours to load, weigh, haul, and unlead.

Refuge Crop Manipulation

Experimental work with widgeon to encourage more use of refuge green crops was continued through January and February. The project entails encouraging widgeon to move inland to flooded rest areas where goese are feeding on barley crops. By providing a condition of protection and minimum disturbance, the species was gradually concentrated at fracts 1-2, Unit I through the use of cats.

By spreading cats on the water daily; widgeon moved in, consumed the cats and gradually joined canada and white-fronted goese grazing in adjacent barley crops.

On February 2, it was estimated 10,000 widgeon were using tract in Unit I daily. On February 9, 8,000 were estimated using the area exclusively. On February 10th we started flushing the widgeon from the general area to keep them from killing the barley. It took three days to discourage use of the green barley field by widgeon. Two days later damages occurred on the Seybert place.

Since we have succeeded in creating this condition two years new it is planned to elaborate on the work and hold the widgeon assumed as possible next winter.

VII OTHER ITEMS

During the period the California Fish and Game Department under the direction of Wendell Miller, transplanted samples of shoal grass shipped in by the State of Texas.

Plantings were wrapped in cheese cloth and submerged near Mullet Island under the supervision of University of California fisheries research biologists.

At the same time we were offered samples for transplanting at refuge Unit I. A meeting was scheduled and preparations made for the work. Miller must have been delayed. He never arrived with or without the planting stock.

It is recalled how transplants of this species set out a few years ago proved unsuccessful.

The Salton Sea Beach Retates, a land subdivision developed by Ed. Jorgensen and Cal Brown of Riverside County, received approval March 17th by the Imperial County Planning Commissioners. The subdivision is along the south east shores of Salton Sea near Bombay beach. It was jointly agreed by the estate and I.I.D. that no land would be sold below -220 feet contour without a waiver exempting I.I.D. from claims in case of rise in the sea.

In early Fobruary it became known that a strong possibility exists for the Miramar Maval Air Station being moved to Salton Sea. Final decision seems to hinge on the City of San Diego which is pressing for an international airport. Such would conflict with the air pattern at Kearney Mesa. Of course Imperial Valley Board of Trade, Navy League, Industrial Development Commission, Chamber of Commerce and County Supervisors are urging use of Salton Sea.

On February 20th Beverley Hills Contractor Barl Brown was awarded a contract to construct a \$ 63,896 beat basin at the Salton Sea State Park about 8 miles Southwest of Meson on Highway 111.

A graveled parkway for 50 cars, concrete boat launching ramp, 3-acre boat basin 10 feet deep by 10 foot wide are included in the project.

future plans call for more camping grounds, and picnicking areas.

_-----

Tractor Operator Carl Ford made local headlines in the Brawley News February 16th when he 'dozed up and killed a 5-foot, limentie diamond-back snake on east Unit "A" of the refuge.

The suit-squabble ever Colorado River water being waged between Arisons and California continued through the period at San Francisco with most of the testimony being presented by Pale Verde, Coachella and Imperial Valley water districts.

With the Commence

The California Fish and Game Department's "Outdoor California" for February 1957 featured an article entitled "Salton Sea Project Seeks Reasons For Steady Decline of Mullet Fishery", by L. J. Hendricks. The article relates how mullet have apparently failed to reproduce or enter the sea.

During the period Biologist Bob Smith and Horton Jensen of the Service and Chester Kibbs of Oregon Fish and Game Department spent three weeks at El Centro awaiting clearance to make the annual waterfowl inventory flight into Mexico. After contacting these men and gathering and returning cameras, sidearms, etc. three times during the interim, it developed that negotiations could not be made with Mexico to count any pert of the countries waterfowl population.

During the period the wrath of angry hunters was unleased full force against the move in California to outlaw Hourning dove hunting. Although a bill was introduced at Sacramento by the proponents of the kill ban, it was squelched by law makers for a two year period pending committee study.

Respectfully submitted,

Heard . O'Nelli Befuge Manager

Approved	

CREDIT DEPARTMENT: Credit is due Mr. Cuddy for compiling banding recovery tabulations, preparing MR Forms, and typing this report.

SALTON SEA REFUGE BANDING RECOVERIES - 1956 - 57

:s	Baldpate	Cinn. Teal	Coot	C-N Teal	Fulwous	Pints 11	Redbead	Paddy	Mallard	Ceneda	aous Joos	#-# 6008e	Shove ler	Canvasback
-					-	. 144		14			03 6			
	!													
	2		his and and over t		And the case of the case of						į			
		1		THE PARTY OF THE	TOTAL SECTION AND ASSESSED.									
	7					17		!						
	200	2	7	<u>.</u> .	1	546	18	1	2	l	3	1	2	
	12		44.0	5		_38) Ber Manach der von	-			
	i					3			i i i				[-
	1	ĺ		1		5	Sheate Mean of					1		
												- أ		
						2					damen ares			
	lasses, mantes					13	and the second				<u></u>			
	3	7	_2	6		58	. 1							
			and the contract of			1	ang anggrass, les gerre	Name alsonore with a sign		-			-	
	Transmission of	1				2			1					
	1				ar ann ann an an an an							<u> </u>		ļ.,
	Da con atomico		India and the			1		amouga soundeadhair t						<u> </u>
		1				1	and a marchist for the con-		-	De reflexaçõe admin		-		ļ
	1					3			-			<u> </u>		<u> </u>
						2		<u> </u>			<u> </u>	<u> </u>		
,				1			-		-	-		_		ļ.,
	-] 	<u> </u>	ļ			-	-
	-			1		2		-	-		-			
٠	·		-	<u> </u>			-		_	ļ.,	-		-	
•	-	1_1		<u> </u>			ļ.	ļ		-		4	<u> </u>	<u> </u>
) 	-	-	5		<u></u>	,	-		-		
	-		-	1_1		_66		į	ļ	L		- -	<u> </u>	-
	_113			8	-	_ 55.		-	-					-
	1				1	1]	-	<u></u>	1		

TOTALS- 272 35 9 67 1 824 19 1 2 2 3 1 2 1

3-1750 Form NR-1 (Rev. March 1953)

WATERFOWL

19 4 388 8 28 8 3 1 * * දු 3 28 38 8 8 2 perio \$ MONTHS OF 888 28 8 8 8 Aertal Cenema 8 8 e porting 22 00 V 8 88 8 88 88 (2) 8 8 88 8 200 8 88 答 o f 2 00,60 00,00 00,00 00,00 8 kg 8 8 22 • • • 3 38 8.00 1.00 1.00 1.00 8 8 R & 32 191 **88** 8 8 8 88 松 Salten See Green-winged teal Blue-winged teal White-fronted Cinnamon teal Other - Mege* Species Ring-necked <u>E</u> Canvasback Bufflehead Goldeneye Whistling Trumpeter Cackling Baldpate Shoveler Ducks: Mallard Redhead Pintail Gadwall Geese: Canada Brant Scaup Black Ruddy 0ther Wood Snow Blue REFUGE Swans: Coots

3 -1750a Cont. NR-1 (Rev. March 1953)

WATERFOWL (Continuation Sheet)

REFUGE FOR				1		HONT	MONTHS OF	- Commer	2 - C	Ì	19
				(2)	((3)	(7)	
•		Weeks	o f	repor	ting	peri	o d	•••	Estimated	: Production	tion
(1) Species	3 /32	**	*	3	*	1489	120	18	waterfowl days use	Broods:	:Broods:Estimated : seen : total
Swans:		Aeria 1			Aseria 1						
Trumpeter		Central			Cetterus						
Geese:											
Canada	0 4			_					33,360		
Gackling Brant											
White-fronted		,							926.05		
Snow	003	' &	*						15.5%		
Other									ş		
Ducks:											
Mallard			CA						Ż		
Gacheall											
Beldrete	-	1			1	1			4		
Pintail	8 8	35	3 5		R.S	38	38		25.4		
Green-winged teal	1,00	3	18		38	R R	38				
Blue-winged teal	}	}			}	}	}				
Cinnamon teal	8	ğ	1550		a	997	29		31.542		
Shoveler	8	897	8		8	8	R		20.08		
WOOD Dodress		,	1			1					
Reducesta Ring-necked		,	g ·			ន-	3		327		
Canvasback	2	n	*			3 0			Rti		
Seaup	2	9	8		ç	Ş	ç		1 §	-	
Goldeneye	2	}	\		}	\@	9		22		
Bufflehead	2		8		2	OI.)		2		
Ruddy	8	1500	8		887	2	R		166,630		
Other	- 1-4			pph of					,		
Pulvous 1. Duak		₩₽ ₩₽	Φ.			#	*		5		
Coot:	8	7650	99		88				166,980		
)	(0000)				•		
		_	_	-	<u>`</u>	•					

SUMMART	Principal feeding areas	Debbling-Refuge Units and agriculture leach fields. Whipses agriculture alfalfa fields	Principal nesting areas talter ter		Reported by Manre J. O'Nelll, Nefuge Manager
(7) Total Production					
(6) Peak Number		10,153	45.99	7,480 - 7,480	
(5) Total Days Use		49,135	**************************************	14,980	1.5
	Swans	ි ම සෙ	Ducks	Coots	

INSTRUCTIONS (See Secs. 7531 through 7534, Wildlife Refuges Field Manual)

In addition to the birds listed on form, other species occurring on refuge during the	reporting period should be added in appropriate spaces. Special attention should be given to those species of local and national significance.	
(1) Species:		

- (2) Weeks of Reporting Period: Estimated average refuge populations.
- Average weekly populations x number of days present for each species. Estimated Waterfowl Day's Use: \mathfrak{S}
- Estimated number of young produced based on observations and actual counts on representative Brood counts should be made on two or more areas aggregating 10% of Estimates having no basis in fact should be omitted. breeding habitat. breeding areas. Productions Ξ
- (5) Total Days Use: A summary of data recorded under (3).

9

- Maximum number of waterfowl present on refuge during any census of reporting period. Peak Number:
- (7) Total Production: A summary of data recorded under (μ) .

Interior Duplicating Section, Washington, D. C. 37944

(Nov. 1945) Form NR-1A 3-1751

MIGRATORY BIRDS

Months of (other than waterfowl)

Estimated Total Number 9 Young Total Production Number | Total # Colonies Nests (2) 100 3 Date 2 Last Seen **4** Number 2 (over) 2% iiiiiss 343445 Date 多 3 Peak Numbers (3) Number 8 88 RARARAR R~ZRBRE REE 25.2 多 Date First Seen (2) Number 8] Water and Marsh Birds Shorebirds, Gulls and balled Mon ALL SEMENATES HALLELLA OLI PATTACIO Common Name the Plone THE PARTY OF Species PARTITION OF bite Palie Mangy 1740 TA CEL 111111 TATA T Parameter S Terns: 177 II.

(1)	(2)	(3)	(4)	(5)	(9)
III. Doves and Pigeons: Mourning dove White-winged dove	Poetiene poetie				
IV. <u>Predaceous Birds:</u> Golden eagle Duck hawk Horned owl Magpie Raven Crow				·	
			Reported by	Stempt J. Office II.	

INSTRUCTIONS

Species:

(1)

I. Water and Marsh Birds (Gaviiformes to Ciconiiformes and Gruiiformes) Use the correct names as found in the A.O.U. Checklist, 1931 Edition, and list group in A.O.U. form, other species occurring on refuge during the reporting period should be added in appro-In addition to the birds listed on Special attention should be given to those species of local and National II. Shorebirds, Gulls and Terns (Charadriiformes) Avoid general terms as "seagull", "tern", etc. Groups: priate spaces. significance.

III. Doves and Pigeons (Columbiformes)

IV. Predaceous Birds (Falconiformes, Strigiformes and predaceous

Passeriformes)

- The first refuge record for the species for the season concerned. First Seen: (3)
- The greatest number of the species present in a limited interval of time. Peak Numbers: (3)
- . The last refuge record for the species during the season concerned Last Seen: (4)
- Estimated number of young produced based on observations and actual counts. Production: (2)
- (6) Total: INT.-DUP. SEC., WASH., D.C.

3-1752 Form NR-2				UPLAND GAME BIRDS	တ္ဆ		<i>√</i> , , , ,			1613
(April 1946)	Refuge galless for			Months of	s of			t C	19/2	
(1) Species	(2) Density	Pr	(3) Young Produced	(4) Sex Ratio	Re	(5) Removals	, m	(6) Total	(7) Remarks	
Common Name	Cover types, total pacreage of habitat B	Acres per Wird	broods obsivid. Estimated Total	Percentage	BuitamH	For Re- stocking	For	Estimated number using Refuge	Pertinent information not specifically requested. List introductions here.	
	S charge in starts	<u> </u>							-	
								·		
						······································				

INSTRUCTIONS

Form NR-2 - UPLAND GAME BIRDS.*

(1) SPECIES: Use correct common name.

<u>ર</u>

DENSITY:

- grass prairie, etc. Standard type symbols listed in Wildlife Management Series No. 7 should be used where possible. Figures submitted should be based on actual of cover types. information is to be prefaced by a statement from the refuge manager as to the number of acres in each cover type found on the refuge; once submitted, this information need not be repeated except as significant changes occur in the area size of sample area or areas should be indicated under Remarks. observations and counts on representative sample areas. swamp, upland hardwoods, reverting agriculture land, bottomland hardwoods, short information but not so much as to obscure the general picture. hunts, etc.). Detailed data may be omitted for species occurring in Applies particularly to those species considered in removal programs (public Density to be expressed in acres per animal by cover types. Cover types should be detailed enough to furnish the desired Survey method used and Examples: spruce limited
- (i) YOUNG PRODUCED: Estimated number of young produced, based upon observations and actual counts in representative breeding habitat.
- £ SEX RATIO other species if available. This column applies primarily to wild turkey, pheasants, etc. Include data on
- 5 REMOVALS: Indicate total number in each category removed during the report period.
- TOTAL: include resident birds plus those migrating into the refuge during certain seasons. Estimated total number using the refuge during the report period. This may
- REMARKS * include other pertinent information not specifically requested. Indicate method used to determine population and area covered in survey.

3

6

^{*} Only columns applicable to the period covered should be used.

V * 1 * 1 * 1 * 1	MAMMALO
*****	משאר

Refuge & Man Bar

3-1754 Form NR-4 (June 1945)

Year ending April 30, 1997

(5)		lurs Jestroyed		
Fure	ped	Total Ref Ture Ship		
y		share		
(μ) Disposition	Share Trapping	Trappers Share		
Á	Shar	Permit Number		
		For Re-	:	
		For Re- stocking		
(3) Removala		Predator		
Rec		Hunting Fur Harvest		
		Acres Per Animal		
(2) Deneity		Cover Types & Total	The control of the Predator Animal Hunter	وجيب مين يولين المراجعة المراجعة والمراجعة وال
(1) Species		Common Neme	Continuedall Accidentall Accidentall Accidental Acciden	

Instructions

Form NR-4 - SMALL MAMMALS (Include data on all species of importance in the management program; i. e., muskrats, beaver, coon, mink, coyote. Data on small rodents may be omitted except for estimated total population of each species considered in control operations.)

Use correct common name. Example: Striped skunk, spotted skunk, short-(Accepted common names in current use are found in the "Field Book of North American Mammals" by H. E. Anthony and the "Manual of the Vertebrate Animals of the Northeastern United States" by David Starr Jordan.) tailed weasel, gray squirrel, fox squirrel, white-tailed jackrabbit, etc.

<u>(၁</u> DENSITY:

number of acres in each cover type found on the refuge; once submitted, mitted should be based on actual observations and counts on representative the area of cover types. Cover types should be detailed enough to furnish the desired information but not so much as to obscure the general picture. this information need not be repeated except as significant changes occur in tion is to be prefaced by a statement from the refuge manager as to the Density to be expressed in acres per animal by cover types. This informa-Detailed data may be omitted for species occurring in limited numbers. Applies particularly to those species considered in removal programs. indicated under Remarks. land hardwoods, short grass prairie, etc. Standard type symbols listed in Examples: spruce swamp, upland hardwoods, reverting agriculture land, bottom sample areas. Survey method used and size of sample area or areas should be Wildlife Management Series No. 7 should be used where possible. Figures sub-

3) REMOVALS:

Hunter. Also show any removals not falling under headingslisted. previous year, including any taken on the refuge by Service Predatory Animal Indicate the total number under each category removed since April 30 of the

DISPOSITION OF JUR:

On share-trapped furs list the permit number, trapper's share, and refuge share ness or damaged condition, and furs donated to institutions or other agencies personnel. Total number of pelts of each species destroyed because of unprimeshould be shown in the column provided. Indicate the number of pelts shipped to market, including furs taken by Service

5 TOTAL POPULATION:

Estimated total population of each species reported on as of April 30.

REMARKS:

any other pertinent information not specifically requested. Indicate inventory method(s) used, size of sample area(s), introductions, and

REFUGE GRAIN REPORT

(I) VARIETTE*	(2) On Hand	(3) Received	(4)		GRAIN DI	(5) GRAIN DISPOSED OF		(6) On Hand	Propose	- 13 →	E USE*
	OF PERIOD	PERIOD	LOTAL	Transferred	Seeded	Fed	Total	End of Period	Seed	Feed	Surplus
Marious Barley	9		100		-				×		
	8		8			2	2	8		н	
	8	4	đ		4			0	н		
		9	180		8		8	3	н		
		1800	31800		160	8	9	8		×	
									-		

(8) Indicate shipping or collection points that I and II charge buildings.

Init I and II atorage buildings (9) Grain is stored at

(10) Remarks

*See instructions on back.

16--61482-1

REFUGE GRAIN REPORT

This report should cover all grain on hand, received, or disposed of, during the period covered by this narrative report.


60 lb., barley—50 lb., rye—55 lb., oats—30 lb., soy beans—60 lb., millet—50 lb., cowpeas—60 lb., and mixed—50 lb. In computing volume of granaries, multiply the cubic contents (cu. ft.) by 0.8 bushels. Report all grain in bushels. For the purpose of this report the following approximate weights of grain shall be considered equivalent to a bushel: Corn (shelled)—55 lb., corn (ear)—70 lb., wheat-

- (1) List each type of grain separately and specifically, as flint corn, yellow dent corn, square deal hybrid corn, garnet wheat, red May wheat, durum wheat, spring wheat, proso millet, combine milo, new era cowpeas, mikado soy beans, etc. Mere listing as corn, wheat, and soybeans will not suffice, as specific details are necessary in considering transfer of seed supplies to other refuges. Include only domestic grains; aquatic and other seeds will be listed on NR-9.
 - Report all grain received during period from all sources, such as transfer, share cropping, or harvest from food patches. (3)
- (4) A total of columns 2 and 3.
- (6) Column 4 less column 5.
- This is a proposed break-down by varieties of grain listed in column 6. Indicate if grain is suitable for seeding new crops. 3
- 3) Nearest railroad station for shipping and receiving.
 - (9) Where stored on refuge: "Headquarters granary," etc.
- (10) Indicate here the source of grain shipped in, destination of grain transferred, data on condition of grain, unusual uses proposed.


16-61482-1 U S. GOVERNMENT PRINTING OFFICE



Farthquake cracks in road along west edge of refuge Tract 1, Unit II. Fissures resulted in "sand boils" which brought salty water and fine sands to surface.




April 25th earthquake damage to "I" road at point 3 miles south of refuge subheadquarters.




L.E. Sinclair ranch cotton crop showing typical "sand boil" damage by quakes. Field is 2 miles south of refuge Unit II. (April 25, 1957)

Jose' Barros unloading chopped green alfalfa during widgeon feeding experiment. (Jan.1957)



Green alfalfa was spread on water. Note drifting feed in background.


Trector Operator Carl Ford points out grade fill stake where service ditch for Unit "A" will be located. Tract 1, Unit I. (March, 1957)



Mr. Ford points out same grade stake as ditch fill is being completed. (April, 1957)




Widgeon and Canada geese using green barley at Tract $l_{\rm I}$, Unit I as a result of crop manipulation efforts. Note varying degrees of utilization. (January, 1957)



Same as above showing pintails and widgeon in green barley field.



Snow goese feeding in green cattails at Tract 8, Unit II. (Photo by Dr. Howard J. Lee, Milwaukee, Wisconsin, February, 1957)



Cormorant which impaled itself on broken tree limb at bird rookery. Limb passed completely through body except for skin on back. (April, 1957)



Calipatria, California Boy Scouts who visited refuge during conservation campaign. (L. to R. Ray Coronado, George Rocha, Bob Nash, Lonnie Dearborn, Johnnie Galleano, Tom Dearborn, Dwight Metzler, Joe Epley, Charles Moe, Dan O'Neill, Robert Gibson).



Local propaganda against ban on Mourning Dove hunting brought forth heated protests.