

W. W. W. W.


SALTON SEA NATIONAL WILDLIFE REFUGE

AND

WATERFOWL DEVELOPMENT AREAS

NARRATIVE REPORT

SEPTEMBER - DECEMBER, 1951



UNITED STATES DEPARTMENT OF THE INTERIOR

FISH AND WILDLIFE SERVICE

BRAWLEY, CALIFORNIA

SALTON SEA NATIONAL WILDLIFE REFUGE

AND

WATERFOWL DEVELOPMENT AREAS

NARRATIVE REPORT

SEPTEMBER - DECEMBER, 1951

* * * * *

* * * * *

UNITED STATES DEPARTMENT OF THE INTERIOR

FISH AND WILDLIFE SERVICE

BRAWLEY, CALIFORNIA

C O N T E N T S

	Page
PERSONNEL	2
I GENERAL CONDITIONS	3
A. Weather conditions	3
B. Precipitation and Water Conditions	4
C. Fires	4
II WILDLIFE	5
A. Migratory Birds	5
1. Populations and Behavior	5
2. Geese	5
3. Ducks	6
4. Disease	9
5. Shorebirds, Gulls and Terns	10
6. Marsh and Water Birds	11
7. Food and Cover	11
B. Upland Game Birds	13
C. Other Birds	14
D. Fur Animals, Predators, etc.	14
E. Fish	15
III REFUGE DEVELOPMENT AND MAINTENANCE	15
A. Physical Developments	15
B. Plantings	16
1. Cultivated Crops	16
C. Receipts of Seed and Stock	17
IV ECONOMIC USE OF REFUGE	18
A. Grazing and Haying	18
V PUBLIC RELATIONS	18
A. Recreational Uses	18
B. Refuge Visitors	19
C. Refuge Participation	19
D. Violations	20
VI OTHER ITEMS	21
A. Items of Interest	21
VII PHOTOGRAPHS	23
VIII PROGRESS MAPS	
IX NR FORMS	

REFUGE PERSONNEL

Regular Personnel

Edward J. O'Neill	Refuge Manager
Clyde W. Stewart	Foreman-Farm Operations
Jerryl W. Saxton	Clerk-Typist
Will T. Wesley	Maintenance Man
Jose Barros	Maintenance Man
Earl M. Barker	Mechanic
Alfred W. McFarland	Tractor Operator
Melvin Ford	Tractor Operator
Carl W. Ford	Tractor Operator
Chesley H. Williams	Tractor Operator
James W. Hamilton	Dragline Operator

Temporary Personnel

John Barros	Irrigator
Sylvester Barros	Irrigator
Ray W. Bennett	Tractor Operator
Raymond L. Gash	Carpenter
William E. Hoff	Laborer
Leon Lesicka	Laborer
Manuel Cardonzo	Irrigator
William L. Lynch	Irrigator
Owen Schutt	Irrigator
Leo E. Cox	Tractor Operator
Paul E. Williams	Tractor Operator

NARRATIVE REPORT

I GENERAL CONDITIONS

A. Weather Conditions

Weather throughout the period was rather pleasant save for a few hot, humid days in September and a number of windy days in December.

Light drizzly rains fell the night of November 22nd, wetting roads and fields badly with the small amount of precipitation received.

On December 13th an unusual, dense fog, blanketed the Valley with heaviest patches occurring near Calxico. Members of the weather station blamed the freak occurrence on a moist air mass in the air.

The temperature dropped the lowest since last December, when on the morning of the 15th same nature emerged sporting a most beautiful sparkling, frosty white coat. In parts of the Valley the mercury dipped as low as 26°. Anti-freeze sales went up as the temperatures went down, and it was not until well into January that tourists were fully aware that this was truly sunny California.

Tabulated below is the weather data for the period as compiled by the El Centro Naval Air Station at El Centro, Calif.

<u>MONTH</u>	<u>MAXIMUM</u>	<u>MINIMUM</u>	<u>PRECIPITATION</u>	<u>DAYS WINDS OVER 25 MPH</u>
September	111	67	tr	3
October	113	54	tr	3
November	87	41	.21	5
December	78	33	<u>.08</u>	10
TOTAL29	

Early on the morning of December 5th an earthquake of considerable force rocked the Imperial Valley and tore a large 100 foot long crack in the Standard and Highline Canal roads and damaged half a dozen homes. The quake shook with a number two intensity and one woman suffered a bruised hip when thrown against

a toppling chest of drawers. Dishes crashed to the floor and furniture shifted back and forth in scores of houses. A District hydrographer reported cracks in the road near the Highline Canal for a distance of 2 or 3 miles. The Mulberry, or east portion of the Valley, appeared to be the center of the shock, which was said to be worse than the 1940 quake.

B. Precipitation and Water Conditions

The rains of the previous period which resulted in cloud burst proportioned run-off continued to hamper operations well into September. After the clouds cleared, all of tracts 5, 6, 12 and 13 in Unit I were reflooded with nearly 1 foot of muddy, yellow water, which refused to drain from the area due to the rise in Salton Sea. Never during the period did the waters recede enough to again expose Tracts 12 and 13, as irrigation water from the balance of Unit I later added to the natural sump area. The dragline operations there were discontinued and may be resumed in the future, if the waters recede.

The shoreline and elevation of Salton Sea dropped very little throughout the summer months and at the close of the period it was evident that a gradual rise was taking place much earlier than in former, similar years, of low precipitation. Some place the blame on the late August rain and others pointed out the tremendous irrigation spillage from the Valley's new cotton crop.

From here it appears as though a combination of irrigation waste water drainage, rains and lack of usual (6 ft. per year) evaporation are all to blame. If cotton is important as a waste water crop it could be that next year will see more of both than ever. Cotton yields have been almost fantastic. Poor, secondary lands have brought off one and one-half bale to the acre. Better lands, inland from the Sea are yielding their 3rd and 4th bale! We can expect to see lots of Cadillac pickups and some broad smiles before the last bale is ginned. One 80 acre parcel of land next to tract 18, Unit I, was purchased a few years back for \$20.00 per acre. This year the land sold for \$100.00 per acre and after the 2nd cotton picking the new owner was sporting a clear title to the land.

C. Fires

Each year the cattails make tremendous growths around fresh water canal and drain estuaries. For years it has been the practice to burn the growth, partly to eliminate black bird roosts near grain crops, but mainly to open areas for operation by the Irrigation District. One such fire just west of Unit I, along the SW shores of Salton sea aroused many a new citizen with the spectacular black, billowing smoke clouds. One bus driver reported

the Atomic Energy Commission was going up in flames. A local newspaper reporter spent most of the night tramping over the jungle-like area to learn that the rumors were false and some farmer was just burning cattail patches.

* * * * *

II WILDLIFE

A. Migratory Birds

1. Populations and Behavior

The increased population figures shown on form 3-1750 are accountable somewhat through general expansion of the refuge units. It could be stated that a true picture of population trends is not reflected as is the overall refuge "use" factor.

2. Geese

As early as the torrid days of September, geese were here. On the 1st, three Lesser Snows were present. These we suspected to be hold-over cripples and did not look upon their presence as a natural occurrence. They were all capable of extended flights as near as could be determined. Main flights started to arrive early in October. The peak came the week of December 9th, when an estimated 8,600 were present.

On September 29th, 17 White-fronted geese showed up. October 6th, 70 were here and on October 13th an estimated 1,000. After that the arrival was steady and by October 27th 1,300 were here. The species reached an all time peak of 1,800 the week of November 3rd.

Two Canada geese were here October 6th. Our notes at that date read "Weather still in upper 80's, prospective hunters arriving in flower-patterned shirts and straw hats." On October 13th, 15 Canada's were present and 22 on October 21st. The increase continued gradual, by November 18th 900 were here, November 30th, 1,580. On December 22nd an estimate of 2,450 was made. This represents an all time high. About that time one visitor from Montana related how the geese in northern states literally passed over the cold, frozen northern marshes.

Returns on banded Canada geese which were killed near the refuge indicated they were banded by Biologist Horton Jensen at Blackfoot Reservoir, Idaho, and by personnel at the Bear River Refuge, Utah.

Three Lesser Canadas were examined closely during the period. One, a cripple, remained for some time in the headquarters enclosure, disappeared and returned a week or so later. Others were checked in the bags of successful hunters within a stones throw of the Unit I boundary.

December 3rd 8 Ross' Geese showed up at Unit I. Two individuals were observed as close as 100 yards. Throughout the balance of the period from 1 to 3 were seen often with Snow and Canada geese in refuge alfalfa plots. Last year Agent Wooten and State Warden Guy Noel reported a picked "small white geese" coming through Customs at the International Boundary. Two years ago Agent A. W. Elder received a report that 2 small geese, believed to be Ross' geese, were taken here by a Los Angeles hunter. Three years ago we learned of three young men who bagged two or three small white geese in the desert west of Imperial. (See pp 5, Salton Sea narrative - September - December, 1950) Although literature does not record the Ross' goose as ranging this far south, it would appear the species does and has done so for a number of years.

The Cackling goose, which regularly winters here in small numbers, is not recorded in literature either. This winter only one was seen on the refuge with Canada geese.

Of interest is the two unusual Canada geese which wintered on the refuge. Both birds were partial albinos - one with a white head and neck but normally marked and colored body. The other individual appeared just the opposite - normal head and neck with very light wings and body plumage. Surprisingly, these birds appeared to be mated with normal geese. At all times both were very wary and difficult to approach. An amateur photographer, Mr. Mikhardt, with the aid of telephoto lens, photographed both of the birds. One report indicated the white-necked bird was seen at the State's Honey Lake Refuge in the fall. Another report came in that the same bird was seen last winter in lower Mexico, during the annual waterfowl inventory. Word reached us late in the season that a number of ninruds in the country had either seen or heard of these freaks of nature. A \$10.00 reward was being offered for the person killing the white headed bird.

3. Ducks

By September the fall movements of ducks into the Salton Sea country were well started also.

Most prominent the first part of the month was the arrival of some 1,200 Cinnamon Teal and 4,500 Pintails. Fulvous Tree ducks reached a count of 460. The Green-winged Teal also showed a population gain with the arrival of 300. Some 160 American Coots were present at the same time. Total waterfowl

population then was estimated to be 7,150 birds.

The following few days in early September saw Cinnamon double in numbers with 2,100 present. First Shovellers of the season arrived when we recorded 2 individuals. Baldpates also were now on the fall list, with 10 present. Fulvous ducks reached a peak of 530. Most outstanding were the Pintails with a population of 8,200. By this time the waterfowl population on the refuge was estimated to be 11,850 birds.

The last of September found a drop in Cinnamons to 400; Shovellers went up to 230; 900 Ruddy ducks; 180 Baldpates; 2,080 Green-winged Teal and 1,500 American Coots. Some 6 Mallards were present and Fulvous ducks dropped to 200.

In mid-October the records read thus: Cinnamon Teal, 500; Shovellers, 200; Ruddy ducks, 500; Baldpates, 190; Green-winged Teal, 400; Mallards, 300 and Coots were up to 2,000. Pintails were steadiest of all species with 10,030 present. Fulvous ducks showed steady decline after this. At this time a total waterfowl population of about 14,000 was present.

Mid-November brought the greatest peak ever to the present refuge setup, an estimated 134,000 ducks and geese present. Cinnamon Teal reached 2,300; Mallards, 300; Gadwalls, 150; Baldpates, 45,000; Pintails, 80,000 and Green-winged Teal 1,200. Coots dropped to about 800.

State Fish and Game Biologist, Mr. William Anderson, banded some 3,500 birds in September and early October on the refuge units. Dry condition of the State units and hunting activities eliminated trapping operations on other than the refuge units. Throughout the season the Federal area accommodated the bulk of the birds as usual.

During the period the Service's L-5 Stinson plane, No. N-720, made two trips into the area. Pilot-Biologist Ross Hanson and the writer continued previous waterfowl census work and made progress photographs of various sections of the refuge area. A survey flight was likewise made of the Colorado River from Blythe to Yuma, to learn more about local distribution of wintering birds.

Tabulated below is a summary of waterfowl populations observed in Imperial Valley during Aerial Surveys;

November 13th and 14th flights: (Late PM and early AM)

<u>STATE IMPERIAL REFUGE</u>	<u>NUMBERS</u>
Baldpates	3000
Pintails	3000
Coots	500
	<u>6500</u>

STATE WATERFOWL MANAGEMENT AREAS NUMBERS

(Hazard and Unit B)

Pintails	350
Canada Geese	20
Snow Geese	80
	<u>450</u>

SALTON SEA REFUGE

Units I and II)

Canada Geese	800
Lesser Snow Geese	3400
Baldpates	30000
Pintails	40000
Teal	1000
Shovellers	200
Ruddy	500
Coot	500
	<u>76400</u>

SALTON SEA AND ADJACENT VALLEY AREAS

Canada Geese	40
Baldpates	14400
Pintails	200
Ruddy	9300
Buffle heads	170
Lesser Scaup	2600
Canvasbacks and Redheads	580
	<u>25290</u>
	27290

ESTIMATED TOTAL 108,640
110,

December 22nd flights: (Early AM and late PM)

STATE IMPERIAL REFUGE NUMBERS

Baldpates	1500
Coot	500
	<u>2000</u>

STATE WATERFOWL MANAGEMENT AREAS

(Hazard and Unit B)

Baldpates	2300 ✓
Canada Geese	30 ✓
White-fronted Geese	20
Unidentified Ducks	200
	<u>2550</u>

<u>SALTON SEA REFUGE</u>	<u>NUMBERS</u>
(Units I and II)	
Canada Geese	2000 ✓
White-fronted Geese	500
Snow Geese	5000
Baldpates	2000
Pintails	3000
Teal	2000
Shovellers	1500
Coots	<u>300</u>
	34300

<u>SALTON SEA AND ADJACENT VALLEY AREAS</u>	
Canada Geese	200
Snow Geese	2500
Baldpates and Pintails	40000
Other and unidentified ducks	<u>4300</u>
	47000

ESTIMATED TOTAL 81,300

The matter of disturbance of Snow geese by airplanes has been mentioned in previous reports as an important harrassment factor. The following was extracted from the latest (1950-1951) report of the Severn Wildfowl Trust, Covent Garden, London, England by Curator Peter Scott:

" the disturbance of the geese by aircraft was nevertheless very great. For some reason wild geese and particularly White-fronted geese, (Barnacle geese behave in the same manner) do not become used to aeroplanes. They take wing at once in great alarm at the approach of any aircraft which is heading toward them at a height of less than 5,000 feet and within a mile. This is particularly noticable in the early part of the season when the flock is not very large. The geese appear to develop a kind of neurosis and fly out into the mudflats where they fancy themselves more secure. So serious did this disturbance become during November and December that the geese were kept continually off their feeding grounds by day and became largely nocturnal."

4. Disease

What appears to have been botulism persisted at Salton Sea during the entire period. First indication of sickness showed up in Pintails and Ruddy ducks during the second week in September. It is believed that an equal number of shorebirds succumbed to the condition, but our observations disclosed about 300 ducks died.

Green-winged Teal from the State units were examined by Dr. E. R. Quortrup, San Diego County Veterinarian (formerly of this Service). Merton Rosen of the State, later examined a Pintail from Unit I of the refuge. Both men pronounced the sickness as being botulism. No formal laboratory tests were conducted.

Coincidental to the local sickness, a report reached Warden George Werden, of sick and dead ducks in the Blythe, California area. Mr. William Anderson of the State investigated and discovered a small "dry" lake, known as Ford Lake, northeast of Blythe. The lake was first filled by heavy rains of the last period, about August 28th. Later, about October 29th, rains again brought the lake up. Anderson found that due to the inaccessibility of the area and muddy shoreline conditions, which prevented driving closer than $\frac{1}{2}$ mile, rescue work was impossible. In October very few sick birds were found. It was estimated that about 100 recently lost birds were there. Some birds appeared to have been dead for several weeks, few were beyond recognition. Dead birds included Pintails, Baldpates, Green-winged Teal, Shovellers and Avocets, in order of abundance.

Creosote bush (*Colvillea*) was the main submerged vegetation but *Baccharis* was also dense. The second storm submerged new vegetation which resulted from the August rains. All told about 500 ducks were lost. The Naval Air Base was contacted and in turn pilots drove off the remaining ducks with airplanes and smoke bombs, to prevent further use and contamination of the waters. Little is known of the numerous playa lakes scattered over the local inaccessible desert areas, but they could contribute considerable in the way of duck losses if Ford Lake is a typical example.

5. Shorebirds, Gulls and Terns

Mid-September saw a good variety of mud-probing shorebirds at Salton Sea. Wilson's Phalaropes, Glossy Ibis, Avocets, Western Sandpipers, Black-necked Stilts, Forrester's Terns, Hudsonian and Long-billed Curlews, Least Sandpipers and Yellowlegs all drew the attention of visiting amateur naturalists and photographers.

The usual Ring-billed Gull population was present and on September 29th we estimated that about 2,000 were present on the refuge units. A few Herring and Glaucous winged gulls were present also.

By the last part of the period the picture was changed. Avocets, Black-necked Stilts and yellowlegs were reduced to a small handful. Hudsonian Curlews were absent as well as Phalaropes, terns and the other more tender little denizens of Salton Sea's fringed shorelines. In their places are found Mountain Plover grouped together in hundreds, Sprague's Pipits and Long-billed Dowitchers.

AMERICAN

From 300 to 400 Black-bellied Plover spent the winter here, ranging far and wide over the refuge and neighboring irrigated farm fields.

Forester's and Black terns, as well as a rare straggler of a Gull-billed tern, were here in September but soon departed except for a few Foresters which either winter here or infrequently revisit the Valley during the winter season.

6. Marsh and Water Birds

Some 60 Sandhill Cranes were reported in December by Mr. Wesley, of the Refuge. They frequented the Mesquite Lake area east of Imperial. Only 5 were seen on the refuge in late December.

On September 30th, Agents Elder and Wooten called our attention to 5 Rosetta Spoonbills which they had just discovered on the mouth of the Alamo River. Upon investigation we managed to find 2 more which brought the total up to 7. These birds were seen repeatedly in September but appeared gone after the first part of October. Several years ago Mr. Ben Skupen, operator of the Alamo River Duck Preserve Club, reported seeing a number of small Flamingos. We wonder now if perhaps the Spoonbills were not the subject of his observations. In June of 1951, a "glimpse" observation of one individual was reported in the Valley. (See May-August, 1951 Salton Sea Narrative, pp 3). Of the Spoonbills seen, we concluded all were immature birds. Undoubtedly they moved north to Salton Sea from the nesting grounds in Mexico, as do the Wood Ibis, Egrets and Pelicans. The species makes an interesting addition to our list of species. There appears to be little difference in the habitat here and in the country they inhabit in Mexico.

In early September about 1,600 Wood Ibis were here. By mid-October there was only a few cripples to be found.

White-faced Glossy Ibis were very low in numbers as compared with former years.

7. Food and Cover

As evidenced by the tabulations on page 16 under Cultivated Crops, the overall production of foods for wild ducks and geese continued to show progress. Birds are becoming more and more aware of the refuge foods and the seclusion from harassment. This fall thousands of Pintails were present and continually using the refuge, almost entirely without the knowledge of the farmers, which they have always plagued at seeding time in the early fall.

In order to achieve better utilization of Unit I dry grain crops, the impoundment pond in Tract 11 was pushed to the fullest extent of available equipment through the early days of the period. Even before completion of dragline and bulldozer work on the upper perimeter, water was running to fill up the lower half of the pond. The fresh water adjacent to grain fields had an almost immediate effect. Ducks, especially Pintails, flocked in by the hundreds.

Both ducks and geese utilized the large, unbroken Tract 4 of Unit I, first in preference to the much better stand of barley on the smaller, west half of Tract 19, which also was near a new (Tract 29-31) fresh water impoundment area. White-fronted geese continued to utilize the Tract 4 field until November 19th, at which time it appeared as though a giant steam roller had passed over the field. Geese use a remarkable technique we had never noted before. In taking the grains where the growth was rather tall, both the Snow and White-fronted geese were seen walking astride individual clumps and actually "walking them down" until they were within easy reach. This may account for the completely flattened appearance some of the dry barley fields had after utilization.

The impoundment area, (Tracts 29-31) in the north end of Unit I produced a good stand of mixed Wild Millet and Sudan-grass in the upper contours. The lower contours produced very little of the emergents but the production of Widgeon grass (*Ruppia*) was surprising. This entire area, planted to bullrush (*Scirpus p.*) rhizomes in the previous period, produced very little of the species. At first growth was good, some plants spread as much as 12 feet in 3 months time, but with them came the ever present cattails which choked out most of the growth except where bullrush escaped competition by retreating, in a few places, up the slopes of contours and dikes.

At Red Hill in Unit II, Wild Millet was likewise planted in the upper contours where it produced well. Later it was dominated by new cattail growth.

Wild Millet, although not highly satisfactory here as a marsh food plant, did prove attractive in early September when taken by Pintails and Fulvous ducks. West of Tract 11, Unit I, a small patch of Wild Millet volunteered mixed with Sea Purslane (*Sesuvium g.*). It appeared green and appetizing to Snow geese which moved in during December and utilized both green seed heads and stems of millet and about 20% of the terminal growth of Sea Purslane.

On several occasions we observed small groups of Baldpates in the refuge dry barley fields. It is possible that were

an adequate supply available, the species might prefer it to the green alfalfa crops. In December, several sacks of feed barley were dumped along one dike to "ohum" Pintails for trapping operations. Later we observed the unpredictable Baldpate gleaning the entire offering. During the hunting season Mr. Walter Collins, farmer near Westmorland, reported killing a Baldpate which was "full of barley grain."

During the period 684 bushels of field run barley, shipped in from Tule Lake Refuge, were scattered along the units for supplementary food. Feeding started October 31st and continued throughout the period. Snow geese reacted favorably, but were mainly interested in the green alfalfa crops on and off the refuge.

Again the Snow and White-fronted geese cut down and utilized basal stem and root portions of many acres of cattails (Typha d). It was conservatively estimated that on the refuge units alone at least 200 acres of green, current years growth was cut down. Both Pintails and Baldpates joined the geese and may have benefited a great deal as rootlets and stem debris became available. Some weights of green samples in a measured quadrat indicated that the 200 acres of growth, of which we estimated about 60% utilization of individual plants, amounted to about 1500 lbs. of green feed per acre. Surprising enough, the total for the entire acreage would amount to an estimated 300,000 pounds or 150 tons of green starchy food!

At times the cattail stems were strewn so thick on the water surface that small, peep sandpipers could walk from contour to contour across some of the checks.

Marsh and shorebird foods appeared adequate in irrigated fields, along the ever expanding shores of Salton Sea and on the refuge impoundment units. Where Widgeon grass and Horned pondweed flourished in the fresh water units and where drains and canals meet the Sea, insect life seems always abundant. There is always a great variety of water temperatures, salinity, etc., to supply almost any condition desired.

B. Upland Game Birds

Gamble Quail held up rather well in population under the pressure of this season's hunting, which ran from November 17th to 26th with a daily limit of 8. Cripples were most common than after any previous season we'd witnessed.

Pheasants enjoyed a fair season in the field as best we could determine from observations. The State Fish and Game again released several thousand hatchery-reared birds which "went like Coney Island Red Hots" as an army of hundreds of mis-guided

nimrods invaded the Valley from November 17th to November 26th, for their 2 birds per day bag. The seasonal limit was 10, each to be tagged; 5 hens were permissible in the bag. We saw but a small number of pheasants bagged, but from the sounds of shotguns out our way it can be said that Mr. Du Pont's cash register was in operation as Roadrunners, Hawks and Owls became fewer and fewer along county roads.

Worst of all was the manner in which the State Commission decided, just before opening day, that hunting could be started at sunrise instead of the originally announced 8:00 AM. Our faces were crimson when an apprehended early shooter showed us the morning Los Angeles newspaper with the season change write up!

Following the late August floods of the previous period, most of the Screwbean and Mesquite bean seeds lying under trees were washed away or covered by sand. Many birds, especially quail, shifted to higher ground or the desert edge where other plants thrived as a compensating factor for lost habitat.

C. Other Birds

Robins returned in the fall and spent the winter here again. Several flocks ranging from 50 to about 300 were seen often. Choice item of food appeared to be dates in the small groves, where most of the birds stayed. In Los Angeles, Kenneth Stager, of the County Museum, and Dr. John Doris, Occidental College, invited the public to observe specific instances of robins nesting and congregating. This year appears to be the first time in recorded history at the metropolis, the cheerful red breast showed up in spring and summertime to nest there. Theories were flying as to sudden appearance of robins. Some said it was the increased humidity; others the green lawn sprinklers and resultant earthworms.

D. Fur Animals, Predators, etc.

No notable change over previous period.

It is interesting to note that the last cotton raised in these parts was almost 15 years ago. White-footed mice however, like the idea and refuge personnel, Jose and Sylvester Barros, found a nest of fine, clean cotton which appears to have been brought in from a field almost $\frac{1}{2}$ mile away. The fiber was as clean as the product from some recleaning plant.

E. Fish

The usual fall spawning run of Mullet from Salton Sea up the New and Alamo Rivers was not so pronounced as in previous years. Fishermen with snag hooks and legal, 6 foot dip nets accounted for only a small number during October and early November.

Three years ago the State Fish and Game Department transported a tank load of Anchovies, Sea Bass, Halibut and other ocean species from San Felipe, Mexico, to Salton Sea. This winter a commercial fisherman drawing up his Mullet nets took a Sea Bass. This represents the first report or indication that any of the original stock still survived.

In Units I and II fresh water impoundment areas small fish are already established. Cormorants and Pelicans spent considerable time in the upper contours next to the irrigation canal inlet and our guess is that when they are drained for cultivation, control of cattails and reseeding, some whoppers will show up in the deeper pools.

* * * * *

III REFUGE DEVELOPMENT AND MAINTENANCE

A. Physical Developments

Early in the period the levee which encloses Tract 11 of Unit I was completed through the combined efforts of D-6 and D-7 bulldozers and the P&H $\frac{1}{2}$ -yard dragline. This was one of the most outstanding accomplishments during the period. Water at this location completed the needs for hungry waterfowl and crop utilization was almost immediately doubled by Pintails.

Two miles of deep, 6 foot drain ditch and $2\frac{1}{2}$ miles of irrigation ditch were constructed at Unit I with the dragline before it was placed under repairs.

At headquarters a 30 x 30 foot enclosure with a water pond and shelter was constructed for retrieved, crippled ducks and geese. Before the end of the period, 9 geese and several ducks had been placed in the enclosure where visitors could view and photograph them at close range. About 3 of the stronger geese and most of the ducks, which were banded, managed to recuperate and fly from the enclosure.

Tract 8 of Unit I was cleared of contours and rank growths of Salt Cedar by grader, releveled by landplane and seeded to barley.

Tracts 4, 5, 7, 8, 24 and 25 of Unit I were deep chiseled, disked, landplaned and planted during the period.

In Unit II developments were continued. Tract 17 was contoured by bulldozer and 18 inch concrete drop box structures and pipe were installed for water circulation.

Tracts 2, 9 and 12 of Unit II were deep tilled, landplaned, etc. and planted.

Construction of an oil house and work benches in the Service building at sub-headquarters were continued.

At headquarters a concrete block fence was erected around the office building and lawn.

The F&H Dragline, transferred in from the Wheeler Refuge was sandblasted in early December, preparatory to painting.

The Autocar truck-trailer electric brakes were repaired and a new floor installed on the trailer. General maintenance work included new seals on the GMC Cargo truck wheels, conversion of same to an oil-grease unit for field use, overhaul of the Ford Jeep motor, etc.

B. Plantings

1. Cultivated Crops

Unit I:

Acres under lease	14600
Acres under fallow	320
Acres in crops	1880
Acres under sump, river, etc. .	11400

The fresh water impoundment area (Tract 28-30), although flooded prior to the period, was seeded with additional, soaked Wild Millet to produce as much food as possible and to determine the feasibility of late plantings. The undertaking netted very little in this instance and seed was either taken by ducks or washed ashore by persistent winds.

Tract 5 of Unit I, seeded to certified Mariout Barley the first week of October produced well, however the utilization by geese, between, during and after each irrigation meant complete loss of the 160 acre tract before the end of the period. We have never seen such heavy, complete green crop use heretofore.

Scarcely a standing stock of grain could be found in Tract 4. Lots of people saw the field being used each morning and evening. Farmers were delighted; - - pseudo sportsmen,

drooling over the boundary signs, muttered something about "injustice", or a trip to the Justice, or something.

Unit II:

Acres under lease	2180
Acres under fallow	320
Acres under crops	880
Acres under swamp, river, etc. .	900

The fresh water impoundment area around Red Hill in Tract 17 was given a light reseeding in spots to continue the experiment with Wild Millet to determine the best season for growth and seed production. The late planting resulted in fair growth gains through September and early October. Seed production was poor.

In mid-October the lower ends of lands and bare spots in the Tract 1-2 Sudangrass field were worked and seeded to California Common Barley. The results were good and geese grazed the crop throughout the entire period.

Three acres of buckwheat, seeded in October in Tract 9, at first progressed well, but shortly before blooming, late in the period, the experimental crop was killed by frost. Recent trials by the local experiment station indicate this species is unsuccessful in our desert climate.

During the September meeting of the Lea Act Committee group, it was decided that hunting would not be recommended for any of Unit II Lea Act acquired lands. Tracts 15 and 22 which were flooded and being held in anticipation of a hunting program, were promptly drained, contours plowed down and the area was tilled and leveled for future cropping. The dense growth of cattails which developed during leaching proved an outstanding food item for all species of geese. At times it was estimated between 6,000 and 7,000 Canadas and Snows were using the field as two and sometimes three tractors worked the land. The basal portion of each available plant stock, regardless of the dehydrated, very hard condition, was completely utilized.

C. Receipts of Seed and Stock

Early in October a carload of barley for supplementary feeding and duck trap baiting was received from the Tule Lake Refuge in accordance with past procedures.

* * * * *

IV ECONOMIC USE OF REFUGE

A. Grazing and Haying

The alfalfa seed crop, anticipated during the previous period, was badly damaged during late August rains and what formerly was a crop in great demand, turned out to be a real problem. The matured crop was well beyond the stage of even fair hay. Farmers called for cattle and sheep to remove the worst crops, but many tons were baled and sold to coastal citrus growers for fertilizer and ground mulching purposes. After several unsuccessful attempts we finally obtained a band of 1,460 head of sheep, owned by Mr. Arnold Shields, which entered Tracts 18-19, Unit I, on October 15th. The crop, despite somewhat dried, defoliated condition, held the grazing animals until October 29th (15 days).

Tracts 3 and 4, Unit II, were in much better condition and Permittee Newton Ruston, neighboring farmer, again harvested both fields as No. 2 hay. During the entire summer-fall season Ruston harvested 138 tons of alfalfa hay.

Mr. Lewis C. Copeland, tractor operator with the California Fish and Game, purchased the rain-damaged crop in Tract 27 of Unit I, which netted him 20 tons. His bid for the hay was the only one received.

* * * * *

V PUBLIC RELATIONS

A. Recreational Uses

With the close of the regular waterfowl season it was announced in local papers that the refuge was opened to tour. Interested individuals and groups met and departed from headquarters at 1:30 PM every Sunday. Many citizens were guided over each unit to view the birds using refuge alfalfa and barley crops and the marsh area. Refuge Mechanic Barker contributed many hours of his time in assisting with refuge visitors. A total of 167 people visited the units during conducted tours.

Some 180 members of the Orange County, San Bernardino County and San Diego County Bird Clubs and University of California students studied birds on the refuge units during the period.

1. <u>Public Use</u>	<u>Visitor Days</u>
a. Hunting Use	None
b. Fishing Use	120
c. Miscellaneous Use	<u>387</u>
Total	507

B. Refuge Visitors

<u>NAME</u>	<u>DATE</u>	<u>IDENTIFICATION</u>	<u>PURPOSE</u>
Wm. Anderson (many visits)	9/2	Calif. F & G	Duck trapping and banding
B. Loveland	9/26	Lea Act Committee	Refuge tour
W. R. Batley	9/26	Lea Act Committee	Refuge tour
C. B. Miller	9/26	Lea Act Committee	Refuge tour
R. Miller	9/26	Lea Act Committee	Refuge tour
M. Ferguson	9/26	Lea Act Committee	Refuge tour
O. Witcher	9/26	Lea Act Committee	Refuge tour
R. Jefferson	9/26	Lea Act Committee	Refuge tour
Leo L. Laythe	9/26	Regional Director	Refuge tour
K. F. MacDonald	9/26	Refuge Supervisor	Refuge tour
Fred Ross	9/26	CF&G La Quinta	Refuge tour
D. Tillotson	9/26	CF&G San Francisco	Refuge tour
Larry Rubke	9/26	CF&G Calipatria	Refuge tour
F. M. Francis	10/13	U. S. Senate	Bird observations
R. Frohling	10/16	USN San Diego	Bird observations
R. Ryder	10/16	USN San Diego	Bird observations
B. McEachern	11/5-26	USF&W Merced	Clerical detail
R. Hanson	11/13	USF&W Sacramento	Aerial surveys,
	12/22	Pilot-Biologist	etc.
C. Leichhardt	12/28	G. M. Supervisor	Waterfowl inventory
Guy Noel	12/28	CF&G Warden	Waterfowl inventory

C. Refuge Participation

The Supervisor of county schools requested participation in the local education program for schools by way of short talks on wildlife conservation. We responded by giving a talk, answering questions and showing the films "Conservation in Action" and "Haunts for the Hunted". From October 17th through October 19th, 5 county schools were visited and 1,540 children of the lower grades fired their surprising lists of questions, listened to the refuge function story and thoroughly enjoyed our wildlife films.

On October 18th some 860 persons comprising personnel and families of the Sandea Corporation, AEC Salton Sea Base, saw the films "Halibut Fishing" and "Conservation in Action".

Personnel, including the Refuge Manager and Jose Barros, participated in a law enforcement meeting of Services, California and Arizona officers at Yuma, Arizona, on November 4th.

During the annual dove hunting season Jose Barros and

Jerryl Sexton participated in patrol and hunters bag inspection as in the past.

D. Violations

The period marks, perhaps, the most successful season of refuge patrol in the history of the refuge. Personnel effected some 45 apprehensions, 30 of which are listed below.

REP'T NO.	NAME & ADD.	VIOLATION	FINE
163	H. Van Diest Paramount, Cal.	Possession of firearms on Federal refuge	\$35.00
164	Earl Croft Paramount, Cal.	Possession of firearms on Federal refuge	35.00
165	Harley Brown Calipatria, Cal.	Hunting after hours on Federal refuge	35.00
166	Rex R. Hope Pomona, Cal.	Discharge of firearms on Federal refuge	25.00
167	R. R. Rousseau Claremont, Cal.	Discharge of firearms on Federal refuge	25.00
168	P. Nordisini Brawley, Cal.	Possession of firearms on Federal refuge	25.00
169	Bob Higgins Brawley, Cal.	Possession of firearms on Federal refuge	25.00
170	Wm. Dube El Centro, Cal.	Possession of firearms on Federal refuge	25.00
171	F. Martin Imperial, Cal.	Possession of firearms on Federal refuge	25.00
172	J. C. Potter Hermosa Beach, Cal.	Shooting after hours on Federal refuge	50.00
173	J. R. Jones Long Beach, Cal.	Taking Glossy Ibis	25.00
180	A. K. Johnson Redondo Beach, Cal.	Hunting after hours on Federal refuge	50.00
181	L. G. Korich Venice, Cal.	Hunting after hours on Federal refuge	50.00
182	R. A. McClelland Venice, Cal.	Hunting after hours on Federal refuge	50.00
183	R. D. Potter Los Angeles, Cal.	Hunting after hours on Federal refuge	50.00
184	L. Johnston Costa Mesa, Cal.	Possession of firearms on Federal refuge	25.00

REP'T

<u>NO.</u>	<u>NAME & ADD.</u>	<u>VIOLATION</u>	<u>FINE</u>
185	O. Deerham Orange, Cal.	Possession of firearms on Federal refuge	\$25.00
186	Joe Lorente Los Angeles, Cal.	Possession of firearms on Federal refuge	25.00
187	C. W. Poole Long Beach, Cal.	Possession of firearms on Federal refuge	25.00
188	V. E. Lind San Diego, Cal.	Possession of firearms on Federal refuge	25.00
189	Don Denham San Bernardino, Cal.	Possession of firearms on Federal refuge	25.00
190	R. H. Arnold Los Angeles, Cal.	Possession of firearms on Federal refuge	25.00
191	J. Facinelli Los Angeles, Cal.	Possession of firearms on Federal refuge	25.00
192	M. Handcock La Mesa, Cal.	Possession of firearms on Federal refuge	35.00
193	Leo Austin El Centro, Cal.	Possession of firearms on Federal refuge	25.00
194	H. B. Turner El Centro, Cal.	Possession of firearms on Federal refuge	25.00
195	L. B. Pearson San Diego, Cal.	Possession of firearms on Federal refuge	35.00
196	M. E. Cox El Cajon, Calif.	Possession of firearms on Federal refuge	35.00
197	Ronald Rince Newport Beach, Cal.	Failure to show license on demand	10.00
198	D. J. Steele Pasadena, Cal.	Possession of firearms on Federal refuge	25.00
199	H. J. Forrester Compton, Cal.	Hunting after hours	35.00

* * * * *

VI OTHER ITEMS

A. Items of Interest

The State shooting grounds adjacent to the Federal units I and II were the subject of an article in the September issue of the Sports Afield magazine. The article, entitled "Poor Man's Million-Dollar Duck Club", was written by Lupi Saldana, Los Angeles Daily News columnist. Subject matter described a hunting trip taken by the author and companions in the State-managed shooting grounds.

"WORD FROM WASHINGTON INDICATES STATE HUNTING GROUNDS AROUND SALTON SEA AGAINST FEDERAL LAW" - - that was the headlines

of an article appearing in the local Imperial Valley Democrat, Brawley, California, December 20, 1951. The article went on - "The Fish and Wildlife Service (should have been California Department of Fish and Game) went at the hunting game in a thorough manner for this season. Many thousands of dollars were spent in planting wild rice and barley surrounded by borders and then making a pool alongside the borders with blinds at suitable distances. The effort was successful and when the ducks and geese came to eat the grain they were slaughtered by the hunters.

Private citizens have not been allowed to do anything like this under threat of prosecution, but a recent report from Washington states that henceforth the Federal law will be enforced and such practice prevented in the future."

A total of 266 blinds were offered to hunters using the State Fish and Game shooting grounds - all for the sum total of \$5.00 per man, 2 men per blind. On the Hazard, 60 blinds were erected and on the Pumice (Unit B) 206. The Poe area (Unit A) remained undeveloped and was offered for free. Shooting took place on all areas on opening and closing days, and on Saturdays, Sundays, Wednesdays and holidays.

No dogs were permitted as usual on the \$5.00 areas and hunters were encouraged to bring hip boots and flashlights.

The old registration place at Heise's Station on highway 99 was changed to the Hazard area. Minors up to 16 years of age were admitted for \$2.50 each.

The old Imperial "Refuge" area near Calipatria remained open to hunting daily as it has been for several years.

An immature white-winged Scoter, shot by Mr. Otto Witcher at the south end of the Salton Sea, during the early part of the hunting season was obtained and turned over to the Los Angeles County museum for a specimen.

Members of a Hollywood motion picture firm, Wild Life Films, currently producing material for a weekly television program entitled "Our Great Out Doors", toured parts of Unit II in October to survey the possibilities of a film on refuge programs and the various forms of wildlife.


A tour of Federal and State Units, followed by a meeting of the Lea Act advisory Committee, took place September 26th. Refuge personnel recorded all meeting minutes and handled

secretarial work for the committee.

Respectfully submitted,



Edward J. O'Neill
Refuge Manager

APPROVED:



SALTON

SEA



PROGRESS MAP
September - December, 1951

SALTON SEA NATIONAL WILDLIFE REFUGE

UNIT II

SCALE 2"=1 MI.

R. 13 E., T. 11 S., SAN BERNARDINO MER. IMPERIAL CO. CALIF.

WATERFOWL

Refuge Salt Lake National Wildlife Months of September to December 1945

(1) Species Common Name	(2) First Seen		(3) Peak Concentration		(4) Last Seen		(5) Young Produced		(6) Total Estimated for Period
	Number	Date	Number	Date	Number	Date	Broods Seen	Estimated Total	
I. <u>Swans:</u> Whistling swan	15 1	10/13 12/22	2458 1	12/22					2500
II. <u>Geese:</u> Canada goose Cackling goose Brant White-fronted goose Snow goose Blue goose	17 20	9/29 10/13	1800 8600	11/3 12/9					2000 10000
III. <u>Ducks:</u> Mallard Black duck Gadwall Baldpate Pintail Green-winged teal Blue-winged teal Cinnamon teal Shoveller Wood duck Redhead Ring-necked duck Canvas-back Scaup Golden-eye Buffle-head Ruddy duck Fulvous tree duck	8 2 previous period previous period previous period previous period previous period previous period previous period 1 5	12/9 9/1 previous period previous period previous period previous period previous period Dec. 7 9/1	8 1150 150 1,5000 80000 2100 2300 2600 50	12/9 11/3 11/11 11/11 11/11 11/18 11/11 11/18 11/18					10 3000 200 50000 100000 5000 8000 4,000 200
IV. <u>Coots:</u>	20 previous period 1 previous period previous period previous period	11/18 previous period 11/18 previous period previous period	30 300 3 2900 530 2000	11/25 11/18 Dec. 7 11/3 9/8 11/3					100 200 10 4,000 1000 6000

3-1750
(July 1946)

(over)

Form NR-1

Total Production:

Geese _____

Ducks _____

Coots _____

SUMMARIES

Total waterfowl usage during period 1. 5. 6. 1. 5.

Peak waterfowl numbers _____

Areas used by concentrations _____

Principal nesting areas this season _____

Reported by _____

INSTRUCTIONS

(1) Species:

In addition to the birds listed on form, other species occurring on refuge during the reporting period should be added in appropriate spaces. Special attention should be given to those species of local and National significance.

(2) First Seen:

The first refuge record for the species during the season concerned in the reporting period, and the number seen. This column does not apply to resident species.

(3) Peak Concentration:

The greatest number of the species present in a limited interval of time.

(4) Last Seen:

The last refuge record for the species during the season concerned in the reporting period.

(5) Young Produced:

Estimated number of young produced based on observations and actual counts on representative breeding areas. Brood counts should be made on two or more areas aggregating 10% of the breeding habitat. Estimates having no basis in fact should be omitted.

(6) Total:

Estimated total number of the species using the refuge during the period. This figure may or may not be more than that used for peak concentrations, depending upon the nature of the migrational movement.

Note: Only columns applicable to the reporting period should be used. It is desirable that the Summaries receive careful attention since these data are necessarily based on an analysis of the rest of the form.

3-1751

Form NR-1A

(Nov. 1945)

MIGRATORY BIRDS

(other than waterfowl)

Refuge Salton Sea National Wildlife Months of September to December 1945

(1) Species Common Name	(2) First Seen		(3) Peak Numbers		(4) Last Seen		(5) Production		(6) Total Estimated Number
	Number	Date	Number	Date	Number	Date	Number Colonies	Total # Nests	Total Young
I. Water and Marsh Birds:									
<u>Roseate Spoonbill</u>	5	9/30 period	7	9/31 Oct.					
<u>Eared Grebe</u>	previous	period	150	Dec.					
<u>Great Blue Heron</u>	previous	period	20	9/29					
<u>American Egret</u>	previous	period	180	9/29					
<u>Brewster's Egret</u>	previous	period	300	9/29					
<u>White-faced Glossy Ibis</u>	previous	period	200	9/29					
<u>Wood Ibis</u>	previous	period	1600	9/10					
<u>Green Heron</u>	previous	period	4	Dec.					
<u>Least Bittern</u>	previous	period	9	Nov.					
<u>Black-crowned Nt. Heron</u>	previous	period	11	9/29					
<u>Clapper Rail</u>	previous	period	2	Sept					
<u>Sora Rail</u>	previous	period	1	Sept					
<u>Florida Gallinule</u>	previous	period	150	Sept					
<u>White Pelican</u>	previous	period	180	Sept					
<u>Cormorant</u>	previous	period	80	Oct.					
<u>Sandhill Crane</u>	60	Dec.	60						
II. Shorebirds, Gulls and Terns:									
<u>Forrester's Tern</u>	previous	period	10	9/29					
<u>Gull-billed Tern</u>	previous	period							
<u>Black Tern</u>	previous	period	230	9/29					
<u>Ring-billed Gull</u>	previous	period	2000	9/29					
<u>Herring Gull</u>	previous	period	10	Nov.					
<u>Glaucous-winged Gull</u>	previous	period	3	Nov.					
<u>Wilson's Snipe</u>	previous	period	1	Sept.					
<u>Black-necked Stilt</u>	previous	period	500	9/29					
<u>American Avocet</u>	previous	period	80	9/29					
<u>Wilson's Phalarope</u>	previous	period	100	9/29					
<u>Western Sandpiper</u>	previous	period	2800	9/29					
<u>Black-bellied Plover</u>	previous	period	100	Dec.					
<u>Least Sandpiper</u>	previous	period	2000	Dec.					
<u>Hudsonian Curlew</u>	50	9/20	50	9/29					
<u>Long-billed Curlew</u>	1500	9/29	1500	9/29					
<u>Long-billed Dowitcher</u>	3000	9/29	3000	9/29					
<u>Yellow Legs</u>	200	9/29	200	9/29					
<u>Mountain Plover</u>	60	12/15	200	12/22					

(over)

(1)	(2)	(3)	(4)	(5)	(6)
III. <u>Doves and Pigeons:</u> Mourning dove White-winged dove	previous period	2000	Oct.		
IV. <u>Predaceous Birds:</u> Golden eagle Duck hawk Horned owl Magpie Raven Crow	2 1 previous period	2 1 4	11/2 9/26		
Reported by <u>Salton Sea Refuge</u>					

INSTRUCTIONS

- (1) Species: Use the correct names as found in the A.O.U. Checklist, 1931 Edition, and list group in A.O.U. order. Avoid general terms as "seagull", "tern", etc. In addition to the birds listed on form, other species occurring on refuge during the reporting period should be added in appropriate spaces. Special attention should be given to those species of local and National significance. Groups: I. Water and Marsh Birds (Gaviiformes to Ciconiiformes and Gruiformes)
II. Shorebirds, Gulls and Terns (Charadriiformes)
III. Doves and Pigeons (Columbiformes)
IV. Predaceous Birds (Falconiformes, Strigiformes and predaceous Passeriformes)
- (2) First Seen: The first refuge record for the species for the season concerned.
- (3) Peak Numbers: The greatest number of the species present in a limited interval of time.
- (4) Last Seen: The last refuge record for the species during the season concerned.
- (5) Production: Estimated number of young produced based on observations and actual counts.
- (6) Total: Estimated total number of the species using the refuge during the period concerned.

**Saltion Sea National Wildlife
Refuge**

(1) Species	(2) Density		(3) Young Produced		(4) Sex Ratio	(5) Removals			(6) Total	(7) Remarks
Common Name	Cover types, total acreage of habitat	Acres per Bird	Number broods obs'd.	Estimated Total	Percentage	Hunting	For Re-stocking	For Research	Estimated number using Refuge	Pertinent information not specifically requested. List introductions here.
	<div>NO CHANGE IN STATUS</div>									

INSTRUCTIONS

Form NR-2 - UPLAND GAME BIRDS.*

- (1) SPECIES: Use correct common name.
- (2) DENSITY: Applies particularly to those species considered in removal programs (public hunts, etc.). Detailed data may be omitted for species occurring in limited numbers. Density to be expressed in acres per animal by cover types. This information is to be prefaced by a statement from the refuge manager as to the number of acres in each cover type found on the refuge; once submitted, this information need not be repeated except as significant changes occur in the area of cover types. Cover types should be detailed enough to furnish the desired information but not so much as to obscure the general picture. Examples: spruce swamp, upland hardwoods, reverting agriculture land, bottomland hardwoods, short grass prairie, etc. Standard type symbols listed in Wildlife Management Series No. 7 should be used where possible. Figures submitted should be based on actual observations and counts on representative sample areas. Survey method used and size of sample area or areas should be indicated under Remarks.
- (3) YOUNG PRODUCED: Estimated number of young produced, based upon observations and actual counts in representative breeding habitat.
- (4) SEX RATIO: This column applies primarily to wild turkey, pheasants, etc. Include data on other species if available.
- (5) REMOVALS: Indicate total number in each category removed during the report period.
- (6) TOTAL: Estimated total number using the refuge during the report period. This may include resident birds plus those migrating into the refuge during certain seasons.
- (7) REMARKS: Indicate method used to determine population and area covered in survey. Also include other pertinent information not specifically requested.

* Only columns applicable to the period covered should be used.

Calendar Year

Refuge ~~Salton Sea National Wildlife Refuge~~

[illegible]

Remarks:

Reported by

INSTRUCTIONS

- (1) SPECIES: Use correct common name; i.e., Mule deer, black-tailed deer, white-tailed deer. It is unnecessary to indicate sub-species such as northern or Louisiana white-tailed deer.
- (2) DENSITY: Detailed data may be omitted for species occurring in limited numbers. Density to be expressed in acres per animal by cover types. This information is to be prefaced by a statement from the refuge manager as to the number of acres in each cover type found on the refuge; once submitted, this information need not be repeated except as significant changes occur in the area of cover types. Cover types should be detailed enough to furnish the desired information but not so much as to obscure the general picture. Examples: spruce swamp, upland hardwoods, reverting agriculture land, bottomland hardwoods, short grass prairie, etc. Standard type symbols listed in Wildlife Management Series No. 7 should be used where possible. Figures submitted should be based on actual observations and counts on representative sample areas. Survey method used and size of sample area or areas should be indicated under Remarks.
- (3) YOUNG PRODUCED: Estimated total number of young produced on refuge.
- (4) REMOVALS: Indicate total number in each category removed during the year.
- (5) LOSSES: On the basis of known records or reliable estimates indicate total losses in each category during the year.
- (6) INTRODUCTIONS: Indicate the number and refuge or agency from which stock was secured.
- (7) TOTAL REFUGE POPULATION: Give the estimated population of each species on the refuge at period of its greatest abundance and also as of Dec. 31.
- (8) SEX RATIO: Indicate the percentage of males and females of each species as determined from field observations or through removals.

(June 1945)

Refuge Salton See National Wildlife Refuge Year ending April 30.

Refuge Salton See National Wildlife Refuge System Year ending April 30, _____

[illegible]

* List removals by Predator Animal Hunter

Reported by

INSTRUCTIONS

Form NR-4 - SMALL MAMMALS (Include data on all species of importance in the management program; i. e., muskrats, beaver, coon, mink, coyote. Data on small rodents may be omitted except for estimated total population of each species considered in control operations.)

(1) SPECIES:

Use correct common name. Example: Striped skunk, spotted skunk, short-tailed weasel, gray squirrel, fox squirrel, white-tailed jackrabbit, etc. (Accepted common names in current use are found in the "Field Book of North American Mammals" by H. E. Anthony and the "Manual of the Vertebrate Animals of the Northeastern United States" by David Starr Jordan.)

(2) DENSITY:

Applies particularly to those species considered in removal programs. Detailed data may be omitted for species occurring in limited numbers. Density to be expressed in acres per animal by cover types. This information is to be prefaced by a statement from the refuge manager as to the number of acres in each cover type found on the refuge; once submitted, this information need not be repeated except as significant changes occur in the area of cover types. Cover types should be detailed enough to furnish the desired information but not so much as to obscure the general picture. Examples: spruce swamp, upland hardwoods, reverting agriculture land, bottom land hardwoods, short grass prairie, etc. Standard type symbols listed in Wildlife Management Series No. 7 should be used where possible. Figures submitted should be based on actual observations and counts on representative sample areas. Survey method used and size of sample area or areas should be indicated under Remarks.

(3) REMOVALS:

Indicate the total number under each category removed since April 30 of the previous year, including any taken on the refuge by Service Predatory Animal Hunter. Also show any removals not falling under headings listed.

(4) DISPOSITION OF FUR:

On share-trapped furs list the permit number, trapper's share, and refuge share. Indicate the number of pelts shipped to market, including furs taken by Service personnel. Total number of pelts of each species destroyed because of unprime-ness or damaged condition, and furs donated to institutions or other agencies should be shown in the column provided.

(5) TOTAL POPULATION:

Estimated total population of each species reported on as of April 30.

REMARKS:

Indicate inventory method(s) used, size of sample area(s), introductions, and any other pertinent information not specifically requested.

Refuge Salton Sea National Wildlife Refuge Year 19~~41~~52

Botulism

Lead Poisoning or other Disease

Period of outbreak September - October

Kind of disease _____

Period of heaviest losses late September

Species affected _____

Losses:

	Actual Count	Estimated
(a) Waterfowl	_____	<u>300</u>
(b) Shorebirds	_____	<u>300</u>
(c) Other	_____	_____

Number Affected Species	Actual Count	Estimated
_____	_____	_____
_____	_____	_____
_____	_____	_____

Number Hospitalized

No. Recovered

% Recovered

Number Recovered _____

(a) Waterfowl

Number lost _____

(b) Shorebirds

(c) Other

Source of infection _____

Areas affected (location and approximate acreage) _____

Water conditions _____

unknown

Water conditions (average depth of water in sickness areas, reflooding of exposed flats, etc.)

Food conditions _____

Conditions generally good. (Rising sea level, fresh water ponds, etc.)

Condition of vegetation and invertebrate life _____

Remarks _____

Remarks _____

REMARKS:

3-1757
Form NR-7
(April 1946)

PLANTINGS
(Marsh - Aquatic - Upland)

Refuge.....Salton Sea National Wildlife.....Year 194.....

Species	Location of Area Planted	Rate of Seeding or Planting	Amount Planted (Acres or Yards of Shoreline)	Amount & Nature of Propagules	Date of Plant- ing	Survival	Cause of Loss	Remarks
				<u>NONE</u>				

TOTAL ACREAGE PLANTED:

Marsh and aquatic.....
Hedgerows, cover patches.....
Food strips, food patches.....
Forest plantings.....

DIRECTIONS FOR PREPARING FORM NR-8
CULTIVATED CROPS

Cultivated Crops Report Form NR-8 should be prepared on a calendar-year basis for all crops harvested or utilized during the calendar year and submitted with the December 31 refuge report.

Permittee - List each permittee separately. If lands of the refuge are farmed by refuge personnel or hired labor, this should be indicated in the Permittee column.

Permit No. - List the number of the Special Use Permit issued to the individual.

Use or Location - The Unit No. or name specified in the Economic Use Plan should be listed in this column.

Crops Grown - A separate line of the form should be used for each crop grown by each permittee or by refuge personnel. This is important, since if each crop grown by each operator is not specifically enumerated, the report will be of no value for statistical purposes.

Average Yield per Acre - It is important that the average yield per acre of each crop grown by each operator should be shown.

Permittee's Share - Only the number of acres harvested or utilized by the permittee for his own benefit should be shown under the Acres column, and only the number of bushels of farm crops harvested by the permittee for himself should be shown under the Bushels Harvested column. It is requested that all crops harvested be reduced to bushels wherever possible, or, as in the case with the harvesting of seed such as that of sweet clover, alfalfa, brome grass, etc., the total harvested crop in pounds may be shown. Timothy, alfalfa, or other hay harvested by the permittee should be shown on Form NR-10 and should not be shown in the Permittee's Share column.

Government's Share or Return - Harvested - Show the number of bushels harvested for the Government and the acreage from which this share is harvested, both for grain raised by refuge personnel and by permittees. Unharvested - show the exact number of acres of crops allowed to remain unharvested as food and cover for wildlife. An estimate of the number of bushels of grain that is available for the wildlife in such unharvested crops should be shown in the Bushels column.

Compensatory Services, or Cash Revenue - Show other services received by the Government in cooperative farming activities, the number of acres of food strips planted for wildlife, the amount of wildlife crops not otherwise reported that are planted by cooperators for the Service, or the cultivation of wildlife plantations. If the permit is on a fee basis, the total cash revenue received by the Service.

REFUGE GRAIN REPORT

Refuge Salton Sea National Wildlife Refuge Months of September through December, 1957

(1) VARIETY*	(2) ON HAND BEGINNING OF PERIOD	(3) RECEIVED DURING PERIOD	(4) TOTAL	(5) GRAIN DISPOSED OF			(6) ON HAND END OF PERIOD	(7) PROPOSED OR SUITABLE USE*		
				Transferred	Seeded	Fed		Seed	Feed	Surplus
Barley	72	1600	1672		62	644	926	X	X	
Milo Maize	30		30				30	X		
Wild Millet	200		200				200	X		

(8) Indicate shipping or collection points Westmorland, California(9) Grain is stored at Headquarters, Unit I - Sub-headquarters, Unit II

(10) Remarks

*See instructions on back.

REFUGE GRAIN REPORT

This report should cover all grain on hand, received, or disposed of, during the period covered by this narrative report.

Report all grain in bushels. For the purpose of this report the following approximate weights of grain shall be considered equivalent to a bushel: Corn (shelled)—55 lb., corn (ear)—70 lb., wheat—60 lb., barley—50 lb., rye—55 lb., oats—30 lb., soy beans—60 lb., millet—50 lb., cowpeas—60 lb., and mixed—50 lb. In computing volume of granaries, multiply the cubic contents (cu. ft.) by 0.8 bushels.

- (1) List each type of grain separately and specifically, as flint corn, yellow dent corn, square deal hybrid corn, garnet wheat, red May wheat, durum wheat, spring wheat, proso millet, combine milo, new era cowpeas, mikado soy beans, etc. Mere listing as corn, wheat, and soybeans will not suffice, as specific details are necessary in considering transfer of seed supplies to other refuges. Include only domestic grains; aquatic and other seeds will be listed on NR-9.
- (3) Report all grain received during period from all sources, such as transfer, share cropping, or harvest from food patches.
- (4) A total of columns 2 and 3.
- (6) Column 4 less column 5.
- (7) This is a proposed break-down by varieties of grain listed in column 6. Indicate if grain is suitable for seeding new crops.
- (8) Nearest railroad station for shipping and receiving.
- (9) Where stored on refuge: "Headquarters granary," etc.
- (10) Indicate here the source of grain shipped in, destination of grain transferred, data on condition of grain, unusual uses proposed.

3-1759
Form NR-9
(April 1946)

COLLECTIONS AND RECEIPTS OF PLANTING STOCK (Seeds, rootstocks, trees, shrubs)

Refuge Salton Sea National Wildlife Refuge Year 1944-51

Species	Collections				Receipts		Total Amounts on Hand	Amount Surplus
	Amount	Date or Period or Collection	Method	Unit Cost	Amount	Source		
		<u>NONE</u>		<u>NONE</u>				

3-1760

Form NR-10
(April 1946)

HAYING AND GRAZING

Refuge Salton Sea National Wildlife Refuge Year 1945


Permittee	Permit No.	Unit or Location	Actual Acreage Utilized	Animal Use Months	Tons of Hay Harvested	Period of Use From - To	Rate	Total Income	Remarks
Arnold Shields	SAL-10	T3, U2 T18-19, U1	80	1990		2/8 2/28 3/9	2 1/2 dph 2 1/2 dph	1492.50	1990 hd. sheep grazing alfalfa
Arnold Shields	SAL-11	T4, U2 T26, U1	80	4000		5/7 5/23	2 1/2 dph	1500.00	2000 hd. sheep grazing alfalfa
Arnold Shields	SAL-12	T18-19, U1	120	56128 and		4/30	2 1/2 dph	1403.20	1800 hd. sheep grazing
N. J. Ruston	SAL-13	T26, Ua 1 T4, Ua 2		81	13/14	6/11	\$14 ton	1147.00	Harvest hay
N. J. Ruston	SAL-14	T18-19 U1		36.4		6/30	\$ 8 ton	291.20	Harvest hay
N. J. Ruston	SAL-15	T3-4 Ua 2		52.4		6/25	\$10 ton	524.00	Harvest hay
N. J. Ruston	SAL-16	T26, Ua 1		22.25		7/30	\$10 ton	222.50	Harvest hay
Joe Lesicka	SAL-17	T26, Ua 1		32		8/1	\$ 7 ton	224.00	Harvest hay
N. J. Ruston	SAL-18	T18-19 U1		28		9/1	\$2.50 T	70.00	Harvest hay
N. J. Ruston	SAL-19	T3-4, U2		7 1/2		10/4	\$ 3 ton	22.50	Harvest hay
L. C. Copeland	SAL-20	T26, U1	120	730 and		10/14	3 1/2 dph	657.00	1460 hd. sheep grazing
L. C. Copeland	SAL-21	T18-19 U1							
Arnold Shields	SAL-22								

Totals:

Acreage grazed.....	600	Animal use months.....	1895 1/2	Total income Grazing.....	5052.70
Acreage cut for hay.....		Tons of hay cut.....	260 1/2	Total income Haying.....	2501.20



Snow, Canada and White-fronted geese on refuge alfalfa crop, Tracts 18-19, Unit I - December, 1951



Snow geese in refuge alfalfa crop, Tract 27, Unit II - Salton Sea in background - Dec. 1951


Canada Geese on refuge mixed barley and alfalfa crop, Tract 3, Unit II - Rock Hill in background
December, 1951

ALAMO RIVER DUCK
"PRESERVE"

STATE'S UNIT B


WEST BOUNDARY
UNIT II

Snow and Canada geese, Tracts 17-18, Unit II - Dec. 1951



BEFORE

Newly completed fresh water impoundment area,
Tracts 28, 29, and 30, Unit I - March 1951
(West view)



AFTER

Same area as above, September 1951, Wild Millet,
Bulrush and Cattail growth. (East view; arrow
indicates dead tree in above photo) "Wm. Anderson,
State Fish & Game, banded 3500 ducks during Sept."



Wintering birds on refuge - -
"the impoundment area in the
northend of Unit I produced a
good stand of mixed Wild Millet
and Sudangrass...."



A tour of Federal and State
Units....by Lea Act Advisory
Committee took place Sept.
26th...." (State's Sec. 29
along K-lateral)

P&H Dragline, trans-
ferred from Wheeler Refuge,
as sandblasted, painted
and overhauled..Mechanic
Barker, left, and Drag-
line operator James W.
Elton, right. Dec.,
1951.