

The Salton Sea: A Brief History and Biology

Richard G. Thiery
Coachella Valley Water District
P.O. Box 1058
Coachella, CA 92236

Introduction

The Salton Sea is a shallow, hypersaline lake located below sea level in the Salton Basin (or Sink) of southern California (Figure 1). The Salton Sea is the largest body of water within the state of California. At a surface elevation of -227 feet, it covers approximately 381 square miles (243,718 acres). Despite its size, the Salton Sea is not well known nor is it well regarded by some who know of its existence. This low level of recognition for such a major body of water is due to: 1) the Sea's distance from large metropolitan areas, 2) its dependence upon agricultural drainage for continued existence, and 3) its water quality problems, both real and perceived.

Locale

The Salton Basin is a part of the San Andreas fault system, the main fault generally running along the eastern side of the basin. The Salton Sea divides the basin into the Coachella Valley to the north and the Imperial Valley to the south. The Mexicali Valley of Mexico is also part of the Salton Basin.

The Imperial Valley is a nationally important agricultural area, sometimes called the country's winter garden because of the wide variety of crops grown in the valley. Water for Imperial Valley is transported from the Colorado River via the All-American Canal which runs near the Mexico border. The major cities are El Centro, Brawley, and Calexico; the latter is contiguous with the Mexican city of Mexicali.

The Coachella Valley is also a major agricultural area, known for its dates and table grapes. Most Coachella Valley agriculture is irrigated with Colorado River water via the Coachella Canal, a branch of the All-American Canal which runs north along the eastern edge of the basin. There is also a substantial groundwater in the Coachella Valley. Besides agriculture, the Coachella Valley supports a significant resort/golfing industry historically centered around Palm Springs but now including such cities as Palm Desert, Rancho Mirage, Indian Wells, and La Quinta.

The Mexicali Valley of Mexico is

Formation

The Salton Basin is actually an extreme extension of the delta of the Colorado River (Figure 2). Periodically, as sediment deposition blocked the direct route to the Gulf of California, the Colorado River has changed course to flow west through the Salton Basin creating a large lake called Lake Cahuilla. Over the past 1200 years Lake Cahuilla has been created four times, reaching a maximum elevation of 30 feet above sea level (Waters 1983). After a time, sediment deposition again forced the Colorado River back to its original route leaving the water in Lake Cahuilla to evaporate. After each filling Lake Cahuilla left its marks in the forms of ancient beach lines, particularly on the east side of the basin near the Coachella Canal, "bathtub rings" on the rock formations along the west side, and also in shallow shoreline fish traps constructed by Indians along rocky shores.

The sub-sea level was first realized by William Blake in 1853. Almost from its discovery then, have dreamed of importing Colorado River water.

The Salton Sea is a partial, modern incarnation of Lake Cahuilla formed when large flows from the Gila River of Arizona flooded the Colorado River and breached the headworks of an early Imperial Valley irrigation system (Figure 3). From _____ 1905 till _____ 1907, most of the Colorado River flowed into the Salton Basin. Not until the Southern Pacific Railroad constructed a trestle completely across the breach and poured tons of boulders into the break was the river redirected away from the Salton Basin (Kennan 1917).

Because, the flooding into the Salton Basin occurred at the turnout of an irrigation canal, the Salton Sea is almost always described as a man-made body of water. An argument can be made however, that in 1905 and 1906 the Colorado River would have flowed into the Salton Basin regardless of man's presence and rather than causing the creation of the Salton Sea, man may have halted another natural refilling of Lake Cahuilla. By examining tree ring data _____ () showed that the Gila River flows of 1906 were extraordinarily high: the highest in 150 years by a factor of 3 (Figure 4). In addition, it should be remembered that, for much of their lengths, the early irrigation canals utilized existing dry river beds, i.e. they followed. Also, the lower. These dikes may have prevented the river from breaching at other points, therefore creating the illusion that the canal turnout was

While filling, the Sea reached a maximum elevation of -196 feet and then began receding due to evaporation (Figure 2).

After its creation, the elevation of the Sea gradually fell, and its salinity rose, due to evaporation. The Sea's elevation began to rise again in the 1920s as agricultural development, and its associated drainage, continued to increase. Around 1950, salinity in the Sea reached oceanic levels and marine fish were successfully introduced creating a popular sportfishery. Together with stable elevation, the fishery made the Salton Sea a popular destination resort. More recently, the Sea's elevation has and the quality of the sportfishery has declined.

Water Sources and Balance

Table 1. (inputs)
rainfall & evaporation

Water Quality

bacti mostly cleaned up and can't survive in SS
occasional ag chemical discharge
residual DDE

Selenium

Colo R as source
low level in SS water (volatilization)
elevated in biota - eating warning

The Salton Sea Ecosystem

Productivity

The Salton Sea is a eutrophic (highly productive) ecosystem. Carpelan (1961a) estimated Salton Sea primary productivity (rate of photosynthesis) to be about four times that of coastal marine water (the most productive region of the oceans). The Salton Sea is also eutrophic by freshwater ecosystem standards (Table 1).

This high productivity is an important feature of the Salton Sea. It is ultimately responsible for the Sea's popularity with both waterfowl and anglers. Why is the Sea so productive?

The two nutrients which normally limit the productivity of aquatic systems are nitrogen and phosphorous. Because the inflow to the Sea consists largely of agricultural drainage means that the Sea receives large amounts of both of these nutrients (Table 2), quantities in fact which are normally considered acceptable only for much deeper lakes (Table 4). Nutrient levels within the Sea itself are characteristic of systems at the upper end of the productivity scale (Table 3).

The path of productivity within the Sea however, is not straightforward. The following is a brief review of the biology of the Salton Sea followed by a discussion of the Sea's trophic (feeding) interactions.

Phytoplankton

The immediate beneficiaries of the high nutrient input to the Salton Sea are planktonic algae, the phytoplankton. Over a dozen species of algae have been identified from the Sea (Carpelan 1958, 1961a; Arnal 1961).

Algae in the Sea include diatoms - algae which produce silicon cases (frustules) - and some green algae. The dominant group however, particularly in summer, is dinoflagellates, motile algae equipped with two flagella. Included in this group, and recorded from the Sea, are *Gymnodinium* and *Gonyaulax*, the two genera responsible for most "red tides" in coastal marine waters. Dinoflagellates utilize xanthophyll pigments, in addition to chlorophyll, for photosynthesis. Dinoflagellates range in color from colorless to brown or red and their occurrence partially explains the sometimes dark color of the Sea.

Zooplankton

Rotifers

Rotifers are small, relatively simple, invertebrates characteristic of freshwater environments. *Brachionus plicatilis*, the species in the Salton Sea, is a medium-sized (7 250 μm = .01 inch), completely planktonic rotifer. It is common in estuaries throughout the world. There is a large scientific literature on this species because of its value as food for fish larvae.

Several facets of the life history of *B. plicatilis* are optimal at high temperature (Rico-Martinez and Dodson 1992, Snell 1986). *B. plicatilis* feeds most effectively on particles over 2 μm , i.e. it has only a mediocre ability to feed on bacteria so algae and protozoa are its dominant food (Vadstein et al. 1993).

Like most rotifers, *B. plicatilis* is generally parthenogenetic (females producing females) with the occasional appearance of males resulting in the sexual production of resting eggs. High salinity and temperature make production of males less likely and inhibit the hatching of resting eggs (Hino and Hirano 1984; Lubzens et al. 1980, 1985, 1993; Minkoff et al. 1983).

Epp and Winston (1977) reported that *B. plicatilis* is essentially an osmoconformer, although slightly hyperosmotic, and interpret this as an indication of marine ancestry for this species. A review of the literature led Miracle and Serra (1989) to suggest that tolerance to high salinity increases at higher temperature. Population growth apparently ceases at 50 ppt (Pozuelo and Lubian 1993).

In the Salton Sea, *B. plicatilis* is most abundant in summer and autumn (Carpelan 1961). It is undoubtedly an important food item for the early life stage of Salton Sea fish.

Copepod

Copepods are small planktonic crustaceans with planktonic larvae (nauplii). The copepod in the Salton Sea was originally described as *Cyclops dimorphus* (Kiefer 1931, Johnson 1953) but has now been synonymized with *Apocyclops dengizicus* (Lindberg 1940, Kiefer 1967). *A. dengizicus* occurs in saline lakes worldwide (Kiefer 1967, Bayly 1972).

A. dengizicus occurs year-round in the Salton Sea (Dexter 1993) but is most abundant from mid-summer through autumn (Carpelan 1961b). It apparently consumes both phytoplankton (Carpelan 1961b) and rotifers (Dexter 1993) in the Salton Sea..

Adult *A. dengizicus* can survive salinities up to 107 ppt for extended periods but population growth stops at salinities over 57 ppt (Dexter 1993).

Benthos

Foraminifera

Foraminifera are amoeba-like protozoa which secrete calcium carbonate cases (tests) with pores through which pseudopods for locomotion and feeding extend. They are characteristically marine organisms.

Foraminifera are abundant members of the benthos of the Salton Sea; over two dozen species have been recorded (Rogers 1949, Arnal 1961). They feed on bacteria and the rain of detritus (dead organic matter) from the productive plankton of the Sea.

Pileworm

The pileworm, *Neanthes succinea*, is a widely distributed north Atlantic species of polychaete annelid apparently introduced to the California coast when Virginia oysters were introduced into San Francisco Bay in 1869 (Carlton 1979). It tolerates salinities as low as 14 ppt (Fong 1991) and can reach 460 mm (18 inches) in length but is generally much smaller in the Salton Sea (Hartman 1936).

N. succinea usually constructs a U-shaped tube in the sediment. It is a deposit feeder, ingesting substrate and digesting the organic fraction; it is relatively unselective as to particle size (Fong 1987). *N. succinea* has a higher optimal temperature and tolerates lower oxygen levels better than other species of *Neanthes* (Kristensen 1983a, b).

In the Salton Sea, *N. succinea* occurs throughout the Sea in winter but is most abundant in mid-depth sediments where there is a confluence of high organic sediment content (which increases with depth) and sufficient oxygen (which decreases with depth). In summer, *N. succinea* disappears from the deeper areas, apparently due to low oxygen (Carpelan and Linsley 1961b). Worms are often missing their tail segments, presumably a result of feeding activity by fish when they protrude from the worm tubes.

Like many nereid worms, *N. succinea* reproduces by metamorphosing into sexually mature forms, epitokes, which then swarm at night, near the water's surface, fertilizing eggs which develop into planktonic larvae. In many species, and in other populations of *N. succinea*, swarming is a relatively infrequent, sometimes annual, event synchronized by photoperiod and phases of the moon (Hardege et al. 1990, Fong 1991). In the Salton Sea, although spawning is most frequent in spring and autumn, it occurs virtually throughout the year (Carpelan and Linsley 1961b) and does not appear strongly related to lunar cycle (Carpelan and Linsley 1961a). The larval stage lasts about two weeks; worms can mature within 90 days.

Atokous (fully grown but sexually immature) *N. succinea* can tolerate salinities up to 67.5 ppt (Hanson 1972). Production of epitokes however, is depressed at 55 ppt and fertilization of eggs at 45 ppt (Kuhl and Oglesby 1979).

Barnacle

Barnacles are highly modified crustaceans; the great 19th century geologist Louis Agassiz described them as "nothing more than a little shrimplike animal, standing on its head in a limestone house and kicking food into its mouth." Like most crustaceans they have a planktonic nauplius larvae but, in addition, there is a non-feeding cypris larva immediately prior to settling and metamorphosis.

Balanus amphitrite is a western Atlantic species of barnacle which has, within the past century, spread throughout the world, apparently by hitchhiking on ships. It appeared in Hawaii in 1915, San Diego in 1921, Suez in 1924, northern Europe in 1929, and San Francisco in 1939 (Morris et al. 1980). It is now considered cosmopolitan in warm seas, particularly bays, where summer temperatures exceed the 20°C required for reproduction.

The Salton Sea population, *B. amphitrite saltonensis* (Rogers 1949), originated around 1943 with barnacles transported via Navy buoys or flying boats from San Diego Bay (Hilton 1945). Flowerdew (1985) compared the genetic composition of six populations of *B. amphitrite* and recommended eliminating *B. a. saltonensis* as a subspecies, but Raimondi (1992) detected differences in larval morphology and development which remained consistent during culturing, and which therefore, are probably attributable to genetic differentiation.

In the Salton Sea, barnacle larvae can be found in the plankton through most of the year but predominantly in spring and early summer (Carpelan 1961).

Other Species

Hurlbert (personal communication) has suggested that some benthic species which are now only locally abundant might become more important if fish are eliminated. These species include the amphipod (crustacean) *Gammarus mucronatas* and the water boatmen (insect) *Trichocorixa* sp.

Fish

Desert Pupfish

The desert pupfish, *Cyprinodon macularius*, is the only fish species endemic (native) to the Salton basin. It is a small (2-3 inch) member of the killifish family (Cyprinodontidae). The range of the desert

pupfish originally extended across northern Mexico and the Colorado River delta into southern Arizona (Miller 1943), and probably included backwaters along the lower Colorado River (Turner 1983).

Desert pupfish are extremely tolerant of a wide range of physical conditions. The adults can survive oxygen concentrations as low as 0.2 mg/liter (Lowe et al. 1967) and salinities from freshwater (Cowles 1934) to almost 90 ppt (Barlow 1958a). Estimates of adult upper temperature tolerance range from 44.6°C (112.3°F) (Lowe and Heath 1969) to 53.3°C (128°F) (Jordan and Richardson 1929). Their eggs can develop in temperatures up to 35.7°C (96.3°F) and salinities of 70 ppt (Kinne and Kinne 1962, Sweet and Kinne 1964). Growth is faster at 35 and 15 ppt than at 55 ppt (Kinne 1960).

Desert pupfish are shallow water fish. They can feed and spawn in water 3 cm (1.2 inches) deep (Schoenherr 1985). The males generally establish territories and spawn in water less than 15 cm (6 inches) deep (Kynard and Garrett 1979); non-territorial males are reported to sleep in "the shallowest possible water" (Barlow 1961). They are capable of burrowing into sediment to escape predators (Barlow 1961), excessively high or low temperatures (Schoenherr 1988), and possibly also to withstand temporary drying of their habitat. Pupfish move to deep waters only to escape high temperatures (Barlow 1958b).

Desert pupfish are omnivorous, consuming algae, detritus, insects, and small crustaceans (Cox 1972, Naiman 1979). They spawn whenever the temperature is greater than 20°C (68°F) (Kinne 1960, Moyle 1976); the adhesive eggs are laid on the bottom or, possibly, on vegetation (Schoenherr 1988).

Through the 1950s, desert pupfish were abundant in the Salton Sea; Barlow (1961) reported schools of over 10,000 juvenile pupfish along the shore of the Sea. During the 1960s however, the desert pupfish population declined markedly (Crear and Haydock 1971). By 1978, desert pupfish could be found only sporadically (Black 1980).

The decline of desert pupfish was probably due to the introduction of two fish species: tilapia and sailfin mollies. The mechanism of replacement is behavioral interference with pupfish reproduction: Matsui (1981) demonstrated that tilapia and sailfin mollies reduce pupfish reproductive success by interfering with the territorial and courting behavior of desert pupfish. Schoenherr (1985) suggested that mollies excluded desert pupfish from Salton Sea shoreline pools while juvenile tilapia did the same in irrigation drains.

Regardless of the cause, the decline of the Salton Sea and other desert pupfish populations caused the species to be listed as endangered by the California Fish and Game Commission in 1980 and by the U.S. Fish and Wildlife Service in 1986 (USFWS 1986). More recently, CDFG surveys found desert pupfish in a majority of the irrigation drains entering the Sea, particularly at the north end of the Sea, and from a majority of shoreline pools along the southern and eastern shore of the Sea (Lau and Boehm 1991). This apparent partial recovery may be related to unusually high overwinter tilapia mortality _____.

Even when they were abundant, desert pupfish were not considered important in the Salton Sea ecosystem. They were not numerous enough to present serious food competition to other species and, although preyed upon by bairdiella, longjaw mudsuckers and wading birds, were "not important in the diets of any of these" (Walker, Whitney and Barlow 1961).

Sargo

The sargo (*Anisotremus davidsonii*) is a large (up to 17 inches) member of the grunt family (Haemulidae). They are commonly found near shallow structures such as rocks, kelp beds, and pilings from Santa Cruz to Baja California (Goodson 1988). Sargo eggs and larvae are planktonic.

The sargo population in the Salton Sea is derived from 65 fish transported from the Gulf of California in 1951; they became abundant by 1960 (Walker et al. 1961). During 1982-83, sargo constituted 28.3% of the sportfish catch, just edging out bairdiella as the second most commonly caught species (Black 1985). An average of 0.413 sargo were caught per hour of angling; they were the fish most commonly caught by jetty anglers. Sargo caught by anglers averaged 23 cm (9.1 inches) in length and 280 g (0.62 lb) in weight, with maxima of 35 cm (13.8 inches) and 1010 g (2.23 lbs).

Hanson (1970) reported that 96-hour mortality (without acclimation) of juvenile sargo increased at salinities over 50 ppt and that none survived at 62.5 ppt. Brocksen and Cole (1972) found sargo food utilization to be more efficient at 33 ppt than at 37-45 ppt.

Lasker et al. (1972) detected increased larval mortality at salinities of 40 ppt and higher; they also reported a high incidence of larval abnormalities, particularly bent tails. Matsui, Lattin et al. (1991a) reported that larval sargo mortality was markedly higher at 45 ppt and above than at 40 ppt. Matsui, Bond

et al. (1991) reported a decline in the abundance of late larvae over the period 1987-89 which they related to increasing salinity during that period. Matsui et al. (1992) reported that about 8.2% of all sargo larvae collected during 1987-89 were deformed, most with retarded nervous system development.

Bairdiella

The bairdiella, *Bairdiella icistia*, known in the Salton Sea area as gulf croaker, is a small member of the croaker family (Sciaenidae), native to the Pacific coast of Mexico. A related species, *Bairdiella chrysura*, the silver perch is common in Atlantic and Gulf of Mexico estuaries. *Bairdiella* can be distinguished from young orangemouth corvina (below) by an anal spine longer than the anal fin rays.

The Salton Sea bairdiella population originated with 67 fish introduced from the Gulf of California in 1950-51 (Walker et al. 1961); they had become abundant by 1953 (Walker et al. 1961). An apparent genetic founder effect in the Sea's bairdiella population is detectable in the form of a low occurrence of rare alleles, although the overall rate of genetic heterozygosity is comparable to other fish populations (Beckwitt 1987).

The bairdiella diet is dominated by pileworms, particularly swimming pileworms, i.e. the spawning epitokes (Quast 1961). *Bairdiella*'s dependence on pileworms is apparently reflected in both their daily and seasonal movement patterns. *Bairdiella* are more active and do most of their feeding after sunset when pileworms spawn. They are more abundant near shore beginning in spring, peaking in late summer, and then apparently spend the winter in deeper parts of the Sea (Whitney 1961a). This correlates with the distribution of pileworms, which are eliminated from the deeper parts of the Sea by low oxygen levels during summer. *Bairdiella*'s condition factor (a measure of plumpness) appears to fluctuates in relation to pileworm spawning (Whitney 1961a); mass bairdiella mortality episodes have been related to poor condition (Quast 1961).

May is the peak spawning month for bairdiella (Whitney 1961a). Spawning takes place in early evening; the eggs are planktonic.

During 1982-83, bairdiella ranked third (just behind sargo) as the species most frequently caught by Salton Sea fishermen, making up 28.1% of the total catch. An average of 0.410 bairdiella were caught per hour of angling (Black 1985). *Bairdiella* caught by anglers averaged 24 cm (9.5 inches) in length and 160 g (0.35 lb) in weight, with maxima of 41 cm (16.1 inches) and 760 g (1.68 lbs).

Hanson (1970) reported that juvenile bairdiella transferred directly to higher salinities tolerated 52.5 ppt for 96 hours, although there was 60% mortality at 55 ppt. Mortality of yearling bairdiella started at 52.5 ppt (40%) and increased until no fish survived at 75 ppt. Brocksen and Cole (1972) found that, over a salinity range of 29 to 45 ppt, bairdiella food utilization was most efficient at 37 ppt. Lasker et al. (1972) detected increased larval mortality at salinities of 40 ppt and higher; they also reported a high incidence of larval abnormalities, particularly bent tails. Matsui et al. (1992) reported that about 2.2% of all bairdiella larvae collected during 1987-89 were deformed, most with retarded nervous system development. Matsui, Bond et al. (1991) found a decline in the abundance of bairdiella eggs and early larvae over the period 1987-89 during which salinity rose from 38 to 44 ppt, although the abundance of late larvae increased.

May (1975a, 1975b) has done detailed work on the effect of salinity and temperature on various aspects of bairdiella reproduction. The activity time of bairdiella spermatozoa falls from 3 minutes at 35 ppt salinity to 1.5 minutes at 55 ppt (May 1975a), although it's considerably longer at lower salinities. Fertilization success and hatching rate decline at salinities over 45 ppt and 40 ppt, respectively. Incubation at 30°C reduces the hatching rate compared to 27°C. Larvae reared at 30°C and salinities over 40 ppt had a notable frequency of spinal deformities. Hatched larvae can tolerate temperatures up to 31°C and salinity up to 45 ppt for 72 hours (May 1975b). May (1976) found that fertilization success was higher in Salton Sea water than in ocean water but larval survivorship much lower. By using charcoal filtration to remove organic contaminants and membrane filtration to eliminate protozoan infestations, May related these results to the ionic composition of Salton Sea water.

Tilapia

Tilapia are members of the family Cichlidae, native to Africa but now distributed worldwide because of their popularity in aquaculture. Because of their importance in aquaculture, there is a large scientific literature on several tilapia species, including *Tilapia mossambica*.

The tilapia in the Salton Sea is a mouthbrooding species most similar to *Tilapia mossambica* and may be derived from several sources. Prior to 1963 the Arizona Game and Fish Department introduced *T. mossambica* into canals and drainage ditches leading to the Colorado River; from there they could have spread to the Imperial Irrigation District (IID) and Coachella Valley Water District (CVWD) irrigation systems (St. Amant 1966b). In 1964, *T. mossambica* were found to have escaped from an aquaculture pond into a small creek and drainage sump in the Mineral Spas area east of the Sea (St. Amant 1966a). In 1966 and 1967 the California Department of Fish and Game stocked *T. mossambica* into Wiest Lake on the Central Main Canal portion of the IID canal system (Hoover 1971). And from 1971 through 1982 *Tilapia zilli* (redbelly tilapia) were imported, under CDFG permit, by IID and CVWD for experiments in aquatic weed control in irrigation canals (Pelzman 1973). Since the Salton Sea tilapia are mouthbrooders, while *T. zilli* keeps its eggs in the nest, importation by the irrigation districts appears to be least likely source of the Salton Sea's tilapia, although *T. zilli* has been found in irrigation drains around the Sea (Schoenherr 1979).

T. mossambica feeds on a variety of foods - zooplankton, phytoplankton, and benthic detritus - and may switch among foods on a seasonal basis (Maitipe and De Silva 1985). *T. mossambica* does not filter planktonic algae as efficiently as other tilapia species (Robinson et al. 1990) but can grow on a pure phytoplankton diet (Gaigher 1982). The dominant food items however, are usually detritus and benthic algae (Man and Hodgkiss 1977, de Moor et al. 1986). Benthic feeding is accomplished by repeatedly biting into the substrate at a 45° angle, leaving the bottom pockmarked with small depressions (Bowen 1979).

Male *T. mossambica* build nests about 50 cm (20 inches) in diameter in shallow water. The nests are usually constructed in clusters near sheltered areas (De Silva and Sirisena 1988). The female lays her eggs, the male swims over them releasing sperm, and the female immediately turns and picks the fertilized eggs up into her mouth. The larvae may receive some nutrient support from the mother during mouthbrooding (Kishida and Specker 1994).

During 1982-82, *T. mossambica* was the species most frequently caught by Salton Sea fishermen in general, and boat and shore anglers in particular, making up almost 41% of the total catch. An average of 0.595 tilapia were caught per hour of angling (Black 1985). Tilapia caught by anglers averaged 27 cm (10.6 inches) in length and 375 g (0.83 lb) in weight, with maxima of 45 cm (17.7 inches) and 1,580 g (3.48 lbs). No redbelly tilapia were found during this creel survey.

T. mossambica shows physiological signs of osmoregulatory collapse at 15°C (Smit et al. 1981); mortality occurs from 9.5°C (Behrends et al. 1990) to 15°C (Stauffer 1986). At low temperatures, stress impairs the immune system and the proximate cause of mortality is often fungal infection (Oldewage and As 1987, Behrends et al. 1990). *T. mossambica* can tolerate temperatures as high as 37°C and their preferred temperature is about 34-35°C (Stauffer 1986), although growth is slower at 35°C than at 30°C (Price et al. 1985).

The salinity tolerance of tilapia has been most recently reviewed by Stickney (1986). Five to 10 week old *T. mossambica* can tolerate 70 ppt for indefinite periods (Potts et al. 1967). Juvenile and adult *T. mossambica* can be acclimated to 60 ppt (Assem and Hanke 1979, Kültz and Onken 1993, Kültz and Jürss 1993). Popper and Lichatowich (1975) reported that *T. mossambica* reproduction was so prolific at salinities up to 49 ppt that it was necessary to introduce predators for population control.

There is conflicting evidence on the optimal salinity for *T. mossambica*. It is more susceptible to starvation (Jürss et al. 1984) and organic pollutants (Dang, 1986) at high salinity (i.e. seawater) and respiration is generally lower in fresh water (Job 1969). On the other hand, growth is generally faster at higher salinities (Canagaratnam 1966, Uchida and King 1962).

Orangemouth Corvina

The orangemouth corvina, *Cynoscion xanthalmus*, is a large member of the croaker family (Sciaenidae), native to the Pacific coast of Mexico. Several related species, including spotted seatrout, *Cynoscion nebulosus*, and weakfish, *Cynoscion regalis*, are commercially fished along the Atlantic coast.

The Salton Sea corvina population originated with approximately 250 fish imported from the Gulf of California from 1950 to 1956. By 1957, the Sea's population was estimated at 800,000 fish (Whitney 1961b).

Young of the year corvina over two inches length feed largely on pileworms, competing for food with bairdiella. By their second year corvina have reached 159 mm length (6.26 inches) (Anderson 1971) and are largely feeding on bairdiella (Whitney 1961b) and, since their introduction, presumably tilapia. Other species such as sargo and mudsuckers are also readily consumed.

Whitney (1961b) reported anecdotal accounts suggesting that corvina may congregate near freshwater inflows to spawn. There are also some indications that Salton Sea corvina spawn in autumn as well as spring. Over 1.5 million planktonic eggs can be produced by a single female (Prentice et al. 1989).

During 1982-83, corvina made up 2.9% of the sportfish catch; an average of 0.042 corvina were caught per hour of angling (Black 1985). Corvina caught by anglers averaged 59 cm (23.2 inches) in length and 2000 g (4.41 lbs) in weight, with maxima of 108 cm (42.5 inches) and 12,000 g (26.46 lbs). Walker et al. (1961) recorded a 32 lb corvina.

Hanson (1970) reported that 100% of juvenile corvina survived 96 hours without acclimation at 52.5 ppt and had survivorship over 90% at 55 and 57.5 ppt, but that mortality was complete at 62.5 ppt. Brocksen and Cole (1972) found that corvina food assimilation was more efficient at a salinity of 37 ppt than at 29 or 45 ppt and that oxygen consumption was lowest at 33 ppt.

Matsui, Lattin et al. (1991b) found that adult corvina continued to grow at salinities up to 55 ppt. Oocytes matured at salinities up to 50 ppt although spawning did not occur at 45 or 50 ppt, and egg mortality remained below 10% at salinities up to 55 ppt. Prorated data from Matsui et al. (1992) indicates that about 1.1% of all corvina larvae collected from the Sea during 1987-89 were deformed, most with retarded nervous system development. Matsui, Bond et al. (1991) found a decline in the abundance of corvina eggs and late larvae in the Sea over the period 1987-89 which they related to rising salinity.

Other Species

There are three other fish species common along the edge of the Sea and in the irrigation drains leading to it.

The longjaw mudsucker (*Gillichthys mirabilis*) is a large (5-8 inches) member of the goby family common on mudflats in California estuaries from San Francisco south and in the Gulf of California. The Salton Sea population is derived from 500 fish transported from San Diego Bay in 1930 (Walker et al. 1961). Mudsuckers survive low tides by burrowing into the substrate. They can tolerate low oxygen levels by breathing air through their highly vascularized mouth (Todd and Ebeling 1966) and are capable of some terrestrial movement (Todd 1968). They can survive for several days in freshwater and have been collected from water of 82.5 ppt salinity (Barlow 1963). Mudsuckers are also tolerant of high sulfide levels (Bagarinao and Vetter 1989). Their hardiness has made mudsuckers a popular baitfish. In the Salton Sea, mudsuckers consume pileworms, insects, smaller fish, and barnacles (Walker et al. 1961). Mudsucker males defend breeding territories around their burrows (Weisel 1947); the slightly adhesive eggs are laid on the sides of the burrows (Moyle 1976); the larvae are planktonic.

The sailfin molly (*Poecilia latipinna*) is a livebearer (family Poeciliidae) native to the Atlantic coast of North America from South Carolina to the Yucatan peninsula (Lee et al. 1980) where it is found in both fresh and brackish water. The sailfin molly is a popular species for scientific studies; its social behavior in particular (e.g. Sumner et al. 1994, Schlupp et al. 1994), has been extensively studied.

The Salton Sea population may have been established by escapees from a tropical fish farm in the 1960s (St. Amant 1966a). A second species of molly, the shortfin molly (*Poecilia mexicana* = *Poecilia sphenops*) is also occasionally collected from irrigation drains around the Sea (Schoenherr 1979).

Sailfin mollies generally feed on detritus and plant material and, opportunistically, on insect larvae (Harrington and Harrington 1961). They die if the water temperatures reaches 15°C (59°F); the juveniles generally prefer temperatures over 30°C (86°F) (Stauffer et al. 1985). Adults can tolerate salinities of 80 (Nordlie et al. 1992) to 87 ppt (Herre 1929).

The porthole livebearer (*Poeciliopsis gracilis*) is another livebearing species (family Poeciliidae). Its native distribution is Central America and southern Mexico (Lee et al. 1980). This species appeared in the Salton Sea in the early 1970s, possibly an escapee from a tropical fish farm (Mearns 1975).

Discussion

Lasker et al. 1972 noted that sargo and bairdiella eggs floated in Salton Sea water but not Pacific Ocean water from La Jolla and suggested that this was an indication of adaptation to Salton Sea salinity.

May 1976: [lack of planktonic fish +] high food availability (zoop) = high larval survivorship; makes up for poor larval survivorship due to odd ion makeup

Bairdiella and *Cynoscion* occur together in estuaries

Bairdiella mortality 1953-56 (Whitney 1961)

most salinity work based on seawater ionic composition - may not completely apply to Salton Sea

Walker, Whitney and Carpelan (1961) recommend addition of mysid/amphipod non carnivorous invertebrate to parallel pileworm in food chain; maybe other worms (*Capitella*)

Conclusions

Two traits stand out in this review of Salton Sea biota. First, the invertebrate fauna of the Salton Sea is, for the most part, a collection of highly successful, widely distributed, predominantly estuarine, species. It is not a simple transplant from the Gulf of California nor a highly specialized hypersaline fauna. Second, much of the seasonality common in the reproduction of marine species has been lost. This is undoubtedly an artifact of the southern California desert climate and has been a factor in the productivity of the Sea's fishery.

The path of productivity from the phytoplankton to the Sea's fishery is convoluted because of the lack of an adult planktivorous fish. The populations of the invertebrate herbivores (rotifers and copepods) can respond quickly to changes in algal abundance. This means there is almost always abundant food for larval fish. But the transfer of productivity via live organisms stops at this point. Without a predator to feed on them, the zooplankton die naturally and sink to the bottom of the Sea as detritus to be consumed by bacteria, foraminifera, and pileworms (Arnal 1961, Carpelan 1961).

The lack of an effective planktivorous fish means that the productivity of the Sea has to travel through the benthos before reaching the fishery. There is however, a question about the effectiveness of bairdiella, historically the most important forage fish, in benthic feeding. Whitney (1961) reported that the condition (plumpness) of bairdiella correlated closely with the occurrence of pileworm swarming. This implies that bairdiella could not reliably feed on pileworms directly from the bottom of the Sea. If pileworm swarming were as highly seasonal in the Salton Sea as it is in other habitats, it is possible that bairdiella would not survive in the Sea. This apparent benthic feeding limitation of bairdiella may also explain their partial replacement by *Tilapia mossambica*. *T. mossambica* has a well-developed ability to forage food directly from benthic substrate (Bowen 1979).

Whether via bairdiella or tilapia, the Salton Sea food chain leading to corvina, the primary sport fish, is five or six steps:

- 1) phytoplankton
- 2) zooplankton
- 3) (bacteria/foraminifera)
- 4) pileworm
- 5) bairdiella/tilapia
- 6) corvina

In most lakes, the chain to reach a similar-sized sport fish would be only four steps:

- 1) phytoplankton
- 2) zooplankton
- 3) planktivorous fish
- 4) piscivorous fish (sport fish)

In conclusion, the successful, eurytopic nature of the Salton Sea's invertebrate fauna seems to indicate continued resilience for the Sea's community as a whole, but the pattern and length of the food chain apparently places the sportfishery at considerable risk.

Search: "fish" and "osmoregulation"

Literature Cited

- Anderson, K.R. 1971. Age and growth of orangemouth corvina in the Salton Sea, California. Transactions of the California-Nevada Chapter of the American Fisheries Society 43-50.
- Arnal, R.E. 1961. Limnology, sedimentation, and microorganisms of the Salton Sea, California. Bulletin of the Geological Society of America 72:427-78.
- Assem, H. and W. Hanke. 1979. Concentrations of carbohydrates during osmotic adjustment of the euryhaline teleost, *Tilapia mossambica*. Comparative Biochemistry and Physiology 64A:5-16.
- Bagarinao, T. and R.D. Vetter. 1989. Sulfide tolerance and detoxification in shallow-water marine fishes. Marine Biology 103:291-302.
- Barlow, G.W. 1958a. High salinity mortality of desert pupfish, *Cyprinodon macularius*. Copeia 1958:231-2.
- Barlow, G.W. 1958b. Daily movements of desert pupfish, *Cyprinodon macularius*, in shore pools of the Salton Sea, California. Ecology 39:580-7.
- Barlow, G.W. 1961. Social behavior of the desert pupfish, *Cyprinodon macularius*, in the field and in the aquarium. American Midland Naturalist 65:339-59.
- Barlow, G.W. 1963. Species structure of the gobiid fish *Gillichthys mirabilis* from coastal sloughs of the eastern Pacific. Pacific Science 17:47-72.
- Bayly, I.A.E. 1972. Salinity tolerance and osmotic behavior of animals in athalassic saline and marine hypersaline waters. Annual Review of Ecology and Systematics 3:233-68.
- Beckwitt, R. 1987. Population genetics of an introduced species: *Bairdiella icistius* in the Salton Sea. Bulletin of the Southern California Academy of Science 86:34-40.
- Behrends, L.L., J.B. Kingsley, and M.J. Bulls. 1990. Cold tolerance in maternal mouthbrooding tilapias: phenotypic variation among species and hybrids. Aquaculture 85:271-80.
- Black, G. 1980. Status of the Desert Pupfish, *Cyprinodon macularius* (Baird and Girard), in California. Inland Fisheries Endangered Species Program Special Publication 80-1. 42 pp. California Department of Fish and Game.
- Black, G. 1985. The Salton Sea Sport Fishery. Region 5 Informational Bulletin 10-5-1985. 43 pp. California Department of Fish and Game.
- Bowen, S.H. 1979. A nutritional constraint in detritivory by fishes: the stunted population of *Sarotherodon mossambicus* in Lake Sibaya, South Africa. Ecological Monographs 49:17-31.
- Boyd, K. and R. Garrett. 1979. Reproductive ecology of the Quitobaquito pupfish from Organ Pipe Cactus National Monument, Arizona. National Park Service Transactions and Proceedings 5:625-629.
- Brocksen, R.W. and R.E. Cole. 1972. Physiological responses of three species of fishes to various salinities. Journal of the Fisheries Research Board of Canada 29:399-405.
- Canagaratnam, P. 1966. Growth of *Tilapia mossambica* Peters in different salinities. Bulletin of the Fisheries Research Station of Ceylon 19:47-50.
- Carpelan, L.H. 1958. The Salton Sea, physical and chemical characteristics. Limnology and Oceanography 3:373-86.
- Carpelan, L.H. 1961a. Phytoplankton and plant productivity. pp. 33-42 in Walker 1961.
- Carpelan, L.H. 1961b. Zooplankton. pp. 49-61 in Walker 1961.
- Carpelan, L.H. and R.H. Linsley. 1961a. The spawning of *Neanthes succinea* in the Salton Sea. Ecology 42:189-90.
- Carpelan, L.H. and R.H. Linsley. 1961b. The pile worm, *Neanthes succinea*. pp. 63-76 in Walker 1961
- Carlton, J.T. 1979. Introduced invertebrates of San Francisco Bay. pp. 427-44. in T.J. Conomos (ed.); San Francisco Bay, the Urbanized Estuary; Pacific Division of the American Association for the Advancement of Science.
- Cowles, R.B. 1934. Notes on the ecology and breeding habits of the desert minnow, *Cyprinodon macularius* Baird and Girard. Copeia 1934:40-2.
- Cox, T.J. 1972. The food habits of the desert pupfish (*Cyprinodon macularius*) in Quitobaquito Springs, Organ Pipe National Monument, Arizona. Journal of the Arizona Academy of Science 7:25-7.

- Crear, D. and I. Haydock. 1971. Laboratory rearing of the desert pupfish, *Cyprinodon macularius*. *Fishery Bulletin* 69:151-6.
- Dang, A.D. 1986. Branchial Na⁺-K⁺-ATPase inhibition in a freshwater euryhaline teleost, tilapia (*Oreochromis mossambicus*), during short-term exposure to toluene or naphthalene: influence of salinity. *Environmental Pollution* 42A:273-86.
- De Silva, S.S. and H.K.G. Sirisena. 1988. Observations on the nesting habits of *Oreochromis mossambicus* (Peters) (Pisces: Cichlidae) in Sri Lankan reservoirs. *Journal of Fish Biology* 33:689-96.
- Dexter, D.M. 1993. Salinity tolerance of the copepod *Apocyclops dengizicus* (Lepeschkin, 1900), a key food chain organism in the Salton Sea, California. *Hydrobiologia* 267:203-9.
- Epp, R.W., and P.W. Winston. 1977. Osmotic regulation in the brackish-water rotifer *Brachionus plicatilis* (Muller). *Journal of Experimental Biology* 68:151-6.
- Flowerdew, M.W. 1985. Indices of genetic identity and distance in three taxa within the *Balanus amphitrite* Darwin complex (Cirripedia, Thoracica). *Crustaceana* 49:7-15.
- Fong, P.P. 1987. Particle-size utilization in the introduced polychaete *Neanthes succinea* in San Francisco Bay. *Pacific Science* 41:37-43.
- Fong, P.P. 1991. The effects of salinity, temperature, and photoperiod on epitokal metamorphosis in *Neanthes succinea* (Frey et Leuckart) from San Francisco Bay. *Journal of Marine Biology and Ecology* 149:177-90.
- Gaigher, I.G. 1982. The growth and production of Mocambique tilapia (*Oreochromis mossambicus*) fed on algae in small tanks. *Water SA* 8:142-144.
- Goodson, G. 1988. *Fishes of the Pacific Coast*. 267 pp. Stanford University Press.
- Hanson, J.A. 1970. Salinity tolerances for Salton Sea fishes. Inland Fisheries Administrative Report No. 70-2. 8 pp. California Department of Fish and Game.
- Hanson, J.A. 1972. Tolerance of high salinity by the pileworm, *Neanthes succinea*, from the Salton Sea California. *California Fish and Game* 58:152-4.
- Hardege, J.D., H.D. Bartels-Hardege, E. Zeeck, and F.T. Grimm. 1990. Induction of swarming of *Nereis succinea*. *Marine Biology* 104:291-5.
- Harrington, R.W., Jr. and E.S. Harrington. 1961. Food selection among fishes invading a high subtropical salt marsh: from onset of flooding through the progress of a mosquito brood. *Ecology* 42:646-66.
- Hartman, O. 1936. New species of polychaetous annelids of the family Nereidae from California. *Proceedings of the U.S. National Museum* 83:467-80.
- Hendricks, L.J. 1961. The threadfin shad, *Dorosoma pretense*. pp. 93-4 in Walker 1961.
- Herre, A.W. 1929. An American cyprinodont in Philippine salt ponds. *Philippine Journal of Science* 38:121-6.
- Hilton, J. 1945. Where barnacles grow on the sage. *Desert Magazine* (March):4-5.
- Hino, A. and R. Hirano. 1984. Relationship between water temperature and bisexual reproduction rate in the rotifer *Brachionus plicatilis*. *Bulletin of the Japanese Society of Scientific Fisheries* 50:1481-1485.
- Hoover, F.G. 1971. Status Report on *Tilapia mossambica* (Peters). California Department of Fish and Game Inland Fisheries Administrative Report 71-16. 32 pp.
- Job, S.V. 1969. The respiratory metabolism of *Tilapia mossambica* (Teleostei) I: the effect of size, temperature and salinity. *Marine Biology* 2:121-6.
- Johnson, M.W. 1953. The copepod *Cyclops dimorphus* Kiefer from the Salton Sea. *American Midland Naturalist* 49:188-92.
- Jordan, D.S. and R.E. Richardson. 1929. Description of a new species of killifish, *Lucania browni*, from a hot spring in lower California. *Proceedings of the U.S. National Museum* 33:319-21.
- Jürss, K., T. Bittorf, T. V"nkler, and R Wacker. 1984. Biochemical investigations into the influence of environmental salinity on starvation of the tilapia *Oreochromis mossambicus*. *Aquaculture* 40:171-82.
- Kennan, G.F. 1917. *The Salton Sea: An Account of Harriman's Fight with the Colorado River*. MacMillan, New York. 106 pp.
- Kinne, O. 1960. Growth, food intake, and food conversion in a euryplastic fish exposed to different temperatures and salinities. *Physiological Zoology* 33:288-317.

- Kristensen, E. 1983a. Ventilation and oxygen uptake by three species of *Nereis* (Annelida: Polychaeta) I: Effects of hypoxia. *Marine Ecology-Progress Series* 12:289-97.
- Kiefer, F. 1931. Zur Kenntnis der freilebenden Süßwassercopepoden, insbesondere der Cyclopiden Nordamerikas. *Zoologische Jahrbuch Abteilung Systematische* 61(5/6):269-71.
- Kiefer, F. 1967. Cyclopiden aus salzhaltigen binnigen binnengewässern Australiens (Copepoda). *Crustaceana* 12:292-302.
- Kinne, O. 1960. Growth, food intake, and food conversion in a euryplastic fish exposed to different temperatures and salinities. *Physiological Zoology* 33:288-317.
- Kinne, O. and E.M. Kinne. 1962. Rates of development in embryos of a cyprinodont fish exposed to different temperature-oxygen combinations. *Canadian Journal of Zoology* 40:231-53.
- Kishida, M. and J.L. Specker. 1994. Vitellogenin in the surface mucus of tilapia (*Oreochromis mossambicus*): possibility for uptake by the free-swimming embryos. *Journal of Experimental Zoology* 268:259-68.
- Kristensen, E. 1983b. Ventilation and oxygen uptake by three species of *Nereis* (Annelida: Polychaeta) II: Effects of temperature and salinity changes. *Marine Ecology-Progress Series* 12:299-306.
- Kuhl, D.L. and L.C. Oglesby. 1979. Reproduction and survival of the pileworm *Nereis succinea* in higher Salton Sea salinities. *Biological Bulletin* 157:153-65.
- Kültz, D. and K.Jürss. 1993. Biochemical characterization of isolated branchial mitochondria-rich cells of *Oreochromis mossambicus* acclimated to fresh water or hypersaline sea water. *Journal of Comparative Physiology B* 163:406-12.
- Kültz, D. and H. Onken. 1993. Long-term acclimation of the teleost *Oreochromis mossambicus* to various salinities: two different strategies in mastering hypertonic stress. *Marine Biology* 117:527-33.
- Lasker, R., R.H. Tenaza, and L.L. Chamberlain. 1972. The response of Salton Sea fish eggs and larvae to salinity stress. *California Fish and Game* 58:58-66.
- Lau, S. and C. Boehm. 1991. A Distribution Survey of Desert Pupfish (*Cyprinodon macularius*) Around the Salton Sea. Section 6 Project EF90XII-1. 21 pp. California Department of Fish and Game.
- Lowe, C.H. and W.G. Heath. 1969. Behavioral and physiological responses to temperature in the desert pupfish *Cyprinodon macularius*. *Physiological Zoology* 42:53-9.
- Lowe, C.H., D.S. Hinds, and E.A. Halpern. 1967. Experimental catastrophic selection and tolerances to low oxygen concentration in native Arizona freshwater fishes. *Ecology* 48:1013-7.
- Lubzens, E., R. Fishler, and V. Berdugo-White. 1980. Induction of sexual reproduction and resting egg production in *Brachionus plicatilis* reared in sea water. *Hydrobiologia* 73:55-58.
- Lindberg, K. 1940. Cyclopides (Crust. Cop.) de l'Inde IV. Une revision des representants indiens et iraniens du sousgenre *Metacyclops* Kiefer du genre *Cyclops* Muller. *Records of the Indian Museum* 42:567-88.
- Lubzens, E., G. Minkoff, and S. Marom. 1985. Salinity dependence of sexual and asexual reproduction in the rotifer *Brachionus plicatilis*. *Marine Biology* 85:123-126.
- Lubzens, E., Y. Wax, G. Minkoff, and F. Adler. 1993. A model evaluating the contribution of environmental factors to the production of resting eggs in the rotifer *Brachionus plicatilis*. *Hydrobiologia* 255/256:127-38.
- Maitipe, P. and S.S. De Silva. 1985. Switches between zoophagy, phytophagy and detritivory of *Sarotherodon mossambicus* (Peters) populations in twelve man-made Sri Lankan lakes. *Journal of Fish Biology* 26:49-61.
- Man, H.S.H. and I.J. Hodgkiss. 1977. Studies on the ichthyo-fauna in Plover Cove Reservoir, Hong Kong: feeding and food relations. *Journal of Fish Biology* 11:1-13.
- Marsh, P.C. and D.W. Sada. 1993. Desert Pupfish (*Cyprinodon macularius*) Recovery Plan. U.S. Fish and Wildlife Service, Phoenix. 67 pp.
- Matsui, M.L. 1981. The Effects of Introduced Teleost Species on the Social Behavior of *Cyprinodon macularius californiensis*. M.S. Thesis. 61 pp. Occidental College, Los Angeles.
- Matsui, M.L., A. Bond, G. Jordan, R. Moore, P. Garrahan, K. Iwanaga, and S. Williams. 1991. Abundance and Distribution of the Ichthyoplankton in the Salton Sea, California in Relation to Water Quality. Final Report, Federal Aid Project F-51-R, Study No. 3, California Department of Fish and Game. 40 pp.

- Matsui, M., J.E. Hose, P. Garrahan, and G.A. Jordan. 1992. Developmental defects in fish embryos from Salton Sea, California. *Bulletin of Environmental Contamination and Toxicology* 48:914-920.
- Matsui, M.L., G.L. Lattin, R. Moore, C. Mulski, and A.B. Bond. 1991a. Salinity Tolerance of *Anisotremus davidsonii*. Final Report, Federal Aid Project F-51-R, Study No. 2. California Department of Fish and Game. 17 pp.
- Matsui, M.L., G.L. Lattin, R. Moore, C. Mulski, and A.B. Bond. 1991b. Salinity tolerance of *Cynoscion xanthulus*. Final Report, Federal Aid Project F-51-R, Study No. 2. California Department of Fish and Game. 24 pp.
- May, R.C. 1975a. Effects of temperature and salinity on fertilization, embryonic development, and hatching in *Bairdiella icistia* (Pisces: Sciaenidae), and the effect of parental salinity acclimation on embryonic and larval salinity tolerance. *Fishery Bulletin* 73:1-22.
- May, R.C. 1975b. Effects of acclimation on the temperature and salinity tolerance of the yolk-sac larvae of *Bairdiella icistia* (Pisces: Sciaenidae). *Fishery Bulletin* 73:249-55.
- May, R.C. 1976. Effects of Salton Sea water on the eggs and larvae of *Bairdiella icistia* (Pisces: Sciaenidae). *California Fish and Game* 62:119-31.
- Mearns, A.J. 1975. *Poeciliopsis gracilis* (Heckel), a newly introduced poeciliid fish in California. *California Fish and Game* 61:251-3.
- Miller, R.R. 1943. The status of *Cyprinodon macularius* and *Cyprinodon nevadensis*, two desert fishes of western North America. Occasional Paper of the University of Michigan Museum of Zoology No. 473, 25 pp.
- Minkoff, G., E. Lubzens, and D. Kahan. 1983. Environmental factors affecting hatching of rotifer (*Brachionus plicatilis*) resting eggs. *Hydrobiologia* 104:61-9.
- Miracle, M.R. and M. Serra. 1989. Salinity and temperature influence in rotifer life history characteristics. *Hydrobiologia* 186/187:81-102.
- Morris, R.H., D.P. Abbott, and E.C. Haderlie. 1980. *Intertidal Invertebrates of California*. pp. 522-4. Stanford University Press.
- Moyle, P.B. 1976. *Inland Fishes of California*. University of California Press, Berkeley. 405 pp.
- Naiman, R.J. 1979. Preliminary food studies of *Cyprinodon macularius* and *Cyprinodon nevadensis* (Cyprinodontidae). *Southwestern Naturalist* 24:538-41.
- Nordlie, F.G., D.C. Haney and S.J. Walsh. 1992. Comparisons of salinity tolerances and osmotic regulatory capabilities in populations of sailfin molly (*Poecilia latipinna*) from brackish and fresh waters. *Copeia* 1992:741-6.
- Oglesby, L.C. 1977. Reproduction and Survival of the Pileworm *Nereis succinea* in Higher Salton Sea Salinities. NTIS. 42 pp.
- Oldewage, W.H. and J.G. van As. 1987. Parasites and winter mortalities of *Oreochromis mossambicus*. *South African Journal of Wildlife Research* 17:7-12.
- Pelzman, R.J. 1973. A Review of the Life History of *Tilapia zillii* with a Reassessment of Its Desirability in California. Inland Fisheries Administrative Report 74-1. California Department of Fish and Game.
- Popper, D. and T. Lichatowich. 1975. Preliminary success in predator control of *Tilapia mossambica*. *Aquaculture* 5:213-4.
- Potts, W.T.W., M.A. Foster, P.P. Rudy, and G.P. Howells. 1967. Sodium and water balance in the cichlid teleost, *Tilapia mossambica*. *Journal of Experimental Biology* 47:461-70.
- Pozuelo, M. and L.M. Lubian. 1993. Asexual and sexual reproduction in the rotifer *Brachionus plicatilis* cultured at different salinities. *Hydrobiologia* 255/256:139-43.
- Prentice, J.A., R.L. Colura, and B.W. Bumguardner. 1989. Observations on induced maturation and spawning of orangemouth corvina. *California Fish and Game* 75:27-32.
- Price, E.E., J.R. Stauffer, Jr. and M.C. Swift. 1985. Effect of temperature on growth of juvenile *Oreochromis mossambicus* and *Sarotherodon melanotheron*. *Environmental Biology of Fishes* 13:149-152.
- Quast, J.C. 1961. The food of the bairdiella. pp. 153-64 in Walker 1961.
- Raimondi, P.T. 1992. Adult plasticity and rapid larval evolution in a recently isolated barnacle population. *Biological Bulletin* 182:210-20.
- Rico-Martinez, R. and S.I. Dodson. 1992. Culture of the rotifer *Brachionus plicatilis* Pallas. *Aquaculture* 105:191-9.

- Robinson, R.L., G.F. Turner, A.S. Grimm, and T.J. Pritcher. 1990. A comparison of the ingestion rates of three tilapia species fed on a small planktonic alga. *Journal of Fish Biology* 36:269-70.
- Rogers, F.L. 1949. Three new subspecies of *Balanus amphitrite* from California. *Journal of Entomology and Zoology* 41:23-32.
- St. Amant, J.A. 1966a. Addition of *Tilapia mossambica* Peters to the California fauna. *California Fish and Game* 52:54-5.
- St. Amant, J.A. 1966b. Progress Report of the Culture of *Tilapia mossambica* (Peters) Hybrids in Southern California. California Department of Fish and Game Inland Fisheries Administrative Report 66-9. 25 pp.
- Schlupp, I., C. Marler, and M.J. Ryan. 1994. Benefit to male sailfin mollies of mating with heterospecific females. *Science* 263:373-4.
- Schoenherr, A.A. 1979. Niche separation within a population of freshwater fishes in an irrigation drain near the Salton Sea, California. *Bulletin of the Southern California Academy of Science* 78:46-55.
- Schoenherr, A.A. 1985. Replacement of *Cyprinodon macularius* by *Tilapia zilli* in an irrigation drain near the Salton Sea. *Proceedings of the Desert Fishes Council* 13:65-6.
- Schoenherr, A.A. 1988. A review of the life history and status of the desert pupfish, *Cyprinodon macularius*. *Bulletin of the Southern California Academy of Science* 87:104-134.
- Smit, G.L., J. Hattingh and J.T. Ferreira. 1981. The physiological responses of blood during thermal adaptation in three freshwater fish species. *Journal of Fish Biology* 19:147-160.
- Snell, T.W. 1986. Effect of temperature, salinity and food level on sexual and asexual reproduction in *Brachionus plicatilis* (Rotifera). *Marine Biology* 92:157-62.
- Stauffer, J.R. 1986. Effects of salinity on preferred and lethal temperatures of the mozambique tilapia, *Oreochromis mossambicus* (Peters). *Water Resources Bulletin* 22:205-8.
- Stauffer, J.R., Jr., C.H. Hocutt and W.F. Goodfellow. 1985. Effects of sex and maturity on preferred temperatures: a proximate factor for increased survival of young *Poecilia latipinna*? *Archiv für Hydrobiologie* 103:129-132.
- Stickney, R.R. 1986. *Tilapia* tolerance of saline waters: a review. *Progressive Fish-Culturist* 48:161-7.
- Sumner, I.T., J. Travis, and C.D. Johnson. 1994. Methods of female fertility advertisement and variation among males in responsiveness in the sailfin molly (*Poecilia latipinna*). *Copeia* 1994:27-34.
- Sweet, J.G. and O. Kinne. 1964. The effects of various temperature-salinity combinations on the body form of newly hatched *Cyprinodon macularius* (Teleostei). *Helgolander Meeresuntersuchungen* 11:49-69.
- Todd, E.S. 1968. Terrestrial sojourns of the long-jaw mudsucker, *Gillichthys mirabilis*. *Copeia* 1968:192-4.
- Todd, E.S. and A.W. Ebeling. 1966. Aerial respiration in the longjaw mudsucker *Gillichthys mirabilis* (Teleostie: Gobiidae). *Biological Bulletin* 13:265-88.
- Uchida, R.N. and J.E. King. 1962. Tank culture of tilapia. *Fishery Bulletin* 62:21-52.
- USFWS. 1986. Determination of endangered status and critical habitat for the desert pupfish. *Federal Register* 51(61):10842-51.
- Vadstein, O., G. Oie, and Y. Olsen. 1993. Particle size dependent feeding by the rotifer *Brachionus plicatilis*. *Hydrobiologia* 255/256:261-7.
- Walker, B.W. (ed.). 1961. The Ecology of the Salton Sea, California, in Relation to the Sportfishery. *Fish Bulletin* No. 113. California Department of Fish and Game. 204 pp.
- Walker, B.W., R.R. Whitney, and G.W. Barlow. 1961. The fishes of the Salton Sea. pp. 77-91 in Walker 1961.
- Walker, B.W., R.R. Whitney, and L.H. Carpelan. 1961. General considerations and recommendations. pp. 185-192 in Walker 1961.
- Waters, M.R. 1983. Late Holocene lacustrine chronology and archaeology of ancient Lake Cahuilla, California. *Quaternary Research* 19:373-87.
- Weisel, G.F., Jr. 1947. Breeding behavior and early development of the mudsucker, a gobiid fish of California. *Copeia* 1947:77-85.
- Wetzel, R.G. 1983. *Limnology*, Second Edition. Saunders, New York. 858 pp.
- Whitney, R.R. 1961a. The bairdiella, *Bairdiella icistius*. pp. 105-151 in Walker 1961.

Whitney, R.R. 1961b. The orangemouth corvina, *Cynoscion xanthulus* Jordan and Gilbert. pp. 165-183 in Walker 1961.

Table 1. Surface flows into the Salton Sea, 1980-1990.

Table 2. Primary productivity (mg carbon/m²/day) of the Salton Sea and classification levels for fresh waters.

<u>Category</u>	<u>Productivity</u>	<u>Salton Sea</u>
Ultraoligotrophic	<50	
Oligotrophic	50-300	
Mesotrophic	250-1000	
Eutrophic	>1000	1032

Salton Sea data modified from Carpelan (1961a)
Productivity categories from Wetzel (1983)

Table 3. Phosphate-phosphorous and total nitrogen inflows to the Salton Sea.

	<u>Alamo R.</u>	<u>New R.</u>	<u>Other Imperial Valley Drains</u>	<u>Whitewater R.</u>	<u>Other Coachella Valley Drains</u>	<u>Total</u>
PO ₄ -P (mg/m ³)	682	906	?	1031	?	
Total N (mg/m ³)	9,745	6,794	?	13,693	?	
Water flow (AF/year)	600,000	450,000	100,000	100,000	50,000	1,300,000
Annual PO ₄ -P load (kilograms)	504,741	502,890		127,172		1,134,893
(tons)	556	555		140		1,251
Annual total N load (kilograms)	7,012,344	3,771,720		1,689,007		12,473,071
(tons)	7,730	4,158		1,862		13,750

Nutrient data from RWQCB, 1980-1992

Table 4. Comparison of Salton Sea phosphate-phosphorous and total nitrogen concentrations (mg/m³) with those from lakes of recognized trophic status.

<u>Classification</u>	<u>Phosphorous</u>			<u>Nitrogen</u>		
	<u>Mean</u>	<u>Range</u>	<u>Salton Sea</u>	<u>Mean</u>	<u>Range</u>	<u>Salton Sea</u>
Oligotrophic	8.0	3-18		661	307-1630	
Mesotrophic	26.7	11-96		753	361-1387	
Eutrophic	84.4	16-386	345	1875	393-6100	
Hypereutrophic		750-2000				13,700

Salton Sea data from RWQCB, 1980-1992 Classification data from Wetzel (1983)

Table 5. Annual areal phosphorous and nitrogen loading (g/m²) of the Salton Sea and recommended loading levels.

Mean Lake Depth	"Permissible"		"Dangerous"		Salton Sea	
	P	N	P	N	P	N
5 meters	0.07	1.0	0.13	2.0		
10 meters	0.10	1.5	0.20	3.0		
50 meters	0.25	4.0	0.50	8.0		
100 meters	0.40	6.0	0.80	12.0		
150 meters	0.50	7.5	1.00	15.0		15.4
200 meters	0.60	9.0	1.20	18.0	1.19	

Nutrient data from RWQCB, 1980-1992

Recommended levels from Wetzel (1983)

- Figure 1. Southern California, showing the location of the Salton Sea.
- Figure 2. The Salton Basin, the Gulf of California, and the Colorado River delta.
- Figure 3. Location of break in dikes allowing flow into Salton Basin (modified from Kennan 1917).
- Figure 4. Historic flows of the Gila River based on tree ring data from ____ 19__.
- Figure 5. Elevation and salinity of the Salton Sea, 1905-1994.
- Figure 6. Examples of four dinoflagellate (left) and three other genera of algae recorded from the Salton Sea.
- Figure 7. A live foraminiferan (left) showing the pseudopods extending through the calcium carbonate test, and the tests of three other genera recorded from the Salton Sea.
- Figure 8. The rotifers *Brachionus plicatilis*, carrying a single egg, and *Synchaeta*.
- Figure 9. An adult female cyclopoid copepod (carrying two egg sacs), a nauplius larva, and a harpacticoid copepod.
- Figure 10. The three life stages of *Neanthes succinea*.
- Figure 11. The three life stages of *Balanus amphitrite saltonensis*.
- Figure 12. A male desert pupfish, *Cyprinodon macularius*.
- Figure 13. A sargo, *Anisotremus davidsonii*.
- Figure 14. A gulf croaker, *Bairdiella icistia*.
- Figure 15. A mozambique mouthbrooder, *Tilapia mossambica*.
- Figure 16. An orangemouth corvina, *Cynoscion xanthulus*.
- Figure 17. A longjaw mudsucker, *Gillichthys mirabilis*.
- Figure 18. Female (top) and male (bottom) sailfin mollies, *Poecilia latipinna*.
- Figure 19. A porthole livebearer, *Poeciliopsis gracilis*.
- Figure 20. A threadfin shad, *Dorosoma petenense*.
- Figure 21. Trophic relationships among the biota of the Salton Sea.
- Figure 22. Possible changes in fish fauna with rising Salton Sea salinity.