

Welcome to the Conservation Lecture Series




Upcoming Lectures

- ❁ **Sierra Nevada Red Fox** – April 11, 1:00-3:00.
Presented by Dr. Ben Sacks
- ❁ **Alameda Striped Racer** – April 24, 1:00-3:00.
Presented by Karen Swaim
- ❁ **California Tiger Salamander** –April 28, 10:00 am-12:00. Presented by Dr. Chris Searcy
- ❁ **Shasta Crayfish** – April 29, 10:00-11:30.
Presented by Dr. Maria Ellis. **Location: Redding**
- ❁ **Desert Tortoise** – May 22 (time TBD). Presented by Becky Jones. **Location: Los Alamitos**
- ❁ **Amargosa Vole** – June 9, 1:00-3:00. Presented by Dr. Janet Foley and Dr. Robert Klinger


Questions?

Contact: margaret.mantor@wildlife.ca.gov




Yellow starthistle biology and control

Joe DiTomaso
UC Davis




Acres of western states infested with major invasive species

Species	Acres infested (x million)	% annual increase
Downy brome	56.0	14
Yellow starthistle	14.8	13-17
Canada thistle	7.1	10-12
Sericea lespedeza	5.5	24
Spotted knapweed	5.2	10-24
Musk thistle	4.7	12-22
Leafy spurge	3.7	12-16
Saltcedar	3.7	1-25
Medusahead	2.4	12
Perennial pepperweed	2.0	11-18
Diffuse knapweed	1.8	16
Russian knapweed	1.2	8-14




Distribution and Relative Abundance of Yellow Starthistle (*Centaurea solstitialis*) in the West - 2001 Survey




□ Borders Western States
Yellow Starthistle
□ Not Found
■ 1-100ac
■ 100-1000ac
■ >1000ac
■ Unreported


Data provided in 2001
by state weed coordinators
and compiled by Eric Lane,
Colorado Weed Coordinator







Available through the Cal-IPC.org website






UC Davis Weed Science Program
Copyright Regents, University of California
Photo by Joe DiTomaso



DiTomaso, Kyser and Piroosko. 2003. Weed Science 51, 334







*UC Davis Weed Science Program
Copyright Regents, University of California
Photo by Tom Lanini*




*UC Davis Weed Science Program
Copyright Regents, University of California
Photo by Joe DiTomaso*




Pattern of Total Soil Water content (cm/150 cm) Over Four Years





Total Soil Water Content by Season Averaged Across Years



Max-min water content

YST = 15.8 cm

WG = 11.8 cm

AG = 9.9 cm




Potential Water Cost Estimates

- ✪ YST maintained a drier soil profile by 6.6 cm H₂O compared to the annual grasses
 - ✪ $(0.066\text{m H}_2\text{O}) \times 10,000 \text{ m}^2/\text{ha} = 660 \text{ m}^3/\text{ha}$
 - 0.21 acre feet of water
 - $(\$12.00\text{-}30.00/\text{Acre ft}) = (\$2.58\text{-}6.47 \text{ lost})$
- ✪ A higher estimate by Gerlach (2000)
 - ✪ $(0.12\text{m H}_2\text{O}) \times 10,000 \text{ m}^2/\text{ha} = 1200 \text{ m}^3/\text{ha}$
 - 0.39 acre feet of water
 - $(\$12.00\text{-}30.00/\text{Acre ft}) = (\$4.72\text{-}11.70 \text{ lost})$
- ✪ Siskiyou County alone, 26.4 million gallons water lost/yr, equivalent to 15-25% of mean annual precipitation



Yellow starthistle
(Centaurea solstitialis)






Root chamber



**Root chamber dimensions: 1 m
wide on both sides X 2.4 m tall**
Sunken telephone booths




Root chamber



**June 3, 2005 –
counting yellow
starthistle roots.**






Root chamber



Senescing yellow starthistle in late summer




Root chamber



Red marks show amount of individual root growth.

Depths correlate with neutron probe soil moisture depths.






Neutron probes







Roots observed on the chamber wall were derived from lateral root development




Roots evenly distributed throughout profile




Roots primarily in upper profile






Next year under drought, YST produced 72% fewer roots compared to previous year


Under drought, *Elymus* produced 50% fewer roots compared to previous year




Roots primarily in upper profile





Roots primarily in lower profile







YST [CO₂] at 120 cm







YST soil water use at 180 cm






Overall






Three distinct stages of development in the life cycle of YST






Conclusions

- Years with little to no deep soil moisture recharge, roots remain in the shallow regions of the soil typically occupied by annual grasses, thereby intensifying competitive interactions.
 - YST would be functionally similar to annual grasses in drought years (also competes with perennials)
 - In normal or wet years would be functionally similar to deep-rooted perennial species, such as *E. glaucus*.
- *G. camporum* is an early-succession species and a good colonizer in newly restored or disturbed plant communities, but is not tolerant to further disturbance or competition with other grassland species in California.



Significance

- ✦ Control of yellow starthistle is most efficient a couple of weeks before plants reach bolting stage
 - ✦ No impact to water use
 - ✦ Greatest benefit to desirable competing species
- ✦ Biomass and reproductive output of plants already determined by the beginning of the flower stage
 - ✦ Can explain the efficacy of burning and mowing during the early flowering stages
 - ✦ Little opportunity for plants to recover after soil moisture content has been maximally reduced



Dittrichia graveolens (stinkwort)





Fig. 1. Approximate rate of spread of *Dittrichia graveolens* in California as represented by the number of California counties where plant collections have been made between 1984 and 2012 (Consortium of California Herbaria 2012).





Fig. 2. Chronological spread of stinkwort in California counties from 1984 to 2012.










Fig. 3. Life cycle of (A) stinkwort (*Dittrichia graveolens*), a late-season winter annual, compared to the life cycles of (B) wild mustard (*Sinapis arvensis* L.), a typical early-season winter annual, and (C) yellow starthistle (*Centaurea solstitialis* L.), a typical late-season winter annual.





Control options

- ❖ Mechanical
 - ❖ Handpulling, hoeing, weed whips
 - ❖ Tillage
 - ❖ Mowing
- ❖ Biological
- ❖ Cultural
 - ❖ Grazing
 - ❖ Prescribed burning
- ❖ Chemical
- ❖ Integrated approaches
 - ❖ Burning and chemical
 - ❖ Revegetation

Harrow



Disk

Mowing



*UC Davis Weed Science Program
copyright Regents, University of California
Photo by Joe DiTomaso*



Rotary mower



Flail mower

Growth form of YST affects mowing success




4" cutting ht

Soil surface




© Weed Science Program
The Regents, University of California
Joe DiTomaso

UC Davis Weed Science Program
Copyright Regents, University of California
Photo by Joe DiTomaso



Yellow starthistle rosette height under shadecloth and in full sun (ca. 100 days after germination)





UC Davis Weed Science Program
Copyright Regents, University of California
Photo by Joe DiTomaso




Hairy weevil
*Eustenopus
villosus*


UC Davis Weed Science Program
copyright Regents, University of California
Photo by Jack Kelly Clark


False
Peacock Fly
*Chaetorellia
succinea*





Biocontrol distribution and attack rates in California






Effect of insect bioagents on YST

Percentage of Heads			
Insect spp.	Cause of damage	Early season	Mid-season
		N=229	N=231
<i>Eustenopus villosus</i>	Feeding/ovipos.	72	9
	Larvae	17	3
<i>Chaetorellia succinea</i>	Larvae	26	41




Yellow starthistle biomass with simulated biocontrol or mowing

Treatment	Shoot wt (g)	Developed capitulum wt (g)	Undeveloped capitulum wt (g)	Total aboveground wt (g)
Untreated	384.3 ± 15.9 a	130.1 ± 30.1 a	1.05 ± 0.2 a	515.4 ± 15.9 a
Bud damage	390.4 ± 59.3 a	91.6 ± 22.5 ab	5.5 ± 0.9 b	487.4 ± 78.9 a
Mowed	108.9 ± 16.1 b	38.6 ± 9.7 b	0.65 ± 0.2 a	148.1 ± 25.3 b
<i>P > F</i>	0.001	0.04	0.008	0.0019



Number of seeds per plant with controls (C), mowed (M), and simulated biocontrol (D)




Rust Fungus

- ✦ *Puccinia jaceae* var. *solstitialis*
- ✦ Found in Turkey; Approved for released by CDFA in 2003
- ✦ Attacks vegetative part of plant

















Field Site














Grazing

Table 1. Comparison of grazing characteristics of cattle, sheep and goats (Frost and Launchbaugh 2003)

Animal	Digestive systems	Feeding behavior	Classification
Cattle	Large rumens adapted to ferment fibrous material	Best for managing fibrous herbaceous vegetation, prefer grasses but will also graze yellow starthistle at the bolting stage	Grass and roughage eaters
Sheep	Large rumen adapted to ferment fibrous material	Can selectively graze and tolerate high fiber content, diet dominated by forbs, will control yellow starthistle when grazed at bolting stage, but not in rosette stage	Forb and roughage eaters, more easily managed by human herders, used for strategic grazing
Goats	Large liver mass that allows processing of secondary compounds less digestible or more toxic to other grazers	Mouths designed to strip leaves from woody plants and chew branches, will also feed on yellow starthistle in the spiny stage	Browsers used often to control woody species



*UC Davis Weed Science Program
Copyright Regents, University of California
Photo by Craig Thomson*

Prescribed
burning









*UC Davis Weed Science Program
Copyright Regents, University of California
Photo by Joe DiTomaso*







Vegetative cover in July






Yellow starthistle seedbank recovery










Herbicide control








Herbicides on YST control




* LSD (P=0.05)




Transline (clopyralid) treated rangeland on right





Milestone and Transline on yellow starthistle control






Stage of growth at treatment stage




Aminopyralid 3 oz/A

Untreated






Aminopyralid and clopyralid on grass forage following year












*LSD (P=0.05)
**LSD (P=0.01)




Yellow starthistle control


Control 60 DAT					
Treatment	Timing	Rate		YST	Filaree
		oz ai/A	product/A		
aminocyclopyrachlor	rosette	0.5	0.625 oz	67 c	27 bc
aminocyclopyrachlor	rosette	1	1.25 oz	90 b	27 abc
aminocyclopyrachlor	rosette	1.5	1.875 oz	100 a	53 ab
aminocyclopyrachlor	rosette	2	2.5 oz	100 a	47 ab
aminocyclopyrachlor	rosette	2.5	3.125 oz	100 a	57 ab
aminocyclopyrachlor	rosette	3	3.75 oz	100 a	73 a
aminopyralid	rosette	0.75	3 fl oz	100 a	60 ab
aminopyralid	rosette	1.25	5 fl oz	100 a	33 abc
untreated	rosette	---	---	0 d	0 c







Late season control of YST



Selective way of using a non-selective herbicide




Rope wick
application of
Roundup
(glyphosate)

ATV or tractor
attachment with rope
wick applicator





Treated with acetic/citric acid combination




Integrated approaches

- ⊕ Burning and chemical
- ⊕ Revegetation and chemical
- ⊕ Mowing and chemical




San Benito site following burn in 1999




YST cover following two years of control

San Benito County



Yuba County





Ripgut brome (*Bromus diandrus*)




Primarily ripgut brome




Medusahead


Taeniatherum caput-medusae




Medusahead



Ripgut brome






Integrated management of YST at Ft. Hunter Liggett

Site	Treatment	Seedlings/m ²	
		Untreated	Treated (% untreated)
Military use			
2000	Burned 1999	117	271 (232%)
	Clopyr. 2000		
2001		478	2 (0.4%)
Wildland site			
2000	Burned 1999	435	547 (126%)
	Clopyr. 2000		
2001	Burned 2001	1560	6 (0.4%)
2002	No treatment	987	45 (5%)

Assuming 75% germination or loss / year with no new input







Brown Brush Monitor

Foreground: Annual grass community
Background: Perennial grass community





Late Season Yellow Starthistle Cover (summer 2000)






In *Centaurea solstitialis* Cover (%)





Native perennial grass (*Elymus glaucus*)



Black = *Grindelia*, perennial native forb

White = yellow starthistle

Light gray = annual species

Dark gray = *Elymus glaucus*, native perennial grass




Best timing of aminopyralid for control of yellow starthistle

		Yellow starthistle cover at each evaluation date			
		July 2010	May 3, 2011	August 1, 2011	May 23, 2012 ^a
Site	Time of application	%			
Mission	November 2009	17.3 b	32.3 b	52.8 a	0
	January 2010	0.3 c	1.3 c	11.7 b	0
	March 2010	0 c	0.1 c	0.7 b	0
	None	43.4 a	50.0 a	47.4 a	0
	Probability F	< 0.0001	< 0.0001	< 0.0001	—
Back	November 2009	20.8 b	15.0 b	89.4 b	0
	January 2010	0.3 c	0.6 c	12.9 c	0
	March 2010	0.2 c	0.2 c	4.6 d	0
	None	84.3 a	50.7 a	100.0 a	0
	Probability F	< 0.0001	< 0.0001	< 0.0001	—



Stipa cernua was the most successful seeded species to establish, but establishment required 3 yr. January or March aminopyralid treatment integrated with perennial grass drill seeding in January was most successful.

		<i>Stipa cernua</i> cover			
		Aminopyralid application timing			
		November	January	March	untreated
Year of evaluation	Drill seed timing	%			
2010	December	0 b	0 b	0.6 b	0 b
	January	0.6 b	2.7 ab	4.6 a	1.1 ab
	March	0 b	2.8 ab	0.3 b	0 b
	unseeded	0 b	0 b	0 b	0 b
2011	December	0 b	0 b	0 b	0 b
	January	0 b	0.6 ab	5.0 a	0.2 b
	March	0 b	0.6 ab	2.2 ab	0 b
	unseeded	0 b	0 b	0 b	0 b
2012	December	1.2 de	3.3 de	3.5 cde	1.0 de
	January	23.5 a	18.8 a	10.8 abc	13.2 ab
	March	4.0 bcde	6.8 bcd	2.0 de	3.7 cde
	unseeded	0 e	0 e	0 e	0 e



Summary of YST management

- ⊕ Numerous successful control options
 - ⊞ Timely mowing, grazing, burning, herbicides
- ⊕ Keys to long term success
 - ⊞ Deplete the seedbank
 - ⊞ Prevent new seed recruitment
 - Off site recruitment
 - Livestock, vehicles, wind
 - On site escapes
 - Skips, fringe areas, fencelines, satellite populations
 - ⊞ Monitoring and detection of YST
 - ⊞ Spot treatment or follow-up program

Thanks for your attention

Questions?

