

# Welcome to the Conservation Lecture Series



[www.dfg.ca.gov/habcon/lectures](http://www.dfg.ca.gov/habcon/lectures)

- See the schedule of upcoming lectures
- Register for lectures
- Watch videos of past lectures

Questions? Contact  
[margaret.mantor@wildlife.ca.gov](mailto:margaret.mantor@wildlife.ca.gov)



# Lectures in April

- **Cactus Wren, Dr. Kristine Preston**  
April 17, 1:00-3:00, **San Diego**
- **Alameda Striped Racer, Karen Swaim**  
April 24, 1:00-3:00, Sacramento
- **California Tiger Salamander, Dr. Chris Searcy**  
April 28, 1:00-3:00, Sacramento
- **Shasta Crayfish, Dr. Maria Ellis**  
April 29, 10:00-11:30, **Redding**

# California's native and nonnative red foxes



photo by K. Aubry, USFS




photo by A. Rich, USFS



photo by B. Sacks

Benjamin N. Sacks

**Mammalian Ecology and Conservation Unit**

Center for Veterinary Genetics

Dept. Population Health and Reproduction



# Overview

- Biogeography
- California's 3 red foxes
- Population trends and status
- Sacramento Valley red fox research
- Sierra Nevada red fox research

# Biogeography and taxonomy

- Evolved in Africa and Eurasia
- Fossil record: arrival in North America 0.5–0.3 MYA
- Until 1959, was considered *Vulpes fulva* in NA
- Currently considered conspecific with Eurasian *V. vulpes*



# DNA as a tool

## Mitochondrial

- Cytoplasmic
- Maternal inheritance
- 16,000+ base-pairs
- 100% inherited together
- Mutations accumulate over time (molecular clock)
- Great for reconstructing ancestry (maternal evolutionary trees)
- but can be misleading...

## Nuclear

- Nucleus (chromosomes)
- Biparental inheritance
- ~3 billion base-pairs (x2)
- Randomly inherited genes
- Each gene has different ancestral history
- Statistical averaging necessary (species trees)
- Great for population assignment, parentage, individual fingerprints, etc.




# Mitochondrial tree

Eurasia and  
NW North America


North America Only

Clades separated by  
300,000 years




Aubry et al. 2009

# Native Red Foxes



(slide by Marcelle Moore)


# Native Red Foxes






# Native Red Foxes

90 - 50 KYA




# Native Red Foxes

10 KYA





# Native Red Foxes



Aubry et al. 2009

# North American red foxes could be distinct species from Eurasian/African ones

## Nuclear gene tree



Statham et al. in review



# California's 3 red foxes



*photo by B. Sacks, Sacramento Valley red fox, Yolo County*



*photo by A. Rich, USFS, Sierra Nevada red fox, Tuolumne County*



*photo by Jim Wiley, San Francisco Bay area non-native*

# Sacramento Valley?

- Present since 1880
- VERY different habitat
- Origin?
- Roest: railway from east





*(Range map approximate only; do not reference)*






# Conventional wisdom

“Only the montane red fox is native to California. In the nineteenth century a population of the eastern red fox was introduced into the lowlands of the state.”




“In about 1885...this species became established in the Sacramento Valley...almost certainly introduced from eastern North America...have now spread widely...”

# Conventional view (1999)



Lewis, J.C., K.L. Sallee, and R.T. Golightly, Jr. 1999. Introduction and range expansion of nonnative red foxes (*Vulpes vulpes*) in California. *American Midland Naturalist* 142:372-381.






Perrine et al. 2007  
Sacks et al. 2010a

Historical (1850-1930)

Modern (2000-2008)




- “Mountain” mtDNA
- Eastern/Alaskan mtDNA



Perrine et al. 2007  
Sacks et al. 2010a




# “Mountain mtDNA”



Perrine et al. 2007  
Sacks et al. 2010a





Perrine et al. 2007  
Sacks et al. 2010a





# Native subspecies of western red foxes

Washington Cascades red fox  
(*V. v. cascadensis*)

Rocky Mountain red fox  
(*V. v. macroura*)

(Range map approximate only; do not reference)

Sacramento Valley red fox  
(*V. v. patwin*)



Sierra Nevada red fox  
(*V. v. necator*)



Sacks et al. 2010a

# Lowland red foxes revisited

Lewis, J.C., K.L. Sallee, and R.T. Golightly, Jr. 1999.  
*American Midland Naturalist* 142:372-381.




Moore 2009

Sacks et al. 2010b, 2011



# Distribution of “pure” (>90%) foxes




- Native genome
- Non-native genome

Moore 2009  
Sacks et al. 2011



# Distribution of hybrids



- Native ancestry
- Non-native ancestry

Moore 2009  
Sacks et al. 2011

# The two native California red foxes are not the same




Photo: B. Sacks, Sacramento Valley RF, Yolo Cty, CA



Photo: B. Freund, Sierra Nevada RF, Oregon Cascades

# SVRF foxes larger



Sacks et al. 2010b





Photo by M. Statham

Sacramento Valley red fox  
Jan 2011



SNRF D110051

Photo: Mourad Gabriel


Sierra Nevada red fox  
Jan 2011

# Population trends and status




# Genetic diversity decline


Genome wide



mtDNA



Sac Valley



# How many Sacramento Valley red foxes today?

- Phase I: Develop species distribution model
  - Determine habitat use relative to availability
  - Apply Maxent approach to predictive modeling
  - Validate with independent data
- Phase II: Assess occupancy in predicted habitat
  - Select “random” subsample of model-predicted habitat to survey
  - Assess red fox presence
    - Baited camera traps
    - Scat searches
  - Factor in detection probability to estimate occupancy fraction
- Occupancy fraction \* predicted habitat area/home range size = estimated number of breeding pairs



# Phase I

# Burrowing Owl

Sports UC Davis offensive lineman waits for second day of NFL Draft

You can help  
Report fox sightings, or  
the locations of dens or  
road kills, online at

## UCD: Valley foxes are genetically distinct

# The Sacramento Bee REGIONAL DIGEST

### UCD prof seeks help with red foxes

DAVIS – A University of California,

The study is important because  
evidence in California are  
so badly "hammered" en-  
cies that some biologists have  
to be culled.  
ary evidence in the study  
suggest an evolutionary rela-  
tion foxes here and the Sierra  
a rare subspecies at home  
See FOXES, Page A11



Yolo Audubon Society  
Vol. 36 No. 9 May-June 2007

## Rethinking the Red Fox in the Sacramento Valley

A biologist

<b>UC DAVIS</b>	<b>News &amp; Information</b>	This service is provided by UC Davis News Service, 530-752-1930
6.6.2007	Search/Archives Facts & Figures	UC Davis Experts Seminars/Events

### Current News

- Agriculture
- Business/Government/Law
- Campus News
- Engineering
- Health
- Science News
- Social Sciences/Arts/Humanities
- Veterinary Medicine

### UC Davis in the News

- Publications
- UC Davis Magazine
- Dateline UC Davis
- California Aggie
- Broadcast
- NewsWatch
- Frontiers
- Special Programming
- UCTV

### Multimedia

### Related News

- UC Davis Athletics
- UC Davis Health System
- UC News Wire

### News Service Resources

### University Communications

### Native Red Foxes Living in Valley; Public's Help Sought

May 29, 2007

Contrary to onetime scientific opinion, red foxes living in the Sacramento Valley are not an artificially introduced species that threaten local ecosystems, according to a UC Davis researcher.

Ben Sacks, an expert in canine genetics and a researcher in the School of Veterinary Medicine, has new findings suggesting that these animals are natives, and close cousins to the native Sierra Nevada red fox.

In other low-elevation parts of the state, red foxes are indeed nonnative, invasive and threaten several endangered bird species. Some ecologists and biologists have called for the nonnative animals to be killed in these areas to preserve the delicate balance of those ecosystems.

Sacks, who has been studying the foxes for the past 10 years, is reaching out to the public for help in reporting all sightings of red foxes in the Sacramento Valley and in the high elevations of the Sierra Nevada and Cascades mountains. He is requesting that anyone who sees any evidence of red foxes in the area, either alive or dead (such as roadkill), tell him about it online.

"It's important that we collect public data as well as continue our own research," said Sacks. "Getting this well-rounded stream of information will help us learn more about how the Sacramento Valley red fox interacts with its local environment."

Specifically, Sacks and his team are interested in understanding how the Sacramento Valley foxes, whose ancestors evolved in cold mountainous climates, adapted to the warm flatlands of the valley.

Sacks and the School of Veterinary Medicine have set up a Web site where the public can report their sightings: <http://foxsurvey.ucdavis.edu>. The site also has helpful tips and photos for distinguishing red foxes from gray foxes and coyotes.

#### Additional information:

- [Fox survey](#)

#### Media contact(s):

- Ben Sacks, School of Veterinary Medicine, (530) 754-9088, [bensacks@ucdavis.edu](mailto:bensacks@ucdavis.edu)
- Sylvia Wright, UC Davis News Service, (530) 752-7704, [swright@ucdavis.edu](mailto:swright@ucdavis.edu)

E-mail this story  
Printable version



This adult red fox, a California native, was photographed hunting ground squirrels in a freshly disked field near Woodland, Calif. (Ben Sacks/UC Davis photo)



Students and Field/lab techs: Marcelle Moore, Karen Converse  
PIs: BN Sacks/HU Wittmer/MJ Statham  
CDFW cooperators: Armand Gonzales  
Funding: USFWS/CDFW/UC Davis CPB/Sac State



# Sacramento Valley Fox Survey

[Home](#) [Report a Sighting](#) [Fox or Coyote?](#) [Status](#)

Contact: [sacvalleyredfox@gmail.com](mailto:sacvalleyredfox@gmail.com)

## Red Fox Sighting Report

Please report road-kills immediately to Ben Sacks, 530-754-9088

Digital photos appreciated

### Animal

Species	Tail	Ears	Body
<input type="radio"/> Red Fox	<input type="radio"/> Did Not See	<input type="radio"/> Did Not See	<input type="radio"/> Mostly Orange
<input type="radio"/> Gray Fox	<input type="radio"/> White Tip	<input type="radio"/> Backside Black Tip	<input type="radio"/> Mostly Gray
<input type="radio"/> Coyote	<input type="radio"/> Black Tip	<input type="radio"/> Backside Orange Tip	
<input type="radio"/> Other <input type="text"/>			
Confidence: <input type="text" value="Choose"/>			

### Date and Time

How Recently:  or Month/Year:

Time of Day:

### Location

Description:  
(distance and direction to nearest town/intersection)


County:

Click on the map to fill in coordinates.

Latitude:

Longitude:


Terrain:



# Search near sighting reports to confirm occurrence








- Road kills
- Den sites


Sacks et al. 2010b

Tuffly, M., and A. Kilgore. 1998. California Central Valley Wetlands and Riparian GIS Data Sets. California Department of Fish and Wildlife. <http://www.dfg.ca.gov/biogeodata/gis/clearinghouse.asp>







# Selection index



Sacks et al. 2010b

Sacks et al., unpublished distribution models






# Phase II (in progress)

- Stratified based on model
- Selected hexes within strata at random
- Seek permission to survey on private land
- Survey
  - Baited camera traps
  - Scat collection and DNA analysis

Project leader: **Kat Miles** (MS student)  
Field/lab techs: Preston Alden, Michelle Holtz, Zach Lounsberry  
PI: BN Sacks  
CDFW cooperators: D. Wright, C. Nguyen, S. Hemingway,  
K. Converse  
Funding: USFWS/CDFW/UC Davis











2014-02-16 8:05:57 PM M 2/3

47°F



H0898

RECONYX

2013-04-21 4:15:01 AM M 1/3

54°F



H0890

RECONYX



# Preliminary findings

- Presence detected in 3/3 positive control hexes
- Presence detected in 0/4 predicted-absence hexes
- Presence detected in 1/7 predicted-presence sites(!!)







# Sierra Nevada red fox

- “Threatened” under California ESA 1980
- Petitioned for listing under the US ESA 2011
- Causes for decline unclear
  - Trapping?
  - Coyote encroachment?
  - Habitat transformation (fire, timber, livestock)?
  - Prey declines?
  - Climate change?
- Factors limiting recovery
  - Inbreeding depression?
  - Less frequent high-prey years?
  - Recreational/other uses, etc.?








1993 re-discovered (Kucera)  
1998-2002 telemetry/photo/genetics  
2007-present photo/genetic monitoring

2010 re-discovered  
2010-present photo/genetic monitoring





Editor

2010

DOC

TT



# Southern Sierra Nevada red fox working group



**UC DAVIS**  
UNIVERSITY OF CALIFORNIA

VETERINARY GENETICS LABORATORY




Home Services Forensics Research Informatics About Contact  Search

*Mammalian Ecology and Conservation Unit*



**CAL POLY**  
SAN LUIS OBISPO



# Two efforts initiated

- Broad survey (Chris Stermer, CDFW)
- Focal study of re-discovered population (UCD, CDFW, USFS)
  - Size, extent, status of population
  - Snowmobile impacts?
  - Marine training impacts?

## FOCAL STUDY

Project leader: Cate Quinn (PhD student)

Field/lab techs: Preston Alden, David Wolfson, Kat Miles

PI: BN Sacks

CDFW collaborators: E. Burkett, C. Stermer

USDA-FS collaborators: A. Rich, S. Lisius, R. Mazur, J. Lowden

Funding: OHV/USDA-FS/USFWS/CDFW/UC Davis



# Methods

- Telemetry
- Noninvasive
  - Remote cameras paired with hair snaggers (for DNA)
  - Scat collection (for DNA)
  - Snowtracking





# Trapping winter 2014





SNRF 10







SNRF CAM3







SNRF CAM5





SNRF07







SNRF TRAP2





# What next?



photo courtesy of Patrick Cross



# Noninvasive genetic monitoring

- **Sources of DNA**
  - Hair (baited snaggers)
  - Scats (foot-searches)
  - Urine (Snow-tracking)
- **Analysis of DNA samples**
  - Mitochondrial
    - Identify species
    - Native/nonnative haplotypes
  - Nuclear microsatellites/sex markers
    - Compare to population database
    - Identify sex and individual
    - Familial relationships
- **Outcomes**
  - Individual timeline, longevity
  - Immigration/births
  - Home ranges







M2011-5

?

South

F-Roadkill



?

Sonora Pass



2011-3

F2011-2

F2010-6

M2011-1

F2010-7



Aug-10

Oct-10

Dec-10

Feb-11

Apr-11

Jun-11

Aug-11


Oct-11




•

•









# Demography

- Adults live a long time
- Few offspring are detected
- Reproduction or neonatal/juvenile survival limits population
- Also appears true in the Lassen population (Figura/Sacks, unpub.)
- Explanations: inbreeding depression? Nutrition? Predation?



(14 microsatellite loci)



(population assignment based on 33 microsatellite loci)


$H_0$ : Individual genotype originated from population

Probability ( $P$ ) of these (or more unusual) genotypes if  $H_0$  is true:

Fox ID	East Coast	Nonnative California	Rocky Mountains	Nevada	Lassen	Sac Valley	Washington Cascades	Sonora Pass
F2010-6	***	***	**	**	***	***	***	0.59
F2010-7	***	***	**	**	***	***	***	0.90
M2011-1	***	***	**	***	***	***	***	0.72
F2011-2	***	***	***	**	***	***	***	0.62
M2011-5	***	***	**	**	***	***	***	0.67
2012-3	***	***	**	**	***	***	***	0.56
F2010-8	***	***	***	**	***	***	***	0.57
M2012-1	***	***	**	0.03	***	***	***	**
M2012-2	***	***	**	0.02	***	***	***	***

\*\* $P \leq 0.01$

\*\*\* $P \leq 0.001$



0.5 1 2 3 4 Kilometers




# Outcomes/concerns


- Failure to survive and reproduce
- Competition with native foxes (esp males)
- Nonnative hybridization and introgression
  - Temporary genetic “rescue”
  - Outbreeding depression/swamping
  - Extirpation

# Last year

- ♀
- ♂
- unknown
- nonnative



# Last year







Parentage tests confirm hybrid pairings in both cases

## LITTER 1

## LITTER 2



# Sierra Nevada red fox

- Small number of individuals
- Inbreeding depression a concern
- Genetic swamping could be a bigger one
- What can we do?
  - Remove nonnative males?
  - Translocations of other native foxes?
  - Nothing (monitor)
    - Hope for the best
    - Learn from the worst

# California red fox recap

- Sacto Valley and SN red foxes are native and have declined
- Nonnative RFs have increased (encroachment minimal)
- Status of Sacto Valley red foxes uncertain (stay tuned)
- Sierra Nevada red foxes
  - Extremely rare
  - Reproduction/recruitment limited
  - Recently exposed to outbreeding
    - “Rescue” from inbreeding depression (at least in the short term)?
    - Swamp existing populations with non-locally adapted genes?
- Recommendations for SNRF
  - Continue noninvasive genetic monitoring
  - Capture and collar critical to study and management
  - Consider and plan translocations





# Primary co-investigators and facilitators

- Worldwide red fox

- Mark Statham
- Keith Aubry
- Sam Wisely

- Sac Valley red fox

- Marcelle Moore\*
- Karen Converse\*
- Kathleen Miles\*
- Heiko Wittmer
- Armand Gonzalez
- David Wright
- Canh Nguyen
- Stacy Hemingway




(non-fox)

- Sierra Nevada red fox

- Cate Quinn\*
- Jocelyn Akins\*
- Preston Alden\*
- David Wolfson
- Chris Stermer
- John Perrine
- Esther Burkett
- Adam Rich
- Sherri Lisius
- Pete Figura



\*Graduate students  
(underlined = current research coordinators)



# Acknowledgements

-Funders: CA OHV, USDA/FS, CDFW, USFWS, UC Davis-CPB, VGL

-Additional partners


- Rachel Mazur (USDA/FS, R4)
- JoAnne Lowden (USDA/FS, R4)
- Andrew Irvin (USMC)
- Greg Gerstenberg (CDFW, R4)
- Tim Taylor (CDFW, R6)
- Lily Douglas (USDI/BLM)
- Lindsay Cline (NPS)
- Rich Callas (CDFW, R1)\
- Kristy Fien (CDFW, GIS)
- Maureen Easton (USDA/FS-R4)
- Diana Craig (USDA/FS-R5)
- Diane McFarlane (USDA/FS-R5)
- Peggy O'Connell (USDA/FS-R5)
- Keaton Norquist (USDA/FS-R5)
- CSERC volunteers
- MANY, MANY fantastic** undergraduate interns and volunteers from UC Davis, Sac State Univ, and American River College







Questions?





# Further reading

- Aubry, K. A., M. J. Statham, B. N. Sacks, J. D. Perrine, and S.M. Wisely. 2009. Phylogeography of the North American red fox: vicariance in Pleistocene forest refugia. *Molecular Ecology* 18: 2668–2686.
- Converse, K.E. 2012. Genetic mating system and territory inheritance in the Sacramento Valley red fox. M.S. thesis, California State University, Sacramento, California.
- Moore, M.M. 2009. Impacts of encroaching non-native red foxes on the native Sacramento Valley red fox. M.S. thesis, California State University, Sacramento, California.
- Perrine, J. D., J. P. Pollinger, B. N. Sacks, R. H. Barrett, and R. K. Wayne. 2007. Genetic evidence for the persistence of the critically endangered Sierra Nevada red fox in California. *Conservation Genetics* 8: 1083-1095.
- Sacks, B. N., Moore M., Statham, M. J., Wittmer H. U. 2011. A restricted hybrid zone between native and introduced red fox (*Vulpes vulpes*) populations suggests reproductive barriers and competitive exclusion. *Molecular Ecology* 20:326-341.
- Sacks, B. N., M. J. Statham, J. D. Perrine, S.M. Wisely, Aubry, K. A. 2010a. North American montane red foxes: expansion, fragmentation, and the origin of the Sacramento Valley red fox. *Conservation Genetics* 11:1523-1539.
- Sacks, B. N., Wittmer H. U. and Statham, M. J. 2010b. *The Native Sacramento Valley red fox*. Report to the California Department of Fish and Game, May 30, 2010, 49 pp
- Sacks, B.N. In Press. Sacramento Valley red fox, *Vulpes vulpes patwin*. Species Account in California Mammal Species of Special Concern, California Department of Fish and Game.
- Statham, M. J., Sacks, B. N., Aubry, K. A., Perrine, J. D., Wisely, S.M. 2012. The origin of recently established red fox populations in the contiguous United States: Translocations or natural range expansions? *Journal of Mammalogy* 93:52-65.
- Statham, M. J., Rich, A. C., Lisius, S. K., Sacks, B. N. 2012. Discovery of a remnant population of Sierra Nevada red fox (*Vulpes vulpes necator*). *Northwest Science* 86:122-132