

TABLE 6a. 2000 Reported Deer Kill by CDF&G Region, Sex, Antler Class and Percent by Antler Class of Bucks (a).

CDF&G Region of Kill	Total Region Doe Kill	Unclassified Point Count (b)		Spike Bucks		2 Point Bucks		3 Point Bucks		4 Point Bucks		> 4 Point Bucks		Total Region Buck Kill	TOTAL REGION DEER KILL
		Region Buck Kill	% of Region Buck Kill	Region Buck Kill	% of Region Buck Kill	Region Buck Kill	% of Region Buck Kill	Region Buck Kill	% of Region Buck Kill	Region Buck Kill	% of Region Buck Kill	Region Buck Kill	% of Region Buck Kill		
Region 1 - Northern California and North Coast		8	0.1%	1	0.0%	3,602	45.9%	2,759	35.1%	1,296	16.5%	184	2.3%	7,850	7,850
Region 2 - Sacramento Valley and Central Sierra	9	4	0.1%	3	0.1%	1,941	53.8%	1,067	29.6%	507	14.1%	86	2.4%	3,608	3,617
Region 3 - Central Coast	71	15	0.3%	3	0.1%	2,984	61.5%	1,390	28.7%	413	8.5%	46	0.9%	4,851	4,922
Region 4 - San Joaquin Valley and Southern Sierra		5	0.2%		0.0%	1,512	55.6%	771	28.3%	351	12.9%	82	3.0%	2,721	2,721
Region 5 - South Coast	212	41	3.1%	15	1.1%	848	63.1%	329	24.5%	96	7.1%	14	1.0%	1,343	1,555
Region 6 - Eastern Sierra and Inland Deserts	4	5	0.5%		0.0%	475	42.9%	376	33.9%	221	19.9%	31	2.8%	1,108	1,112
STATEWIDE TOTALS:	296	78	0.4%	22	0.1%	11,362	52.9%	6,692	31.2%	2,884	13.4%	443	2.1%	21,481	21,777

TABLE 6a NOTES:

(a) Table 6a does not include PLM kill data. PLM kill analysis is included in TABLES 7a through 7f.

(b) Unclassified Point Count may be bucks without a reported point count or bucks with no points (antlerless and either-sex hunts).