TABLE 6a. 2002 Reported Deer Kill by CDF&G Region, Sex, Antler Class and Percent by Antler Class of Bucks (a).

		Unclassified Point Count (b)		Spike Bucks		2 Point Bucks		3 Point Bucks		4 Point Bucks		> 4 Point Bucks			
CDF&G	Total		% of		% of		% of		% of		% of		% of	Total	TOTAL
Region	Region	Region	Region	Region	Region	Region	Region	Region	Region	Region	Region	Region	Region	Region	REGION
of	Doe	Buck	Buck	Buck	Buck	Buck	Buck	Buck	Buck	Buck	Buck	Buck	Buck	Buck	DEER
Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	Kill	KILL
Region 1 - Northern California and North Coast		32	0.5%			2,968	47.0%	2,116	33.5%	1,022	16.2%	181	2.9%	6,319	6,319
Region 2 - Sacramento Valley and Central Sierra	59	10	0.3%	3	0.1%	1,678	51.3%	990	30.2%	494	15.1%	99	3.0%	3,274	3,333
Region 3 - Central Coast	65	20	0.5%	13	0.3%	2,543	61.8%	1,104	26.8%	390	9.5%	45	1.1%	4,115	4,180
Region 4 - San Joaquin Valley and Southern Sierra	3	4	0.2%			1,251	56.4%	618	27.9%	285	12.8%	60	2.7%	2,218	2,221
Region 5 - South Coast	141	31	3.6%	4	0.5%	567	66.1%	193	22.5%	55	6.4%	8	0.9%	858	999
Region 6 - Eastern Sierra and Inland Deserts	8	2	0.3%	6	0.9%	278	40.8%	223	32.7%	141	20.7%	31	4.6%	681	689
STATEWIDE TOTALS:	276	99	0.6%	26	0.1%	9,285	53.2%	5,244	30.0%	2,387	13.7%	424	2.4%	17,465	17,741

TABLE 6a NOTES:

2002TABLE6a.xls Page 1 of 1

⁽a) Table 6a does not include PLM kill data. PLM kill analysis is included in TABLES 7a through 7f.

⁽b) Unclassified Point Count may be bucks without a reported point count or bucks with no points (antlerless and either-sex hunts).