

TABLE 6a. 2003 Reported Deer Kill by CDF&G Region, Sex, Antler Class and Percent by Antler Class of Bucks (a).

CDF&G Region of Kill	Total Region Doe Kill	Unclassified Point Count (b)		Spike Bucks		2 Point Bucks		3 Point Bucks		4 Point Bucks		> 4 Point Bucks		Total Region Buck Kill	TOTAL REGION DEER KILL
		Region Buck Kill	% of Region Buck Kill	Region Buck Kill	% of Region Buck Kill	Region Buck Kill	% of Region Buck Kill	Region Buck Kill	% of Region Buck Kill	Region Buck Kill	% of Region Buck Kill	Region Buck Kill	% of Region Buck Kill		
Region 1 - Northern California and North Coast		5	0.1%			2,309	45.0%	1,858	36.2%	823	16.0%	134	2.6%	5,129	5,129
Region 2 - Sacramento Valley and Central Sierra	58	12	0.4%	5	0.2%	1,514	49.7%	929	30.5%	488	16.0%	101	3.3%	3,049	3,107
Region 3 - Central Coast	68	15	0.4%	3	0.1%	2,099	58.9%	1,068	30.0%	335	9.4%	45	1.3%	3,565	3,633
Region 4 - San Joaquin Valley and Southern Sierra	1	2	0.1%			1,087	53.3%	608	29.8%	282	13.8%	59	2.9%	2,038	2,039
Region 5 - South Coast	161	10	1.2%	5	0.6%	560	64.7%	207	23.9%	78	9.0%	5	0.6%	865	1,026
Region 6 - Eastern Sierra and Inland Deserts	4	1	0.1%			275	38.6%	262	36.8%	152	21.3%	22	3.1%	712	716
STATEWIDE TOTALS:	292	45	0.3%	13	0.1%	7,844	51.1%	4,932	32.1%	2,158	14.1%	366	2.4%	15,358	15,650

TABLE 6a NOTES:

(a) Table 6a does not include PLM kill data. PLM kill analysis is included in TABLES 7a through 7f.

(b) Unclassified Point Count may be bucks without a reported point count or bucks with no points (antlerless and either-sex hunts).