

# **Environmental Impacts Associated with Construction and Operation of Evaporation Ponds to Control Salinity of the Salton Sea**

Doyle Stephens, USGS  
DRAFT-June 30, 1999

## **Introduction**

The Bureau of Reclamation is evaluating the feasibility of constructing two evaporation ponds having a total surface area of 34.3 square miles at the southwest shore of the Salton Sea. The total capacity of the pond complex would be 245,226 acre feet. Water would be pumped from the Salton Sea into the ponds where it would evaporate, concentrating the salts. Inflow to the Sea is less saline and would replace water pumped to the ponds creating a net loss of salt from the Sea. Earlier variations of this alternative appeared in Bureau of Reclamation (1997, 1998) and involved construction of in-lake dikes isolating areas ranging from 30 to 127 square miles. Due to seismic activity in the area, dikes required to partition large sections of the Sea would have to be built using coffer dams that would need a seismic stability of Richter 5?? Dikes for shallow and smaller ponds could be constructed without coffer dams.

## Description of the ponds

A south pond with an approximate area of 20? acres would be constructed north of the inflow of the New River to a point immediately south of San Felipe Creek. A west pond with an area of 15 acres would be constructed about 3 miles north of San Felipe Creek to Salton City (figure 1). The mouth of San Felipe Creek would not be obstructed to allow Desert pupfish access to the backwater areas. Earthen dikes for both ponds would be constructed along the -250 foot contour of the Sea. Operating elevation of the ponds would be -227 foot, and maximum water depth at the dike would be about 23 feet. The pond bottom would slope shoreward and water depth would be only several inches at the shoreward extreme. The ponds would be unlined.

## Construction of the dikes

Organic-rich muck covers about 750 acres of the Sea bottom where the dike footing would be placed. The material is not structurally stable and would be removed to a depth of 5 feet using a suction dredge. Approximately 7 million cubic yards of muck would be removed. The material would be returned to the Sea between parallel silt curtains suspended by floats from the water surface. The silt curtain would allow water to pass but would contain the solids in a path of unknown width on the Sea bottom. The base area for the dike would be compacted and fill added to bring the dike top elevation to -220 feet. The outer (seaward) bank of the dike would be armored with stone riprap. The top surface of the dike (30 feet wide) would be graded to create a road for access and maintenance. The base of the dike would be 300 feet wide with a finished side slope of 3.5:1. Dikes would be constructed from the shore outward, so coffer dams would not be required.

An estimated 21 million cubic yards of fill and 440,000 cubic yards of riprap would be obtained from an area located west of Salton Sea Beach within the Torres-Martinez Indian Reservation. The Tribe owns mineral rights in the area and is desirous of selling the material. A haul road would be constructed to facilitate movement to the construction site. Due to the massive amounts of fill and riprap required, construction of the ponds would require about 4 years.

### Operation of the Ponds

Spreadsheets simulating pond operation prepared by Paul Weghorst (Bureau of Reclamation, Denver) assume reduction of current inflows to the Salton Sea to 1.06 million acre feet annually and increases in salinity of the Sea to 50,000 ppm by the time the ponds are ready for operation. Under these conditions, the ponds would initially contain 16,671,500 tons of salt. Between 97,150 and 115,700 acre feet of water would be pumped annually from the Sea into the ponds to replace water lost through evaporation. Annual salt removal by the ponds would be about 6.9 to 8.0 million tons. Annual inflow of salt to the Salton Sea would be about 4.03 million tons. The concentration of salt in the ponds would approach 63,000 ppm within about 23 years. After an arbitrary period of 30 years, chosen as a likely period of stability to earthquakes, pumping would discontinue and the ponds would be allowed to dry. The dikes would be reinforced and the ponds abandoned. Current plans do no call for capping to isolate the salt block in the ponds.

### **Potential Environmental Impacts**

Impacts identified during a workshop held at San Diego State University on June 17 are identified as bullet items below. A list of participants appears at the end of this document.

#### **Borrow pit and haul road**

- Presence of sensitive or T and E species in area

Development of a borrow area to provide 21 million cubic yards of fill would require literature search for the presence of sensitive species and threatened or endangered species in the area. This has been done by Tetra Tech using a computerized database maintained by the California Department of Game and Fish. If threatened or endangered species were present, Section 7 consultation with the US Fish and Wildlife Service may be required in the planning stage.

- Presence of cultural resources in the area

A survey should be done of cultural resources that may be located in the area of the borrow pit and haul road. Since the potential site for the pit is located on land owned by the Torres-Martinez Tribe, this may have already been done. Tetra Tech indicated they would pursue this information.

- Presence of wetlands or permanent streams

The presence of any wetlands or permanent streams in the area should be determined and mitigation may be required if damage done to these resources. A

potential problem of pit construction is inflow of surface water or ground water that may inhibit removal of material and could attract waterfowl to the pit site.

- Use of 100 yd<sup>3</sup> trucks on private road vs conventional trucks on hiway

Use of special 100 yd<sup>3</sup> trucks would require 273,000 truck trips to move the 21 million yd<sup>3</sup> of fill (This assumes 30% more material required to all for compaction when placed). Movement of the 440,000 yds<sup>3</sup> of riprap would require 4,400 truck trips. Assuming a three year active hauling period, about 250 truck trips per day would be required. As trucks of this capacity are not permitted on public highways, a separate haul road would be required for the 3 year period of movement of material. Use of conventional (10 yd<sup>3</sup>) trucks would require 10 times as many truck trips.

- Presence of contaminants in borrow fill material used for dikes

The tribal land under consideration as a source of dike fill and riprap has been surveyed for geologic potential, particularly for precious metal production (Toby Mancuso, Mancuso Resource Development Services, oral commun. June 1999). While there may be gold present in part of the deposit, the fill material and riprap could be removed without affecting the metal deposits. The riprap would be removed from a Jurassic deposit of quartz monzonite. Most of the fill material is derived from weathering of the monzonite and is of recent origin (Quaternary). Concentrations of leachable trace metals and metaloids such as selenium are expected to be quite small. Detailed geochemistry of the ore deposit is believed available from the Bureau of Indian Affairs in Denver.

- Compaction of haul road and need for restoration

This is being addressed by Tetra Tech.

### **Construction Impacts- Pond construction**

- Return of 7 million yds<sup>3</sup> sludge to the Salton Sea- turbidity, nutrient or trace element release, blanket of fish habitat, dissolved oxygen demand or hydrogen sulfide release

The most comprehensive inventory of the quality of bottom sediment in the Salton Sea was done by Levine-Fricke Reconnaissance in 1999. Surficial sediment samples taken with a box corer were collected from several sites within the proposed area of the evaporation ponds: sites 37-40 and 43-46. Three cores (8 to 10 feet depth) were taken southwest of the New River delta. Concentrations of trace elements in these samples did not exceed the lower effect range given in Long et al. (1995) except for selenium at one site. The sample from site 45, several miles east of the San Felipe alluvial area contained a selenium concentration of 0.9 mg/kg, 0.2 mg/kg greater than the lowest effect level given in Long et al. (1995). Elevated levels of avian teratogenesis have been associated with sediment selenium concentrations as low as 0.9 mg/kg in ponds near Tulare, California (Skorupa, 1998).

As the sludge may be aerated during dredging, there is some potential that reduced forms of selenium such as selenides may be oxidized and could become mobile in water when sediment is released back to the Salton Sea. Concentrations of all other

trace elements in bottom sediment analyzed by LFR did not appear to be problematic if the material were redeposited in the Sea.

The high organic material content of the sediment and its reduced state, as evidenced by hydrogen sulfide odor, indicate there will be considerable oxygen demand as the material is released back to the Sea. This will cause localized oxygen depletion with potentially adverse effects on biota. The amount of hydrogen sulfide released from the sediment is not known, but it may cause localized odor problems and can be toxic to biota.

- 750 acres of sea bottom + 140 acres shoreline disrupted- effects on habitat or wetlands

Polychaetes (*Neanthes succinea*) are the principle food item for many fish and shorebirds in the Salton Sea and densities average 1,000 individuals/m<sup>2</sup> throughout much of the Sea (Dexter et al., 1999). Walker et al. (1961) estimated the spring standing crop of *Neanthes* at 300 pounds/acre. Replacement of soft bottom material with dike fill could result in loss of about 3 billion polychaetes in the footprint of the dikes. It is not known if polychaetes would colonize the seaward side of the dike surface as it would be covered with riprap. An additional 34.3 square miles of Sea bottom converted to hypersaline pond would also be devoid of polychaetes. This would constitute a loss of about  $8.8 \times 10^{10}$  polychaetes

- Vehicle traffic and dust generation- effects on humans

This will be addressed by Tetra Tech as it is common to all constructed alternatives.

- Desert pupfish in San Felipe Creek and local drains

Pupfish use of 45.5% and 72% of Imperial Valley drains has been documented in two studies (Remington and Hess, 1993; Lau and Boehm, 1991, both cited in Hurlbert, 1997). Pupfish are believed to live in San Felipe Creek where high flows periodically move them to the Salton Sea and they return to the creek when flows subside (Glenn Black, personal commun.).

- Bird use of existing area, specifically near mouth of San Felipe Cr. and New River  
Studies underway by Point Reyes Bird Observatory (1999), indicate the majority of nesting colonies of birds are concentrated near the river deltas such as the New and Alamo Rivers at the southern end of the sea. The only systematic study of black rail distribution (a bird listed by California as Threatened) reported 13 of the 23 birds were observed at the mouth of the New River (Point Reyes Bird Observatory, 1999).

Tetra Tech is pursuing acquisition of additional information from the Redlands database and directly from Point Reyes Bird Observatory.

- Noise and activity effects on bird populations using the area

Construction of a haul road and activities associated with pond construction will generate noise and human presence. These activities may result in loss of habitat, habitat fragmentation, and isolation of some biota. Several studies indicate that species richness and density of individual species decrease with increasing proximity to roads and construction (Van der Zande, 1980; Reijnen and Foppen, 1994). Noise appears to be a

major factor in reduced populations of birds near roads. Van der Zande et al. (1980) reported that species density losses for some species of up to 56% were found within a 1.24 mile disturbance zone adjacent to major highways and up to 34% for lightly traveled dirt roads.

## **Pond Operation**

- Effectiveness of fish screen on inflow –

The issue of entrainment of small fish and fish larvae is being addressed by Tetra Tech as water intakes are part of several alternatives under consideration.

- Seismic stability during operation and after closure

This issue is being addressed by Tetra Tech as seismic concerns are part of several alternatives under consideration.

- Loss of waterfowl and fish habitat along shoreline and at river/stream deltas-

Fresh water and brackish water areas of the New and Alamo Rivers are classified as two of the most important fish habitats of the Salton Sea (Costa-Pierce, 1999) and the mouth of the New River is part of Unit 1 of the Sonny Bono Memorial National Wildlife Refuge. The Trifolium 2 drain enters the Sea immediately west of the mouth of the New River and 62 species of birds including several sensitive-status species have been documented using this water source (Hurlbert, 1997). Burrowing owls were quite numerous (83 counted) along the banks of this drain and the drain ranked in the top three for bird use of all drains surveyed (Hurlbert, 1997).

- Pond salinity will reach 60,000 ppm having adverse effects on feeding and nesting waterfowl

This may be addressed by posing several questions:

- (1) Are water birds attracted to hypersaline ponds if less saline water is available?

Investigations of bird use of evaporation ponds in the San Joaquin Valley have shown considerable attraction to the ponds for feeding, roosting, cover or escape from predators. Personal communications from the San Francisco Bay National Wildlife Refuge (Bradford et al., 1989) indicated the highest diversity of food items was found in ponds with salinities close to that of sea water. At salinities greater than 200,000 ppm, productivity decreased and brine shrimp levels declined. However, these extremely saline ponds were still used by birds for roosting, especially during high tide when feeding areas were inundated.

Use of even highly concentrated saline ponds in Hawaii was reported in a study of bird use at increasing salinities. Black necked stilts and phalaropes were found in ponds having salinities up to 240,000 ppm. Generally, the only sites that birds did not use were crystallization ponds (>300,000 ppm) and bittern ponds (concentrated magnesium and potassium salts after removal of sodium chloride) (Harvey, 1989). The western snowy plover also has been reported to be attracted to evaporation ponds in the San Joaquin Valley (Ivy, 1984). As the Salton Sea supports the largest population of wintering snowy plovers in western North America (Shuford et al., 1995 as cited in Point Reyes Bird



Observatory, 1999), it is reasonable to expect seasonally large numbers of these birds near the evaporation ponds.

(2) What dietary items may be present in evaporation ponds?

Feeding activities of water birds in saline ponds in prairie wetlands was found to vary with the succession of organisms as salinity increased. Breeding dabbling ducks were observed to concentrate and feed in saline lakes of high salt content. Blue-winged teal shifted from a diet dominated by amphipods, gastropods, and insects on lakes where specific conductance averaged 1,234  $\mu\text{S}$  (about 950 ppm) to a diet dominated by Anostraca crustaceans (primarily *Brachinecta* and *Artemia*) on lakes averaging 32,583  $\mu\text{S}$  (about 24,500 ppm) (Swanson et al., 1988).

Organisms potentially useable for food by water birds undergo succession of species as the salinity increases and some species will be more attractive than others as food items. At a Salton Sea salinity of about 46 g/L several species of zooplankton reported by Carpelan (in Walker, 1961-salinity of 34 g/L) still are present (Dexter et al., 1999) and would likely survive the initial evaporation stages in ponds. The rotifer, *Brachionus plicatilis* is reported to be tolerant of salinities up to 90,000 ppm (Hammer, 1986), and was dominant zooplankton in the Sea in summer of 1997. Winter populations of rotifers may be dominated by *Synchaeta* sp. The copepod, *Apocyclops dengizicus* is present primarily during summer and it and a harpacticoid copepod (*Cletocamptus deitersi*) may be predaceous on rotifers and artemia nauplii (Dexter, 1993). It is unlikely that the polychaete, *Neanthes succinea*, currently the major invertebrate utilized by birds in the Salton Sea, would be present in the evaporation ponds as reproduction is inhibited at salinities exceeding 50,000 ppm (Kuhl and Oglesby, 1979).

Brine shrimp (*Artemia franciscana*, the species that would occur in Salton Sea) are present in saline water from about 30,000 ppm to near saturation (Hammer, 1986) and would be the principle zooplankton present in the evaporation ponds at higher salinities. They have been reported as dominant food items in prairie ponds where the specific conductance was 35,000  $\mu\text{S}$  (salinity about 26,000 ppm) (Swanson et al., 1988). They are capable of adapting to water in which sodium sulfate is precipitating as long as sodium chloride is the dominant salt (Croghan, 1958). However, artemia are not tolerant of high concentrations of potassium salts (Hammer and Parker, 1984) and toxicity is dependent on molar ratios of sodium to potassium being less than about 12 (Bowen and Carl, 1992). Microcosm experiments using Salton Sea water done at San Diego State University (Dexter et al., 1999) showed that at a salinity of 65 g/L, artemia were present at high densities and were accompanied by brinefly larvae (*Ephydra riparia*) and some surface-dwelling insects (*Trichocorixa reticulata*). Some of the ephydra are very tolerant of high salinity. Ephydra occur and reproduce in the north arm of Great Salt Lake at a salinity of 330,000 g/L but numbers are fewer than in the less saline south arm of the lake (Post, 1977). Water birds are known to feed heavily on artemia and ephydra in Great Salt Lake (commun. D. Paul, Utah Div. of Wildlife Resources). Artemia also are a highly preferred food for eared grebes in saline environments (Jehl, 1988).

(3) What are the effects of salinity on water birds feeding in evaporation ponds?

Adult water birds contain a supraorbital salt gland that actively removes salt obtained through diet or drinking water. Ducklings do not have functional salt glands until they are 6 days old and must have ready access to fresh water in order to survive (Swanson et al., 1988). Mitcham and Wobeser (1988) found that mallard ducklings given water with a specific conductivity of 35,000  $\mu\text{S}$  died within 60 hours. Assuming salinity for water high in sodium sulfate is 75% of the value for conductivity (Hem, 1985) this would be equivalent to a salinity of 26,250 ppm. Clearly, ducklings raised on Salton Sea need access to fresh water during their development.

Salt encrustation of the feathers may prevent birds from flying from ponds when water is nearing saturation with respect to salt (primarily this is sodium chloride or sodium sulfate). Accompanying the encrustation is sodium toxicity as birds attempt to groom their feathers. Mortality due to salt encrustation on waterfowl has been reported at specific conductances of 77,000 to 90,000 (salinities of 58,000 to 68,000 ppm) in a small sodium-sulfate lake in Canada (Wobeser and Howard, 1987). Sodium toxicity to waterfowl has been associated with ingestion of water containing 17,000 mg/L of sodium in North Dakota (Windingstad et al., 1987) and in playa lakes containing total solids concentrations of 200,000-300,000 ppm in New Mexico (Meteyer et al., 1997).

- Variable salinity and ionic composition in the ponds depending on stage of evaporation.

The amount and types of salts in solution and precipitation depends on composition of salts in the water, average pond temperature, daytime and night time pond temperatures, evaporation rate, and depth of pond (Butts, 1980).

<u>% of Water Evap.</u>	<u>Salt Precipitate</u>	<u>Composition</u>
75	Halite	$\text{NaCl}$
70	Mirabilite	$\text{Na}^2\text{SO}^4 \cdot 10 \text{H}^2\text{O}$
>65	Epsomite	$\text{MgSO}^4 \cdot 7 \text{H}_2\text{O}$
80	Schoenite	$\text{K}^2\text{SO}^4 \cdot \text{MgSO}^4 \cdot 6 \text{H}^2\text{O}$
83	Hexahydrate	$\text{MgSO}^4 \cdot 6 \text{H}^2\text{O}$
87	Kainite	$\text{KCl} \cdot \text{MgSO}^4 \cdot 2.75 \text{H}^2\text{O}$
90	Carnallite	$\text{KCl} \cdot \text{MgCl}^2 \cdot 6 \text{H}^2\text{O}$

Invertebrates such as artemia that are intolerant of high potassium concentrations, particularly in low sodium waters tend to disappear as halite begins precipitating and the ratio of sodium to potassium declines.

- Use of dikes for bird nesting

Colonial nesting birds such as gulls, terns, and skimmers utilize a variety of island-like (surrounded by water) features: earthen levee remnants, low-lying boulder and barnacle bars, and clumps of vegetation. Similarly, cormorants and pelicans may nest on rocky ledges (Point Reyes Bird Observatory, 1999). The long dikes surrounding the evaporation ponds may appear to be islands and may draw nesting birds; and the exposed riprap may attract pelicans. As the dikes have land access and may have human usage, nesting areas on the dikes would place birds at risk if access were not controlled. Management of the dikes for bird production should be considered as an enhancement of

their construction. Increased access to fresh water may be needed to mitigate the high salinity in water and food from the ponds.

- Evaporative concentration of trace elements in the ponds.

There could be a 6 or 7 fold concentration of dissolved elements present in Salton Sea water as it evaporates in the ponds. While some elements may complex with the salts and not be readily available, others may remain in solution and be potentially toxic to organisms using the ponds, and some metals may precipitate. Stiller and Sigg (1990) found that when the salinity of the Dead Sea reached 340 g/L, precipitation of halite occurred which co-precipitated many heavy metals from the water. Most of the cadmium, much of the lead, and varying amounts of zinc and copper were removed primarily as salts of chloride.

- Desert pupfish access and mobility near San Felipe Cr.

Little is known about the movements of pupfish in San Felipe Creek. An unobstructed access to the creek from the sea likely would be required as the fish prefer shallow backwaters areas where predators are fewer. They are reportedly quite tolerant of salinities as high as twice that of sea water (70,000 ppm) (Walker et al., 1961, p.80) and could possibly live and spawn in early stage evaporation ponds. They are tolerant of temperature extremes that might occur in shallow evaporation ponds.

- Circulation changes in delta of New River

This will be addressed by Tetra Tech as part of the modeling effort using the Orlob model for circulation patterns in the Salton Sea.

- Location of pump intakes

This will be addressed by Tetra Tech as part of the design of the fish screens.

- Stability of dikes and ability to pump water if sea elevation drops below -227

This will be addressed by Tetra Tech.

- Abandonment of ponds with no cap: creation of hypersaline habitat during wet weather

Heavy rainfall on the uncovered surface of the salt block could solubilize the salt and create a hypersaline pond that may attract water birds and wildlife.

- Creation of deeper water habitat at toe of dikes and barnacle colonization of riprap.

The seaward extent of the dikes will create water with a depth of about 23 feet, covering much of the riprap facing. Barnacles (*Balanus amphitrite*) in the Sea form dense colonies on rock substrates reaching densities of 33,000 organisms per m<sup>2</sup> (D. Dexter, personal commun. June 1999). While the barnacles avoid mud and clay substrate, they appear to be substrate limited in the Sea and will quickly colonize most other substrates.



The appearance of large numbers of barnacles in deeper water may attract Sargo which feed on the barnacles (G. Black, personal commun. June 1999).

## References Cited

- Bowen, S.T., and Carl, J.R., 1992, *Artemia* populations: differences in tolerance of high potassium. p. 105-116, *In* Robarts, R.D., and Bothwell, M.L., eds., *Aquatic ecosystems in semi-arid regions: Implications for resource management*: National Hydrology Research Institute Symposium Series 7, Environment Canada, Saskatoon.
- Bradford, D.F., Drezner, D., Shoemaker, J.D., and Smith L., 1989, Evaluation of methods to minimize contamination hazards to wildlife using agricultural evaporation ponds in the San Joaquin Valley California: ESE Final Report No. 89-64, California Dept. of Water Resources, Sacramento, CA.
- Bureau of Reclamation, 1997, Salton Sea alternative evaluation, Final Draft Report: Lower Colorado Region, 223 p.
- Bureau of Reclamation, 1999, Salton Sea Alternatives-Final Preappraisal Report: Lower Colorado Region, 108 p.
- Butts, D.S., 1980, Chemistry of Great Salt Lake brines in solar ponds. p. 169-173 *In* Gwynn, J.W. ed., *Great Salt Lake-a scientific, historical and economic overview*: Utah Geological and Mineral Survey Bulletin 116.
- Costa-Pierce, Barry, 1999, Fish and fisheries of the Salton Sea: unpubl. Report to Salton Sea Science Subcommittee, 21 p.
- Croghan, P.C., 1958, The survival of *Artemia salina* (L.) in various media: *Journ. of Experimental Biology*, v. 35, p. 213-218.
- Dexter, D, 1993, Salinity tolerance of the copepod, *Apocyclops dengizicus* (Lepeschkin, 1900), a key food chain organism in the Salton Sea, California: *Hydrobiologia*, v. 267, p. 203-209.
- Dexter, D., Tiffany, M.A., Reifel, K., and Hurlbert, S.H., 1999, A synthesis of our knowledge of the biological limnology of the Salton Sea: unpubl. Report to Salton Sea Science Subcommittee, 19 p.
- Hammer, U.T., 1986, *Saline lake ecosystems of the world*: W. Junk, publisher, The Hague, 616 p.
- Hammer, U.T., and Parker, R.D., 1984, Limnology of a perturbed highly saline Canadian lake: *Arch. fur Hydrobiologie*, V. 102. P. 31-42.
- Harvey, T., 1989, US Fish and Wildlife Service, Hawaiian Islands. Telephone conservation with L. Smith of UCLA, April 7.

- Hem, J.D., 1985, Study and interpretation of the chemical characteristics of natural water: U.S. Geological Survey Water-Supply Paper 2254, 264 p.
- Hurlbert, Allen (editor), 1997, Wildlife use of agricultural drains in the Imperial Valley, California: Department of Interior, National Irrigation Water Quality Program, 98 p.
- Ivey, G.L., 1984, Some recent records for the snowy plover in the San Joaquin Valley, California: *Western Birds*, v. 15, p. 189.
- Jehl, J.R., Jr., 1988, Biology of the Eared Grebe and Wilson's Phalarope in the nonbreeding season: A study of adaptations to saline lakes: *Studies in Avian Biology*, v. 12, p. 1-74.
- Kuhl, D.I., and Oglesby, L.C., 1979, Reproduction and survival of the pileworm *Neanthes succinea* in higher Salton Sea salinities: *Biological Bulletin* v. 157, p. 153-165.
- Long, E.R., MacDonald, D.D., Smith, S.L., and Calder, F.D., 1995, Incidence of adverse biological effects within ranges of chemical concentrations in marine and estuarine sediments: *Environmental Management*, v. 18, p. 81-97.
- Meteyer, C.U., Dubielzig, R.R., Dein, F.J., Baeten, L.A., Moore, M.K., Jehl, J.R., Jr., and Wesenberg, K., 1997, Sodium toxicity and pathology associated with exposure of waterfowl to hypersaline playa lakes of southeast New Mexico: *Journ. of Veterinary Diagnostic Investigations*, v. 9, p. 269-280.
- Mitcham, S.A., and Wobeser, G., 1988, Toxic effects of natural saline waters on mallard ducklings: *Journ. of Wildlife Diseases*, v. 24, p. 45-50.
- Point Reyes Bird Observatory, 1999, A synthesis of avian concerns at the Salton Sea: unpubl. Report to the Salton Sea Science Subcommittee, 18 p.
- Post, F.J., 1977, The microbial ecology of the Great Salt Lake: *Microbial Ecology* v. 3, p. 143-165.
- Reijnen, R., Foppen, R., and Meeuwsen, H., 1996, The effects of traffic on the density of breeding birds in Dutch agricultural grasslands: *Biological conservation*, V. 75, p. 255-260.
- Skorupa, J.P., 1998, Selenium: p. 139-184 in *Guidelines for Interpretation of the biological effects of selected constituents in biota, water, and sediment*: Department of Interior, National Irrigation Water Quality Program.
- Stiller, M., and Sigg, L., 1990, Heavy metals in the Dead sea and their coprecipitation with halite. p. 23-33 In Comin, F. and Northcote, T., eds., *Saline Lakes IV: Hydrobiologia*, v. 197.

- Swanson, G.A., Winter, T.C., Adomaitis, V.A., LaBaugh, J.W., 1988, Chemical characteristics of prairie lakes in south-central North Dakota-Their potential for influencing use by fish and wildlife: USFWS, fish and Wildlife Tech. Report 18, 44 p.
- Swarth, C.W., C. Akagi, and P. Metropulos, 1982, The distribution patterns and ecology of waterbirds using the Coyote Hills Salt Ponds: U.S. Dept. of Interior, Fish and Wildlife Service, San Francisco Bay National Wildlife Refuge.
- Van der Zande, A.N., Keurs, W.J., and van der Weijden, J., 1980, The impact of roads on the densities of four bird species in an open field habitat-evidence of a long-distance effect: *Biological Conservation*, v. 18, p. 299-320.
- Walker, B.W., 1961, The ecology of the Salton Sea, California, in relation to the sport fishery: State of California, Dept. of fish and Game, Fish Bulletin No. 113, 204 p.
- Windingstad, R.M., Kartch, F.X., Stroud, R.K., and Smith M.R., 1987, Salt toxicosis in waterfowl in North Dakota: *Journ. of Wildlife Diseases*, v. 23, p. 443-446.
- Wobeser, G., and Howard, J., 1987, Mortality of waterfowl on a hypersaline wetland as a result of salt encrustation: *Journ. of Wildlife Diseases*, v. 23, p. 127-134.

**Participants-Salton Sea Evaporation Ponds Workshop  
SDSU, June 15, 1999**

Doyle Stephens, chair, USGS 801-975-3396  
 Chuck Keene, S4, Calif. Dept of Water Resources, 818-543-4600 ext 246  
 Glenn Black, S4, Calif. Dept. of Fish and Game, 909-597-5043  
 John Elder, S4, USGS, 608-821-3854  
 Tom Whitehead, Tetra Tech 415-974-1221  
 Bill Thompson, USBR, 303-445-3128  
 Carol Roberts, USFWS, 760-431-9440  
 Dan Chambers, U. of Redlands, 909-335-5268  
 Phil Pryde, Audubon California, 619-594-5525  
 Richard Vogl, Levine-Fricke Recon, 949-442-7276