

# STREAM INVENTORY REPORT

## North Fork DeHaven Creek

### WATERSHED OVERVIEW

North Fork DeHaven Creek is a tributary to DeHaven Creek (Figure 1). Elevations range from 100 feet at the mouth of the creek to 1,400 feet in the headwater areas. North Fork DeHaven Creek's legal description at the confluence with DeHaven Creek is T21N R17W S22. Its location is 39°39'36"N. latitude and 123°44'32"W. longitude according to the USGS Lincoln Ridge 7.5 minute quadrangle.

### HABITAT INVENTORY RESULTS

The habitat inventory of October 9, 1996 was conducted by Dave Wright. The total length of surveyed stream in North Fork DeHaven Creek was 2,318 feet (0.44 miles, 0.70 KM) (Table 1). There were no side channels in this creek.

Flow measured at the mouth of North Fork DeHaven Creek on October 9, 1996 was 0.32 cubic feet per second (cfs).

North Fork DeHaven Creek consists of one reach: A B3 for the entire length of stream surveyed.

Table 1 summarizes the Level II riffle, flatwater and pool habitat types. By percent occurrence, riffles comprised 28%, flatwater 36% and pools 34% of the habitat types (Graph 1). By percent total length, riffles comprised 36%, flatwater 46% and pools 8% (Graph 2).

Nine Level IV habitat types were identified and are summarized in Table 2. The most frequently occurring habitat types were step runs 26%, low gradient riffles 24% and plunge pools 20% (Graph 3). The most prevalent habitat types by percent total length were step runs at 41%, low gradient riffles at 34% and dry units at 11% (Table 2).

Table 3 summarizes main, scour and backwater pools which are Level III pool habitat types. Scour pools were most often encountered at 59% occurrence and comprised 62% of the total length of pools.

Table 4 is a summary of maximum pool depths by Level IV pool habitat types. In second order streams, pools with depths of two feet (0.61 m) or greater are considered optimal for fish habitat. In North Fork DeHaven Creek, two of the 17 pools (12%) had a depth of two feet or greater (Graph 4).

The depth of cobble embeddedness was estimated at pool tail-outs. Of the pool tail-outs measured, 0% had a value of 1, 0% had a value of 2, 0% had a value of 3 and 100% had a value of 4 (Graph 5).

## **North Fork DeHaven Creek**

Of the Level II habitat types, pools had the highest mean shelter rating at 6 (Table 1). Of the Level III pool habitat types, main channel pools had the highest mean shelter rating at 8 (Table 3).

Of the 17 pools, none were formed by large woody debris (LWD) (Table 4).

Table 6 summarizes dominant substrate by Level IV habitat types. Of the low gradient riffles fully measured, 50% had silt/clay and another 50% had sand as the dominant substrate (Graph 6).

Mean percent closed canopy was 81%: 48% coniferous trees and 33% deciduous trees. Mean percent open canopy was 19% (Graph 7, calculated from Table 7).

Table 7 summarizes the mean percent substrate/vegetation types found along the banks of the stream. Mean percent right bank vegetated was 72% while mean percent left bank vegetated was 81%. Grass was the dominant bank vegetation type in 50% of the units fully measured. The dominant substrate composing the structure of the stream banks was cobble/gravel, found in 59% of the units fully measured.

## **DISCUSSION**

The information gathered in the process of habitat typing will provide Georgia-Pacific with baseline data on the current condition of this creek and the available habitat for salmonids.

When reviewing North Fork DeHaven Creek data it is important to consider the short distance surveyed. The survey was limited to approximately 2,300 feet with only 50 units; therefore, many of the determinations for the indicated parameters were based on only one or two completely measured units. Determinations based on such a limited sample size may lack statistical validity and therefore are of questionable analytical value.

### **Level II habitat types by percent occurrence and length**

Flatwater habitat types comprised a relatively high percentage of the units by both percent occurrence and length at 36% and 46% respectively (Table 1 and Graph 1). These unit types usually do not provide optimal spawning or rearing habitat for salmonids. Riffle habitat units comprised a moderate percentage by both percent occurrence and length at 28% and 36% respectively. Pools comprised a high percentage by percent occurrence and a low percentage by length at 34% and 8% respectively. Riffles usually provide good spawning habitat while pools provide important rearing habitat. In addition, Mundie (1969) reported that invertebrate food production is maximized in riffles while pools provide an optimum feeding environment for coho. In fact, the most productive streams are those consisting of a pool to riffle ratio of approximately one to one (Ruggles 1966).

## **North Fork DeHaven Creek**

### **Pool Depth**

According to Flosi and Reynolds (1994), a stream with at least 50% of its total habitat comprised of primary pools is generally desirable. Primary pools are at least two feet deep in first and second order streams and at least three feet deep in third order streams. The information from Graph 4 on maximum depth in pools was used to determine percent of primary pools. North Fork DeHaven Creek, a second order stream, is comprised mainly of shallow pools with 12% of the pools having a maximum depth of two feet or greater.

### **Instream Shelter**

Instream shelter ratings are derived from two measurements: instream shelter complexity and instream shelter percent cover. The first is a value rating which provides a relative measure of the quality and composition of the shelter, and the second is a measure of the area of a habitat unit covered by shelter. The various types of instream shelter include LWD, small woody debris, boulders, root wads, terrestrial vegetation, aquatic vegetation, bedrock ledges and undercut banks. Of the Level II habitat types, pools had the highest shelter rating at 6. Of the Level III habitat types, main channel pools had the highest shelter rating at 8. These values are very low as shelter values of 80 or higher are considered optimal for good rearing habitat (Flosi and Reynolds 1994).

### **Large Woody Debris**

The presence of large woody debris in streams is a significant component of fish habitat. Woody debris creates areas of low flow, providing a refuge for fish during periods of high flow (Robison and Beschta, 1990). Woody debris also provides cover for fish, lowering the risk of predation. The percent of pools formed by LWD in North Fork DeHaven Creek was 0%, indicating that there is a lack of sufficient LWD in this creek. Based on data from Georgia-Pacific's 1995 Aquatic Vertebrate Study, the only coho found in the Ten Mile River Basin were in stream reaches where approximately 50% of pools were formed by large woody debris. Those reaches that did not support coho had a significantly lower percentage of pools formed by large woody debris (Ambrose et al, 1996). This suggests that a low percentage of LWD formed pools could adversely affect juvenile coho populations (C.S. Shirvel 1990).

The above LWD analysis pertains only to pools formed by logs or root wads as described in Flosi and Reynolds (1994): lateral scour pool-log enhanced, lateral scour pool-rootwad enhanced, backwater pool-log formed and backwater pool-rootwad formed. Other pools containing LWD as a component were not included in the calculation. For example, plunge pools may be formed by boulders, bedrock or LWD, but are not described as such by habitat unit types. Therefore, the LWD formed pool calculation is limited to four pool types and does not quantify the total amount of LWD in North Fork DeHaven Creek.

### **Canopy**

There are two important benefits of canopy cover in coastal streams. Canopy keeps stream temperatures cool as well as providing nutrients in the form of leaf litter and organic material

## **North Fork DeHaven Creek**

(Bilby 1988). This leaf litter, organic material, and their associated nutrients are utilized as a food source by benthic macroinvertebrates (aquatic insects). The macroinvertebrates, in turn, are major food sources for most fish species in forested areas (Gregory et al., 1987). Mean percent canopy cover for the North Fork DeHaven Creek was 81%. This is relatively high since a canopy cover of 80% or higher is considered optimum (Flosi and Reynolds, 1994).

Coniferous trees occupied a larger portion of the canopy than did deciduous trees. Coniferous trees comprised 48% and deciduous trees 33% of the canopy. The significance of this is that wood from coniferous species does not deteriorate as rapidly as wood from alder and most other deciduous species (Sedell, *et al.* 1988). Therefore, more LWD would be available in the future for fish cover and LWD formed pools in this creek and others dominated by coniferous species.

### **Embeddedness**

High embeddedness values (silt levels), such as those found in North Fork DeHaven Creek, have been associated with many negative impacts to salmonids. These negative impacts can be observed in important environmental components of salmonid habitat, such as pool habitats, dissolved oxygen levels and water temperatures.

The impact high silt levels have on pool habitat is that they fill in and eventually eliminate pools. As already mentioned, pools provide important habitat for rearing salmonids.

High silt levels also impact oxygen levels in the water. They do so by reducing water circulation within the substrate, thus lowering the oxygen levels needed by salmonid eggs (Sandercock, 1991). This can hinder the survival of the eggs deposited in redds, as well as the survival of juvenile salmonids.

Water temperature is impacted by high silt levels in several ways. Hagans et al (1986) reported the following impacts to water temperatures: 1) the loss of a reflective bottom; 2) darker sediment (as opposed to clean gravels) storing heat from direct solar radiation which is then transferred to the water column; and 3) a reduction in the flow of water through the substrate interstitial spaces thereby exposing more of the water column to direct solar radiation.

Another means by which water temperatures are increased is through the widening of stream channels: over time, high silt levels increase the substrate surface level of the creek, resulting in a wider, shallower stream channel (Flosi and Reynolds 1994). In shallow streams more surface area is exposed to the sun relative to the volume of water, leading to an increase in solar heating which in turn leads to higher water temperatures.

Substrates embedded with silt in varying degrees were given corresponding values as follows: 0-25%= value 1, 26 - 50% = value 2, 51 - 75% = value 3 and 76 - 100% = value 4. According to Flosi and Reynolds (1994), creeks with embeddedness values of two or higher are considered to have poor quality fish habitat. In North Fork DeHaven Creek, 100% of the pool tail-outs measured had embeddedness values of two or more.

## North Fork DeHaven Creek

It is important to consider, however, that the above embeddedness values were obtained in the summer during low flow conditions. In winter and spring, flows are usually higher due to the rainy season and the lowered evapotranspiration of the trees. This higher flow can carry away some of the previously deposited silt to sites further downstream. Therefore, embeddedness values may fluctuate throughout the year along different sections of the stream.

### Substrate

In North Fork DeHaven Creek, none of the low gradient riffles had gravel or small cobble as the dominant substrate. The absence of gravel and small cobble in riffles indicates that there is an insufficient amount of substrate available as potential spawning habitat in this creek. It is also important to note that 50% of the low gradient riffles surveyed were dominated by silt and clay and another 50% by sand. Another point to consider is that regardless of the amount of substrate or spawning habitat available, this habitat may not be suitable for salmonids if it is highly embedded.

Overall, North Fork DeHaven Creek appears to have a low percentage of primary pools and a lack of LWD formed pools. This stream also appears to have low shelter values, high embeddedness values and insufficient substrate for spawning. This creek does appear to have sufficient canopy.

### RECOMMENDATIONS

- 1) Due to marginal habitat and small size of this creek, the net results of any expense or effort directed towards creek restoration, other than maintaining good canopy cover, would not be worth the expended effort.

### COMMENTS AND LANDMARKS

The following landmarks and possible problem sites were noted. All distances are approximate and taken from the beginning of the survey reach.

Position  
(ft):

Comments:

---

1456	Six percent slope over 100 feet. Possible barrier.
1487	Seven foot high plunge into two foot pool, possible barrier.
1532	No fish observed for last 10 units.
1560	Solid mud bottom. Mud is eight inches thick. Seven foot high plunge into 3' deep pool. Possible barrier.

**North Fork DeHaven Creek**

2318            End of survey due to diminished habitat. No fish observed for 20-30 units.

North Fork DeHaven Creek

