

STREAM INVENTORY REPORT

Little Bear Creek

INTRODUCTION

A stream inventory was conducted during the summer of 1995 on Little Bear Creek. The inventory was conducted in two parts: habitat inventory and biological inventory. The objective of the habitat inventory was to document the habitat available to anadromous salmonids in Little Bear Creek. The objective of the biological inventory was to document the presence and distribution of juvenile salmonid species. There is no known record of adult spawning surveys having been conducted on Little Bear Creek.

The objective of this report is to document the current habitat conditions, and recommend options for the potential enhancement of habitat for chinook salmon, coho salmon and steelhead trout. Recommendations for habitat improvement activities are based upon target habitat values suitable for salmonids in California's north coast streams.

WATERSHED OVERVIEW

Little Bear Creek is tributary to Usal Creek, a tributary to the Pacific Ocean, located in Mendocino County, California (Figure 1). Little Bear Creek's legal description at the confluence with Usal Creek is T23N R18W. Its location is 39°51'45" north latitude and 123°50'29" west longitude. Little Bear Creek is a first order stream and has approximately 1.5 miles of blue line stream according to the USGS Hales Grove and Piercy 7.5 minute quadrangles. Little Bear Creek drains a watershed of approximately 0.7 square miles. Summer base runoff is approximately 0.2 cubic feet per second (cfs) at the mouth. Elevations range from about 100 feet at the mouth of the creek to 1800 feet in the headwater areas. Redwood and Douglas fir forest dominates the watershed. The watershed is privately owned and is managed for timber production. Foot access is available from a private road by crossing Usal Creek to the mouth of Little Bear Creek.

METHODS

The habitat inventory conducted in Little Bear Creek follows the methodology presented in the *California Salmonid Stream Habitat Restoration Manual* (Flosi and Reynolds, 1991 rev. 1994). The California Conservation Corps (CCC) Technical Advisors and Watershed Stewards Project/AmeriCorps (WSP/AmeriCorps) members that conducted the inventory were trained in standardized habitat inventory methods by the California Department of Fish and Game (DFG). Little Bear Creek personnel were trained in May, 1995, by Gary Flosi. This inventory was conducted by a two-person team.

Little Bear Creek

SAMPLING STRATEGY

The inventory uses a method that samples approximately 10% of the habitat units within the survey reach (Hopelain, 1994). All habitat units included in the survey are classified according to habitat type and their lengths are measured. All pool units are measured for maximum depth. Habitat unit types encountered for the first time are further measured for all the parameters and characteristics on the field form. Additionally, from the ten habitat units on each field form page, one is randomly selected for complete measurement.

HABITAT INVENTORY COMPONENTS

A standardized habitat inventory form has been developed for use in California stream surveys and can be found in the *California Salmonid Stream Habitat Restoration Manual*. This form was used in Little Bear Creek to record measurements and observations. There are nine components to the inventory form.

1. Flow:

Flow is measured in cubic feet per second (cfs) at the bottom of the stream survey reach using standard flow measuring equipment, if available. In some cases flows are estimated.

2. Channel Type:

Channel typing is conducted according to the classification system developed and revised by David Rosgen (1985 rev. 1994). This methodology is described in the *California Salmonid Stream Habitat Restoration Manual*. Channel typing is conducted simultaneously with habitat typing and follows a standard form to record measurements and observations. There are five measured parameters used to determine channel type: 1) water slope gradient, 2) entrenchment, 3) width/depth ratio, 4) substrate composition, and 5) sinuosity.

3. Temperatures:

Both water and air temperatures are measured and recorded at every tenth habitat unit. The time of the measurement is also recorded. Both temperatures are taken in degrees Fahrenheit at the middle of the habitat unit and within one foot of the water surface.

4. Habitat Type:

Habitat typing uses the 24 habitat classification types defined by McCain and others (1988). Habitat units are numbered sequentially and assigned a type identification number selected from a standard list of 24 habitat types. Dewatered units are labeled "dry". Little Bear Creek habitat typing used standard basin level measurement criteria. These parameters require that the minimum length of a described habitat unit must be equal to or greater than the stream's mean wetted width. Channel dimensions were measured using hip chains, range finders, tape measures, and stadia rods. All units were measured for mean length; additionally, the first occurrence of each unit type and a randomly selected 10% subset of all units were sampled for

Little Bear Creek

all features on the sampling form (*Sampling Levels for Fish Habitat Inventory*, Hopelain, 1995). Pool tail crest depth at each pool unit was measured in the thalweg. All measurements were in feet to the nearest tenth.

5. Embeddedness:

The depth of embeddedness of the cobbles in pool tail-out reaches is measured by the percent of the cobble that is surrounded or buried by fine sediment. In Little Bear Creek, embeddedness was ocularly estimated. The values were recorded using the following ranges: 0 - 25% (value 1), 26 - 50% (value 2), 51 - 75% (value 3), 76 - 100% (value 4). Additionally, a rating of "not suitable" (NS) was assigned to tail-outs deemed unsuited for spawning due to inappropriate substrate particle size, having a bedrock tail-out, or other considerations.

6. Shelter Rating:

Instream shelter is composed of those elements within a stream channel that provide salmonids protection from predation, reduce water velocities so fish can rest and conserve energy, and allow separation of territorial units to reduce density related competition. The shelter rating is calculated for each fully-described habitat unit by multiplying shelter value and percent cover. Using an overhead view, a quantitative estimate of the percentage of the habitat unit covered is made. All cover is then classified according to a list of nine cover types. In Little Bear Creek, a standard qualitative shelter value of 0 (none), 1 (low), 2 (medium), or 3 (high) was assigned according to the complexity of the cover. Thus, shelter ratings can range from 0-300 and are expressed as mean values by habitat types within a stream.

7. Substrate Composition:

Substrate composition ranges from silt/clay sized particles to boulders and bedrock elements. In all fully-described habitat units, dominant and sub-dominant substrate elements were ocularly estimated using a list of seven size classes and recorded as a one and two respectively.

8. Canopy:

Stream canopy density was estimated using modified handheld spherical densimeters as described in the *California Salmonid Stream Habitat Restoration Manual*, 1994. Canopy density relates to the amount of stream shaded from the sun. In Little Bear Creek, an estimate of the percentage of the habitat unit covered by canopy was made from the center of approximately every third unit in addition to every fully-described unit, giving an approximate 30% sub-sample. In addition, the area of canopy was estimated ocularly into percentages of coniferous or deciduous trees.

9. Bank Composition and Vegetation:

Bank composition elements range from bedrock to bare soil. However, the stream banks are usually covered with grass, brush, or trees. These factors influence the ability of stream banks to withstand winter flows. In Little Bear Creek, the dominant composition type (options 1-4) and the dominant vegetation type (options 5-9) of both the right and left banks for each fully-

Little Bear Creek

described unit were selected from the habitat inventory form. Additionally, the percent of each bank covered by vegetation was estimated and recorded.

BIOLOGICAL INVENTORY

Biological sampling during stream inventory is used to determine fish species and their distribution in the stream. In Little Bear Creek fish presence was observed from the stream banks, and two sites were electrofished using one Smith-Root Model 12 electrofisher. These sampling techniques are discussed in the *California Salmonid Stream Habitat Restoration Manual*.

DATA ANALYSIS

Data from the habitat inventory form are entered into Habitat, a dBASE 4.2 data entry program developed by Tim Curtis, Inland Fisheries Division, California Department of Fish and Game. This program processes and summarizes the data, and produces the following six tables:

- Riffle, flatwater, and pool habitat types
- Habitat types and measured parameters
- Pool types
- Maximum pool depths by habitat types
- Dominant substrates by habitat types
- Mean percent shelter by habitat types

Graphics are produced from the tables using Lotus 1,2,3. Graphics developed for Little Bear Creek include:

- Riffle, flatwater, pool habitats by percent occurrence
- Riffle, flatwater, pool habitats by total length
- Total habitat types by percent occurrence
- Pool types by percent occurrence
- Total pools by maximum depths
- Embeddedness
- Pool cover by cover type
- Dominant substrate in low gradient riffles
- Percent canopy
- Bank composition by composition type
- Bank vegetation by vegetation type

HABITAT INVENTORY RESULTS

The habitat inventory of July 20, 1995, was conducted by Kyle Young (WSP/AmeriCorps) and Craig Mesman (CCC). The total length of the stream surveyed was 922 feet with an additional 24 feet of side channel.

Little Bear Creek

Flow was measured at the bottom of the survey reach with a Marsh-McBirney Model 2000 flowmeter at 0.18 cfs on August 1, 1995.

Little Bear Creek is a G4 channel type for the entire 922 feet of stream reach surveyed. G4 channels are entrenched, gully-like, step/pool channels with low width/depth ratios, moderate gradients, and gravel-dominant substrates.

Water temperature was 56 degrees Fahrenheit. Air temperatures ranged from 63 to 67 degrees Fahrenheit.

Table 1 summarizes the Level II riffle, flatwater, and pool habitat types. Based on frequency of **occurrence** there were 39% pool units, 35% riffle units, and 26% flatwater units (Graph 1). Based on total **length** of Level II habitat types there were 36% pool units, 34% riffle units, and 30% flatwater units (Graph 2).

Thirteen Level IV habitat types were identified (Table 2). The most frequent habitat types by percent **occurrence** were low-gradient riffles, 32%; runs, 16%; and mid-channel pools, 11% (Graph 3). Based on percent total **length**, low-gradient riffles made up 32%, step runs 16%, and runs 14%.

A total of 24 pools were identified (Table 3). Main channel pools were most frequently encountered at 46% and comprised 46% of the total length of all pools (Graph 4).

Table 4 is a summary of maximum pool depths by pool habitat types. Depth is an indicator of pool quality. Three of the 24 pools (12%) had a depth of two feet or greater (Graph 5).

The depth of cobble embeddedness was estimated at pool tail-outs. Of the 20 pool tail-outs measured, 2 had a value of 1 (10%); 9 had a value of 2 (45%); 9 had a value of 3 (45%); and none had a value of 4 (0%) (Graph 6). On this scale, a value of 1 indicates the highest quality of spawning substrate.

A shelter rating was calculated for each habitat unit and expressed as a mean value for each habitat type within the survey using a scale of 0-300. Riffle habitat types had a mean shelter rating of 23, and pool habitats had a mean shelter rating of 22 (Table 1). Of the pool types, the scour pools had the highest mean shelter rating at 29. Main channel pools had a mean shelter rating of 23 (Table 3).

Table 5 summarizes mean percent cover by habitat type. Small woody debris is the dominant cover type in Little Bear Creek. Graph 7 describes the pool cover in Little Bear Creek.

Table 6 summarizes the dominant substrate by habitat type. Gravel was the dominant substrate observed in the one low-gradient riffle measured (100%) (Graph 8).

The mean percent canopy density for the stream reach surveyed was 99%. The mean percentages of deciduous and coniferous trees were 63% and 37%, respectively. Graph 9 describes the canopy in Little Bear Creek.

Little Bear Creek

For the stream reach surveyed, the mean percent right bank vegetated was 71%. The mean percent left bank vegetated was 70%. The dominant elements composing the structure of the stream banks consisted of 10% bedrock, 0% boulder, 83% cobble/gravel, and 7% sand/silt/clay (Graph 10). Deciduous trees were the dominant vegetation type observed in 67% of the units surveyed. Additionally, 20% of the units surveyed had coniferous trees as the dominant vegetation, including down trees, logs, and root wads (Graph 11).

BIOLOGICAL INVENTORY RESULTS

Two sites were electrofished on August 1, 1995, in Little Bear Creek. The sites were sampled by Craig Mesman (CCC) and Kyle Young (WSP/AmeriCorps).

The first site sampled included habitat units 15-27, a series of pools, runs, and riffles approximately 243 feet from the confluence with Usal Creek. This site had a length of 145 feet. The site yielded thirty-one 0+ steelhead, one 1+ steelhead, and three Pacific giant salamanders.

The second site included habitat units 27-36, a series of pools, runs, and riffles located approximately 388 feet above the creek mouth. This site had a length of 170 feet. Numerous Pacific giant salamanders were observed, but no fish were sampled.

DISCUSSION

Little Bear Creek is a G4 channel type for the entire 922 feet of stream surveyed. The suitability of G4 channel types for fish habitat improvement structures is as follows: good for bank-placed boulders; fair for low-stage weirs, opposing wing deflectors, and log cover; and poor for medium-stage weirs, boulder clusters, and single wing deflectors.

The water temperature recorded on the survey day July 20, 1995, was 56 degrees Fahrenheit. Air temperatures ranged from 63 to 67 degrees Fahrenheit. This is a good water temperature for salmonids. To make any further conclusions, temperatures would need to be monitored throughout the warm summer months, and more extensive biological sampling would need to be conducted.

Flatwater habitat types comprised 30% of the total **length** of this survey, riffles 34%, and pools 36%. The pools are relatively shallow, with only 3 of the 24 (12%) pools having a maximum depth greater than 2 feet. In general, pool enhancement projects are considered when primary pools comprise less than 40% of the length of total stream habitat. In first and second order streams, a primary pool is defined to have a maximum depth of at least two feet, occupy at least half the width of the low flow channel, and be as long as the low flow channel width. Installing structures that will increase or deepen pool habitat is recommended for locations where their installation will not conflict with the modification of the numerous log debris accumulations (LDA's) in the stream.

Little Bear Creek

Nine of the 20 pool tail-outs measured had embeddedness ratings of 3. Only 2 had a 1 rating. Cobble embeddedness measured to be 25% or less, a rating of 1, is considered to indicate good quality spawning substrate for salmon and steelhead. In Little Bear Creek, sediment sources should be mapped and rated according to their potential sediment yields, and control measures should be taken.

The mean shelter rating for pools was low with a rating of 22. The shelter rating in the flatwater habitats was lower at 12. A pool shelter rating of approximately 100 is desirable. The relatively small amount of cover that now exists is being provided primarily by small woody debris in all habitat types. Additionally, large woody debris contributes a small amount. Log and root wad cover structures in the pool and flatwater habitats are needed to improve both summer and winter salmonid habitat. Log cover structure provides rearing fry with protection from predation, rest from water velocity, and also divides territorial units to reduce density related competition.

The one low-gradient riffle measured had gravel as the dominant substrate. This is generally considered good for spawning salmonids.

The mean percent canopy density for the stream was 99%. This is a relatively high percentage of canopy. In general, revegetation projects are considered when canopy density is less than 80%.

The percentage of right and left bank covered with vegetation was moderate at 71% and 70%, respectively. In areas of stream bank erosion or where bank vegetation is not at acceptable levels, planting endemic species of coniferous and deciduous trees, in conjunction with bank stabilization, is recommended.

RECOMMENDATIONS

- 1) Little Bear Creek should be managed as an anadromous, natural production stream.
- 2) The limited water temperature data available suggest that maximum temperatures are within the acceptable range for juvenile salmonids. To establish more complete and meaningful temperature regime information, 24-hour monitoring during the July and August temperature extreme period should be performed for 3 to 5 years.
- 3) Where feasible, design and engineer pool enhancement structures to increase the number of pools. This must be done where the banks are stable or in conjunction with stream bank armor to prevent erosion.
- 4) Increase woody cover in the pools and flatwater habitat units. Most of the existing cover is from small woody debris. Adding high quality complexity with woody cover is desirable and in some areas the material is locally available.
- 5) Inventory and map sources of stream bank erosion and prioritize them according to present and potential sediment yield. Identified sites, like the site at 697', should then be treated to reduce the amount of fine sediments entering the stream.

Little Bear Creek

- 6) Active and potential sediment sources related to the road system need to be identified, mapped, and treated according to their potential for sediment yield to the stream and its tributaries.
- 7) There are several log debris accumulations present on Little Bear Creek that are retaining large quantities of fine sediment. The modification of these debris accumulations is desirable, but must be done carefully, over time, to avoid excessive sediment loading in downstream reaches.

PROBLEM SITES AND LANDMARKS

The following landmarks and possible problem sites were noted. All distances are approximate and taken from the beginning of the survey reach.

Position (ft):	Comments:
0'	Begin survey at confluence with Usal Creek. Channel type is G4.
388'	Six foot jump with gravel retained 6' deep. Jump is obstructed by debris. Probable barrier. No fish sampled or observed upstream of this point.
524'	LDA 10' high x 20' wide x 30' long.
568'	LDA 7' high x 30' wide x 15' long retaining gravel 7' deep at base. Possible barrier.
697'	LDA 8' high x 20' wide x 30' long retaining gravel 8' deep at base. Possible barrier. Right bank road failure 40' high x 60' long.
723'	Five foot jump with no jump or landing pools.
922'	Six foot jump over logs. End of survey due to diminished habitat and flows.

REFERENCES

- Flosi, G., and F. Reynolds. 1994. California salmonid stream habitat restoration manual, 2nd edition. California Department of Fish and Game, Sacramento, California.
- Hopelain, J. 1995. Sampling levels for fish habitat inventory, unpublished manuscript. California Department of Fish and Game, Inland Fisheries Division, Sacramento, California.

Little Bear Creek

LEVEL III and LEVEL IV HABITAT TYPE KEY

HABITAT TYPE	LETTER	NUMBER
RIFFLE		
Low Gradient Riffle	[LGR]	1.1
High Gradient Riffle	[HGR]	1.2
CASCADE		
Cascade	[CAS]	2.1
Bedrock Sheet	[BRS]	2.2
FLATWATER		
Pocket Water	[POW]	3.1
Glide	[GLD]	3.2
Run	[RUN]	3.3
Step Run	[SRN]	3.4
Edgewater	[EDW]	3.5
MAIN CHANNEL POOLS		
Trench Pool	[TRP]	4.1
Mid-Channel Pool	[MCP]	4.2
Channel Confluence Pool	[CCP]	4.3
Step Pool	[STP]	4.4
SCOUR POOLS		
Corner Pool	[CRP]	5.1
Lateral Scour Pool - Log Enhanced	[LSL]	5.2
Lateral Scour Pool - Root Wad Enhanced	[LSR]	5.3
Lateral Scour Pool - Bedrock Formed	[LSBk]	5.4
Lateral Scour Pool - Boulder Formed	[LSBo]	5.5
Plunge Pool	[PLP]	5.6
BACKWATER POOLS		
Secondary Channel Pool	[SCP]	6.1
Backwater Pool - Boulder Formed	[BPB]	6.2
Backwater Pool - Root Wad Formed	[BPR]	6.3
Backwater Pool - Log Formed	[BPL]	6.4
Dammed Pool	[DPL]	6.5

Little Bear Creek

