

Mallard feeding on kokanee salmon eggs, Taylor Creek, California

LORI D. DIETER, ANDREW M. ATKINSON, AND DANIEL R. YPARRAGUIRRE*

California Department of Fish and Wildlife, Gray Lodge Wildlife Area, 3207 Rutherford Road, Gridley, CA 95948, USA (LDD, AMA)

California Department of Fish and Wildlife, Wildlife and Fisheries Division, 1416 Ninth Street, Sacramento, CA 95814 USA (DRY)

*Correspondent: dan.yparraguirre@wildlife.ca.gov

Key words: *Anas platyrhynchos*, California, food habits, kokanee, mallard, *Oncorhynchus nerka*, salmon eggs

Mallards (*Anas platyrhynchos*) are an abundant breeding duck throughout California (Kozlik 1974) and feed widely on both animal and plant materials (Martin et al. 1961, Swanson et al. 1985, Bellrose 1980, Cederholm et al. 1989). Munro (1923) appears to be the first published record of mallards consuming salmon eggs. In Alaska, Gleason (2007) reported mallards eating salmon carcasses, as well as a personal communication that the use of salmon eggs is a common foraging strategy of mallards. Gleason (2007) also included additional second-hand reports that anecdotally suggest that salmon form an important food source for mallards. Other reports (Munroe 1943, Willson and Halupka 1995), combined with observations from Canada (J. S. Gleason, U.S. Fish and Wildlife Service, personal communication June 2014) and California (E. G. Hunt, CDFW retired, personal communication June 2014) suggest that this foraging strategy is more common than is indicated in the literature. In this note we call attention to the personal observations or personal communications included in previously published papers (Munroe 1923, Gleason 2007). Further, we describe an additional observation of a male mallard consuming eggs of kokanee salmon (*Oncorhynchus nerka*) in a tributary to Lake Tahoe, California. Kokanee were introduced to Lake Tahoe in 1944 when fry were accidentally released (Cordone et al. 1971). Kokanee were planted annually thereafter in most of the major tributaries to the lake, and Taylor Creek, El Dorado Co., California became the major spawning area (Cordone et al. 1971).

At 1208 on 6 October 2013 authors Dieter and Atkinson observed a male mallard at Taylor Creek (38° 56' 6" N, 120° 3' 25" W) fly into the creek and actively begin stirring up kokanee redds with his feet and consuming the eggs as they floated to the surface (Figure 1). This behavior continued for approximately 20 minutes while dozens of fish were spawning, and within 3 m of many other people observing the annual kokanee run.



FIGURE 1.—Male mallard (*Anas platyrhynchos*) stirring kokanee (*Oncorhynchus nerka*) redds with his feet. The mallard repeatedly stirred the bottom and fed on the kokanee eggs as they floated up from the substrate. Taylor Creek, El Dorado County, California, 6 October 2013. Photograph by L. D. Dieter.

The kokanee seemed habituated to this activity, staying within the area disturbed by the mallard; one male kokanee was observed proximate to (i.e., <20 cm) the duck's feet without apparent alarm. On each dabble, the mallard ingested several kokanee eggs. The stirring and dabbling behavior continued as the duck moved upstream through the spawning kokanee. The duck, although habituated to people, appeared to be a normal wild mallard (Figure 2).



FIGURE 2.—The mallard observed stirring kokanee redds with his feet appeared to be a normal wild bird, albeit tolerant of human presence. Note the slim body conformation and wild plumage, both of which are atypical of domesticated birds. Taylor Creek, El Dorado County, California, 6 October 2013. Photograph by L. D. Dieter.

Mallards are well known to exploit seasonally abundant food resources (Heitmeyer 2006, Lafferty et al. 2013) in order to meet nutrient reserves necessary for reproduction and migration (Krapu 1981, Ankney et al. 1991). The ability to use various habitats and food sources likely explains the widespread abundance of mallards; information included herein further confirms this plasticity in their food habits, and provides an additional description of a foraging behavior that previously has been documented poorly in the formal literature.

LITERATURE CITED

- ANKNEY, C. D., A. D. AFTON, AND R. T. ALISAUSKAS. 1991. The role of nutrient reserves in limiting waterfowl reproduction. *Condor* 93:1029-1032.
- BELLROSE, F. C. 1980. Ducks, geese and swans of North America. Third edition. Stackpole Books, Harrisburg, Pennsylvania, USA.
- CEDERHOLM, C. J., D. B. HOUSTON, D. L. COLE, AND W. J. SCARLETT. 1989. Fate of coho salmon (*Oncorhynchus kisutch*) carcasses in spawning streams. *Canadian Journal of Fisheries and Aquatic Sciences* 46:1347-1355.
- CORDONE, A. J., S. J. NICOLA, P. H. BAKER, AND T. C. FRANTZ. 1971. The kokanee salmon in Lake Tahoe. *California Fish and Game* 57:28-43.
- GLEASON, J. S. 2007. Mallards feeding on salmon carcasses in Alaska. *Wilson Journal of Ornithology* 119:105-107.
- HEITMEYER, M. E. 2006. The importance of winter foods to mallards in the Mississippi Alluvial Valley. *Journal of Wildlife Management* 70:101-110.
- KOZLIK, F. M. 1974. Waterfowl of California. Sixth edition. California Department of Fish and Game, Sacramento, USA.
- KRAPU, G. L. 1981. The role of nutrient reserves in Mallard reproduction. *Auk* 98:29-38.
- LAFFERTY, K. D., J. P. McLAUGHLIN, AND J. E. DUGAN. 2013. Novel foraging in the swash zone on Pacific sand crabs (*Emerita analoga*) by mallards. *Wilson Journal of Ornithology* 125:423-426.
- MARTIN, A. C., H. S. ZIM, AND A. L. NELSON. 1961. American wildlife and plants, a guide to wildlife food habits. Dover Publications, New York, USA.
- MUNRO, J. A. 1923. A preliminary report on the relation of various ducks and gulls to the propagation of sock-eye salmon at Henderson Lake, Vancouver Island, B.C. *Canadian Field Naturalist* 37:107-116.
- MUNRO, J. A. 1943. Studies of waterfowl in British Columbia: mallard. *Canadian Journal of Research* 21:223-260.
- SWANSON, G. A., M. I. MEYER, AND V. A. ADOMAITIS. 1985. Foods consumed by breeding mallards on wetlands of south-central North Dakota. *Journal of Wildlife Management* 49:197-203.
- WILLSON, M. F., AND K. C. HALUPKA. 1995. Anadromous fish as keystone species in vertebrate communities. *Conservation Biology* 9:489-497.

Received 13 July 2014

Accepted 9 September 2014

Corresponding Editor was V. Bleich