

Project Overview

The Ocean Salmon Project (OSP), based in Santa Rosa, provides fisherydependent and fishery-independent data needed for the management of ocean, river, and tribal salmon fisheries. OSP produces annual estimates of ocean salmon harvest, fishery effort, ocean abundances, and fishery impacts on salmon stocks of special concern (ESA-listed stocks) and those managed by the Pacific Fishery Management Council and the California Fish and Game Commission. These stocks include fall Chinook (Oncorhynchus tshawytscha) from the Klamath-Trinity basin, Central Valley, and coastal rivers; spring and winter Chinook from the Central Valley; and coho (O. kisutch) statewide. Project biologists collaborate with other department, state, federal, and tribal salmon projects and provide technical assistance to the Pacific States Marine Fisheries Commission, the Pacific Fishery Management Council, the California Fish and Game Commission, non-governmental organizations, academia, and California constituents.

Ocean Fisheries Monitoring

- OSP has two primary objectives: 1) estimate total salmon harvest and fishing effort by management area and half month period for commercial and recreational ocean salmon fisheries, and 2) assess the contributions of important salmon stocks to these fisheries by time and management area
- A minimum of 20% of all ocean salmon fisheries are dockside sampled at major fishing ports all along the coast
- Commercial: $\geq 20\%$ of all salmon pounds by time and management area
- Recreational: \geq 20% of all salmon-targeting trips (charter and party boats) or available fishing days (private skiffs) by time and management area
- OSP field staff, in conjunction with California Recreational Fisheries Survey (CRFS) staff, measure and collect the heads from all adipose fin-clipped salmon (indicating the presence of a coded-wire tag) observed during sampling, and also collect other biological data, including fin clips for genetic analysis and scales for untagged-stock age structure, as needed













Ocean Salmon Project

California Department of Fish & Wildlife, Marine Region

Central Valley Support

- Project staff assist sampling returning spawners at several Central Valley hatcheries • Project staff collect scale samples from returning Chinook for age structure analyses and heads are collected from all adipose fin-clipped salmon for coded-wire
- tag recovery and scale-age validation
- During years of low Chinook abundance, project staff conduct real-time coded-wire tag analyses of returning spawners for broodstock selection in order to preserve the genetic integrity of different stocks and increase the available broodstock



Coded-Wire Tag Processing and Analysis

- In 2006, a Constant Fractional Marking program was initiated to increase the coded-wire tag rate of hatchery produced fall Chinook and tag each release group representatively; at least 25% of all fall Chinook produced in California's hatcheries are adipose fin-clipped and implanted with a microscopic ($\leq 1 \text{ mm}$) coded-wire tag which includes a unique code that provides release information
- OSP's laboratory now manually extracts, reads under a microscope, and validates coded-wire tags from over 30,000 ocean-caught adipose fin-clipped salmon and nearly 20,000 Central Valley Chinook annually
- An additional 20,000 Central Valley coded-wire tags are processed annually at a separate state lab with project staff providing data validation, management, and analysis
- Coded-wire tag recovery information, combined with harvest and effort estimates by time, management area, and fishery, provide biologists and fishery managers with the information needed to design fishing seasons that allow harvest of abundant stocks while protecting weaker stocks
- Project biologists determine the contribution and stray rates of hatchery and natural-origin Chinook returning to spawn in the Central Valley and Klamath-Trinity basins, and evaluate hatchery release strategies
- Each year, project staff upload statewide ocean and inland coded-wire tag recovery and associated catch-sample data to the Regional Mark Processing Center, a coastwide centralized public data warehouse maintained by the Pacific States Marine Fisheries Commission













Central Valley Scale Aging

- spawning escapement
- stocks
- An annual standardized Central Valley scale collection program was initiated in 2006 to allow for age-specific spawning escapement estimates by stock
- Central Valley scale samples are aged and validated with coded-wire tag information using state of the art digital imaging techniques along with computer aided data storage and retrieval methods



- Total and age-specific ocean and river harvest, fishery effort, spawning escapement, and associated coded-wire tag data collected and maintained by project staff provide the scientific basis for fishery management decisions
- Project staff use data to evaluate salmon cohort reconstructions, forecast Klamath River and Sacramento River fall Chinook ocean abundances, and update multi-year ocean harvest models which are used to analyze fishing season alternatives proposed by the fishing industry
- Project staff participate on Klamath-Trinity and Central Valley technical teams and the Pacific Fishery Management Council's Salmon Technical Team, which are responsible for ensuring the best available science is used for management decisions and that conservation objectives for all California salmon stocks are met
- Project staff host and actively participate in at least four annual public hearings to assist in the development of salmon fishing season alternatives and allow for constituent comment





- Melodie Palmer-Zwahlen **Brett Kormos** Jennifer Simon James Phillips **Barry Miller Alex Letvin**
 - Kandice Morgenstern Vanessa Gusman **Roxanne Jordan Adam Stewart** Katie Sowul

• Age-specific escapement data allow for cohort reconstruction which may result in more reliable forecasts of ocean abundance, fishery harvest, and expected

Age-specific data for Klamath-Trinity fall Chinook have been available for decades, but lacking for Central Valley Chinook, limiting scientific understanding of these



Fisheries Management

Data and analyses conducted by project staff are publicly available through the Pacific Fishery Management Council and the Regional Mark Processing Center